

# Experimental Access to the ⚠ Emergence of Mass ⚠

Ralf W. Gothe for the CLAS Collaboration


UNIVERSITY OF  
SOUTH CAROLINA




Emergent Mass and its Consequences in the Standard Model,  
September 17-21, 2018, ECT\*, Trento, Italy

- **$\gamma_v$ NN\* Experiments:** The best access to the baryon and quark structure?
- **Analysis and New Results:** Exclusive, quasi-free, and final state interaction!
- **Outlook:** New experiments with extended scope and kinematics!


# From Meson-Baryon to Quark Degrees of Freedom




- Study of the distance dependent structure of baryons into the domain where dressed quarks are the dominant active degree of freedom.
- Explore the formation of excited nucleon states in interactions of dressed quarks and their emergence from QCD.



# Hadron Structure with Electromagnetic Probes



- Study the structure of the nucleon spectrum in the domain where dressed quarks are the major active degree of freedom.
- Explore the formation of excited nucleon states in interactions of dressed quarks and their emergence from QCD.




# Spectroscopy

# Build your Mesons and Baryons ...

Three Generations of Matter (Fermions)


	I	II	III	
mass→	2.4 MeV	1.27 GeV	171.2 GeV	0
charge→	$\frac{2}{3}$	$\frac{2}{3}$	$\frac{2}{3}$	0
spin→	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
name→	<b>u</b> up	<b>c</b> charm	<b>t</b> top	<b>γ</b> photon
Quarks	4.8 MeV	104 MeV	4.2 GeV	0
	$-\frac{1}{3}$	$-\frac{1}{3}$	$-\frac{1}{3}$	0
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
	<b>d</b> down	<b>s</b> strange	<b>b</b> bottom	<b>g</b> gluon
Leptons	<2.2 eV	<0.17 MeV	<15.5 MeV	91.2 GeV
	0	0	0	0
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
	<b>ν<sub>e</sub></b> electron neutrino	<b>ν<sub>μ</sub></b> muon neutrino	<b>ν<sub>τ</sub></b> tau neutrino	<b>Z<sup>0</sup></b> weak force
	0.511 MeV	105.7 MeV	1.777 GeV	80.4 GeV
	-1	-1	-1	$\pm 1$
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
	<b>e</b> electron	<b>μ</b> muon	<b>τ</b> tau	<b>W<sup>±</sup></b> weak force



$$\mathcal{L} = \frac{1}{4g^2} G_{\mu\nu}^a G_{\mu\nu}^a + \sum_j \bar{q}_j (i\gamma^\mu D_\mu + m_j) q_j$$

where  $G_{\mu\nu}^a \equiv \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + if_{abc} A_\mu^b A_\nu^c$   
and  $D_\mu \equiv \partial_\mu + it^a A_\mu^a$   
*That's it?*

Frank Wilczek, Physics Today, August 2000




Bosons (Forces)




# N and $\Delta$ Excited Baryon States ...

Simon Capstick

➤ Orbital excitations  
(two distinct kinds in contrast to mesons)



➤ Radial excitations  
(also two kinds in contrast to mesons)



# Quark Model Classification of N\*


Dietmar Menze

Lowest Baryon Supermultiplets

SU(6)xO(3) Symmetry


Particle Data Group

- \*\*\*\*
- \*\*\*
- \*\*



# $N^*$ Spectrum in LQCD

The strong interaction physics is encoded in the nucleon excitation spectrum that spans the degrees of freedom from meson-baryon and dressed quarks to elementary quarks and gluons.





LQCD predicts states with the same quantum numbers as CQMs with underlying  $SU(6) \times O(3)$  symmetry.

R. Edwards et al.  
arXiv:1104.5152, 1201.2349

# Evolution of the Early Universe

Volker Burkert




Dramatic events occur in the microsecond old Universe

- Transition from the QGP to the baryon phase is dominated by excited baryons.
- A quantitative description requires more states than found to date → missing baryons.
- During the transition the quarks acquire dynamical mass and become confinement.

# Quark Model Classification of N\*


BnGa energy-dependent coupled-channel PWA of CLAS  $K^+\Lambda$  and other data




# Electron Scattering



# Baryon Excitations and Quasi-Elastic Scattering




Paticle and Nuclei, Povh et al., MAMI B





Deep Inelastic Scattering  
 S. Stein et al., PR **D22** (1975) 1884

# Baryon Excitations and Quasi-Elastic Scattering




quasi-elastic off  
point-like  
constituents





Deep Inelastic Scattering  
M. Breidenbach et al.,  
Phys. Rev. Lett. **23** (1969) 935


# Baryon Excitations and Quasi-Elastic Scattering





hard and confined



quasi-elastic



hard



soft


Deep Inelastic Scattering  
S. Stein et al., PR **D22** (1975) 1884

# Structure Analysis of the Baryon


Demolition of a chimney at the "Henninger Brewery" in Frankfurt am Main, Germany, on 2 December 2006




hard and confined



hard and soft




quasi-elastic




# Transition Form Factors

# Hadron Structure with Electromagnetic Probes




- Study the structure of the nucleon spectrum in the domain where dressed quarks are the major active degree of freedom.
- Explore the formation of excited nucleon states in interactions of dressed quarks and their emergence from QCD.




# Data-Driven Data Analyses

## Consistent Results

Single Pion




Int. J. Mod. Phys. E, Vol. 22, 1330015 (2013) 1-99




# Data-Driven Data Analyses

## Consistent Results

Double Pion



Int. J. Mod. Phys. E, Vol. 22, 1330015 (2013) 1-99




# Data-Driven Data Analyses

## Consistent Results





- Single meson production:  
Unitary Isobar Model (UIM)  
Fixed- $t$  Dispersion Relations (DR)
- Double pion production:  
Unitarized Isobar Model (JM)
- Coupled-Channel Approaches:  
EBAC  $\Rightarrow$  Argonne-Osaka  
JAW  $\Rightarrow$  Jülich-Athens-Washington  $\Rightarrow$  JüBo  
BoGa  $\Rightarrow$  Bonn-Gatchina

Int. J. Mod. Phys. E, Vol. 22, 1330015 (2013) 1-99



# Electrocouplings of $N(1440)P_{11}$ from CLAS Data




□ PDG   
 ●  $N\pi$  (UIM, DR)   
 ▲  $N\pi\pi$  (JM) 2012   
 ■  $N\pi\pi$  (JM) preliminary

Consistent results obtained in the low-lying resonance region by independent analyses in the exclusive  $N\pi$  and  $p\pi^+\pi^-$  final-state channels – that have fundamentally different mechanisms for the nonresonant background – underscore the capability of the reaction models to extract reliable resonance electrocouplings.

Phys. Rev. C 80, 055203 (2009) 1-22 and Phys. Rev. C 86, 035203 (2012) 1-22

# Electrocouplings of $N(1440)P_{11}$ History




- Lowest mass hybrid baryon should be  $J^P=1/2^+$  as Roper.
- In 2002 Roper  $A_{1/2}$  results were consistent with a hybrid state.

# Transition Form Factors and QCD Models

Roper resonance  $P_{11}(1440)$

PDG 2013 update



+  $q^3g$   
 +  $q^3q\bar{q}$   
 + N-Meson  
 + ...

or

-  $q^2q$   
 - ...


... all have distinctively different  $Q^2$  dependencies

- $A_{1/2}$  has zero-crossing near  $Q^2=0.5$  and becomes dominant amplitude at high  $Q^2$ .
- Consistent with radial excitation at high  $Q^2$  and large meson-baryon coupling at small  $Q^2$ .
- Eliminates gluonic excitation ( $q^3G$ ) as a dominant contribution.


Nick Tyler closes the 1-2  $GeV^2$  gap for single pion production.

# Transition Form Factors and QCD Models

Roper resonance  $P_{11}(1440)$



I.T. Obukhovsky





- $A_{1/2}$  has zero-crossing near  $Q^2=0.5$  and becomes dominant amplitude at high  $Q^2$ .
- Consistent with radial excitation at high  $Q^2$  and large meson-baryon coupling at small  $Q^2$ .
- Eliminates gluonic excitation ( $q^3G$ ) as a dominant contribution.

Nick Tyler closes the **1-2 GeV<sup>2</sup> gap** for single pion production.

# Transition Form Factors and QCD Models

Roper resonance  $P_{11}(1440)$

G. Ramalho





- $A_{1/2}$  has zero-crossing near  $Q^2=0.5$  and becomes dominant amplitude at high  $Q^2$ .
- Consistent with radial excitation at high  $Q^2$  and large meson-baryon coupling at small  $Q^2$ .
- Eliminates gluonic excitation ( $q^3G$ ) as a dominant contribution.

Nick Tyler closes the **1-2 GeV<sup>2</sup> gap** for single pion production.

# Roper Transition Form Factors in DSE Approach


**N(1440)P<sub>11</sub>**

J. Segovia *et al.*, Phys. Rev. Lett. **115**, 171801



- DSE Contact       $x=Q^2/m_N^2$
- DSE Realistic
- Inferred meson-cloud contribution
- Anticipated complete result



Importantly, the existence of a zero in  $F_2$  is not influenced by meson-cloud effects, although its precise location is.



# Roper Transition Form Factors in DSE Approach

$N(1440)P_{11}$


J. Segovia and C.D. Roberts, arXiv:1607.04405



# DSE and EBAC/ANL-Osaka Approaches

... more  $(\pi, \pi\pi)$ ,  $(\pi, \pi\eta)$ , and  $(\pi, KY)$  data needed

Semi-quantitative agreement with the first DSE 3-dressed-quark core radial excitation.




Int. J. Mod. Phys. E, Vol. 22, 1330015 (2013)

# Structure of the Roper Resonance from LQCD

Hamiltonian effective field theory approach based on an extension of chiral perturbation theory that incorporates the Lüscher relation connecting the energy levels observed in finite volume to the scattering phase shifts.


Input: fit to experimental phase shift, inelasticity, and T-matrix data



Adelaide and Graz Groups




# Structure of the Roper Resonance from LQCD

Hamiltonian effective field theory approach based on an extension of chiral perturbation theory that incorporates the Lüscher relation connecting the energy levels observed in finite volume to the scattering phase shifts.



Adelaide and Graz Groups

# Electrocouplings of $N(1520)D_{13}$ and $N(1535)S_{11}$



— Argonne Osaka / EBAC DCC MB dressing  
(absolute values)

— E. Santopinto, M. Giannini, hCQM  
PRC 86, 065202 (2012)


— S. Capstick, B.D. Keister (rCQM)  
PRD51, 3598 (1995)

■  $\pi^+\pi^-p$  2012    ▲  $\pi^+\pi^-p$  2010    ●  $N\pi$  2009


▲  $\eta p$   
CLAS/Hall-C

# Interplay between Meson-Baryon Cloud and Quark Core

$N(1520)3/2^-$



$N(1675)5/2^-$



..... Argonne-Osaka MB dressing (absolute values)


— E. Santopinto and M. Giannini, PRC 86 (2012) 065202

The almost direct access to

- quark core from the data on  $N(1520)3/2^-$
- meson-baryon cloud from the data on  $N(1675)5/2^-$

sheds light on the transition from the confined quark to the colorless meson-baryon structure and its dependents on the  $N^*$  quantum numbers.


# Evidence for the Onset of Precocious Scaling?



➤  $A_{1/2} \propto 1/Q^3$


➤  $A_{3/2} \propto 1/Q^5$

I. G. Aznauryan *et al.*, Phys. Rev. C80, 055203 (2009)



# Evidence for the Onset of Precocious Scaling?


Ye Tian



V. Mokeev, [userweb.jlab.org/~mokeev/resonance\\_electrocouplings/](http://userweb.jlab.org/~mokeev/resonance_electrocouplings/) (2016)

# Evidence for the Onset of Precocious Scaling?

Ye Tian




V. Mokeev, [userweb.jlab.org/~mokeev/resonance\\_electrocouplings/](http://userweb.jlab.org/~mokeev/resonance_electrocouplings/) (2016)

# N(1520)D<sub>13</sub> Helicity Asymmetry


L. Tiator

$$A_{\text{hel}} = \frac{A_{1/2}^2 - A_{3/2}^2}{A_{1/2}^2 + A_{3/2}^2}$$



# $\gamma NN^*$ Helicity Asymmetries

Ye Tian




V. Mokeev, [userweb.jlab.org/~mokeev/resonance\\_electrocouplings/](http://userweb.jlab.org/~mokeev/resonance_electrocouplings/) (2016)

# New Experimental Results & Approaches


# Single $\pi^-$ Electroproduction off the Deuteron

Ye Tian



# Single $\pi^-$ Electroproduction off the Deuteron


Ye Tian



Below a missing momentum of 0.2 GeV the **measured data** coincides with the resolution smeared **theoretical Fermi momentum distribution**.

# Single $\pi^-$ Electroproduction off the Deuteron

Ye Tian



Gary Hollis inclusive of the **bound nucleon** in the Deuteron with correction of Fermi smearing.

# Single $\pi^-$ Electroproduction off the Deuteron

Ye Tian

$W = 1212 \text{ MeV}$

$\Delta W = 25 \text{ MeV}$

$Q^2 = 0.5 \text{ GeV}^2$

$\Delta Q^2 = 0.2 \text{ GeV}^2$

$\cos(\theta) = -0.7$











$\Delta \cos(\theta) = 0.2$

$\cos(\theta) = 0.7$

$\phi = 20^\circ$










$\Delta \phi = 40^\circ$

$\phi = 340^\circ$



# Single $\pi^-$ Electroproduction off the Deuteron

Ye Tian



# Single $\pi^-$ Electroproduction off the Deuteron

Ye Tian



Inclusive:  
Gary Hollis

$Q^2 = 0.5 \text{ GeV}^2$

$W = 1212 \text{ MeV}$

$Q^2 = 0.7 \text{ GeV}^2$

$Q^2 = 0.9 \text{ GeV}^2$




# Single $\pi^-$ Electroproduction off the Deuteron

$Q^2 = 0.5 \text{ GeV}^2$

$W = 1212 \text{ MeV}$


$W = 1488 \text{ MeV}$



Ye Tian

Inclusive:  
Gary Hollis



$W = 1662 \text{ MeV}$



# $N\pi^+\pi^-$ Electroproduction Kinematic Coverage

Gleb Fedotov


Phys. Rev. C 98, 025203 (2018)



$\pi\pi^+\pi^-$  event yields over  $W$  and  $Q^2$ . Gray shaded area new  $e1e$  data set, hatched area at low  $Q^2$  already published  $e1c$  data by G. Fedotov *et al.* and hatched area at higher  $Q^2$  already published data in one large  $Q^2$  bin by M. Ripani *et al.*

# Integrated $N\pi^+\pi^-$ Cross Sections

Gleb Fedotov




Bound Proton:  
I. Skorodumina

Black hatched already published data (Fedotov *et al.*, PRC79, 015204 (2009)) and red hatched new  $e1e$  data in the overlap region.

# $N\pi^+\pi^-$ Single-Differential Cross Sections


Gleb Fedotov



# Exclusive $\pi^+\pi^-$ Electroproduction off the Deuteron

Iuliia Skorodumina

$P_X$  of  $ep(n) \rightarrow e'p'(n)\pi^+\pi^-$







# Effective FSI Correction in $p(n)\pi^+ \pi^-$

Iuliia Skorodumina

$$\frac{d\sigma_{corrected}}{dW dQ^2 d\tau} = \frac{d\sigma_{not\ corrected}}{dW dQ^2 d\tau} F_{fsi}(\Delta W, \Delta Q^2)$$



$$F_{fsi}(\Delta W, \Delta Q^2) = \frac{\text{Area under green}}{\text{Area under red}}$$



Preliminary

# Comparison with Free Proton Cross Section

Iulia




**Black bullets** – free proton cross sections ( $e1e$  at  $E_{\text{beam}} = 2.039 \text{ GeV}$ )  
 error bars show both statistical and systematical uncertainties  
 G. Fedotov under paper review

**Red bullets** – bound proton quasi-free cross sections ( $e1e$  at  $E_{\text{beam}} = 2.039 \text{ GeV}$ )  
 error bars show statistical uncertainty only

# $\phi$ -dependent $N\pi\pi$ Single-Differential Cross Sections

$Q^2, W$  bin =  $[2.4, 3.0) \text{ GeV}^2, [1.725, 1.750) \text{ GeV}$

Arjun Trivedi  
Evgeny Isupov



● normalized

● hole filled


● TWOPEG

$$\left( \frac{d^2\sigma}{dX_{ij}d\phi_i} \right) = \underline{R2_T X_{ij} + R2_L X_{ij}} + R2_{LT}^{c, X_{ij}} \cos \phi_i + R2_{TT}^{c, X_{ij}} \cos 2\phi_i + \delta_{X_{ij}\alpha_i} (R2_{LT}^{s, \alpha_i} \sin \phi_i + R2_{TT}^{s, \alpha_i} \sin 2\phi_i)$$

# $\phi$ -dependent $N\pi\pi$ Single-Differential Cross Sections

$Q^2, W$  bin =  $[2.4, 3.0)\text{GeV}^2, [1.725, 1.750)\text{GeV}$

Arjun Trivedi



$$\left(\frac{d^2\sigma}{dX_{ij}d\phi_i}\right) = R2_T^{X_{ij}} + R2_L^{X_{ij}} + \underline{R2_{LT}^{c,X_{ij}} \cos \phi_i} + \underline{R2_{TT}^{c,X_{ij}} \cos 2\phi_i} + \delta_{X_{ij}\alpha_i} (R2_{LT}^{s,\alpha_i} \sin \phi_i + R2_{TT}^{s,\alpha_i} \sin 2\phi_i)$$


# $\phi$ -dependent $N\pi\pi$ Single-Differential Cross Sections

$Q^2, W$  bin =  $[2.4, 3.0) \text{ GeV}^2, [1.725, 1.750) \text{ GeV}$

Arjun Trivedi

Chris McLauchlin extracts the **beam helicity dependent** differential cross sections.

Preliminary




$$\left( \frac{d^2\sigma}{dX_{ij}d\phi_i} \right) = R2_T^{X_{ij}} + R2_L^{X_{ij}} + R2_{LT}^{c, X_{ij}} \cos \phi_i + R2_{TT}^{c, X_{ij}} \cos 2\phi_i + \delta_{X_{ij}\alpha_i} \left( \underline{R2_{LT}^{s, \alpha_i} \sin \phi_i} + \underline{R2_{TT}^{s, \alpha_i} \sin 2\phi_i} \right)$$


# CLAS12



# 12 GeV CEBAF






- Luminosity  $> 10^{35} \text{ cm}^{-2}\text{s}^{-1}$
- Hermeticity
- Polarization
  
- Baryon Spectroscopy
- Elastic Form Factors
- N to N\* Form Factors
- GPDs and TMDs
- DIS and SIDIS
- Nucleon Spin Structure
- Color Transparency
- ...




# New Forward Time of Flight Detector for CLAS12

ToF12 Time Resolution Measurements





# Anticipated $N^*$ Electrocouplings from Combined Analyses of $N\pi/N\pi\pi$





Open circles represent projections and all other markers the available results with the 6-GeV electron beam

- Examples of **published and projected results** obtained within 60d for three prominent excited proton states from analyses of  $N\pi$  and  $N\pi\pi$  electroproduction channels. Similar results are expected for many other resonances at higher masses, e.g.  $S_{11}(1650)$ ,  $F_{15}(1685)$ ,  $D_{33}(1700)$ ,  $P_{13}(1720)$ , ...
- The approved CLAS12 experiments E12-09-003 (NM,  $N\pi\pi$ ) and E12-06-108A (KY) are currently **the only experiments** that can provide data on  $\gamma_v NN^*$  electrocouplings for almost all well established excited proton states at the highest photon virtualities ever achieved in  $N^*$  studies up to  $Q^2$  of 12  $\text{GeV}^2$ , see <http://boson.physics.sc.edu/~gothe/research/pub/whitepaper-9-14.pdf>.

# Summary

- First high precision photo- and electroproduction data have become available and led to a new wave of significant developments in reaction and QCD-based theories.
- New high precision hadro-, photo-, and electroproduction data off the proton and the neutron will stabilize coupled channel analyses and expand the validity of reaction models, allowing us to
  - investigate and search for baryon hybrids (E12-16-010) ,
  - establish a repertoire of high precision spectroscopy parameters, and
  - measure light-quark-flavor separated electrocouplings over an extended  $Q^2$ -range, both to lower and higher  $Q^2$ , for a wide variety of  $N^*$  states (E12-16-010 A).
- Comparing these results with LQCD, DSE, LCSR, and rCQM will build further insights into
  - the strong interaction of dressed quarks and their confinement,
  - the origin of 98% of nucleon mass, and
  - the emergence of bare quark dressing and dressed quark interactions from QCD.
- A close collaboration of experimentalists and theorists has formed, is growing, and is needed to push these goals, see Review Article *Int. J. Mod. Phys. E*, Vol. 22, 1330015 (2013) 1-99, that shall lead to a strong QCD theory that describes the strong interaction from current quarks to nuclei.



**ECT\*2015, INT2016, NSTAR2017, APCTP2018 ...**