


PAULI EXCLUSION PRINCIPLE VIOLATION AND QUANTUM GRAVITY WITH VIP-2

*Fabrizio Napolitano on behalf of the
VIPCollaboration*



fabrizio.napolitano@lnf.infn.it

Nuclear and Atomic Transitions as Laboratories for High Precision


Tests of Quantum Gravity Inspired Models - ECT Trento 20/09/2022*

The Pauli Exclusion Principle

In an atom there cannot be two or more equivalent electrons for which the values of all four quantum numbers coincide. If an electron exists in an atom for which all of these numbers have definite values, then the state is occupied.

W. Pauli, Über den Zusammenhang des Abschlusses der Elektronengruppen im Atom mit der Komplexstruktur der Spektren, Zeitschrift für Physik 31 (1925) 765.

Pauli Archive, holding: fierz_0092-064




The Pauli Exclusion Principle

In an atom there cannot be two or more equivalent electrons for which the values of all four quantum numbers coincide. If an electron exists in an atom for which all of these numbers have definite values, then the state is occupied.

W. Pauli, Über den Zusammenhang des Abschlusses der Elektronengruppen im Atom mit der Komplexstruktur der Spektren, Zeitschrift für Physik 31 (1925) 765.

Pauli Archive, holding: fierz_0092-064



.. The impression that the shadow of some incompleteness [falls] here on the bright light of success of the new quantum mechanics seems to me unavoidable.

W. Pauli, Nobel lecture 1945

The Pauli Exclusion Principle (PEP)

Spin-statistic connection:

half-integer spin particles → antisymmetric wave function & Fermi-Dirac stat

Integer spin particles → symmetric wave function & Bose statistics

Lüders and Zumino: spin-statistics lays on few, general assumption:


Lorentz/Poincaré Symmetry, CPT, unitarity, locality & causality

Theories of Statistics Violation


O.W. Greenberg: AIP Conf.Proc.545:113-127,2004

“Possible external motivations for violation of statistics include: (a) violation of CPT, (b) violation of locality, (c) violation of Lorentz invariance, (d) extra space dimensions, (e) discrete space and/or time and (f) non-commutative spacetime.....”

The Pauli Exclusion Principle (PEP)



The Pauli Exclusion Principle (PEP)



BSM theories embedding extra dimensions, non commutative and/or discrete spacetime could have effect on PEP

How to model PEP violations

- *Ignatiev & Kuzmin model: Fermi oscillator with a third state*

(Ignatiev, A.Y., Kuzmin, V. , *Quarks '86: Proceedings of the 229 Seminar, Tbilisi, USSR, 1517 April 1986*)

$$\begin{array}{ll}
 a^+|0\rangle = |1\rangle & a|0\rangle = 0 \\
 a^+|1\rangle = \beta|2\rangle & a|1\rangle = |0\rangle \\
 a^+|2\rangle = 0 & a|2\rangle = \beta|1\rangle
 \end{array}$$

β quantifies the degree of violation in the transition

- *Greenberg & Mohapatra: Local Quantum Field Theory, q parameter deforms anticommutators [Phys. Rev. Lett. 1987,59,2507]:*

$$a_k a^+{}_l - q a^+{}_l a_k = \delta_{k,l}$$


- *Rahal & Campa: global wave function of the electrons not exactly antisymmetric, PEP holds as long as the number of wrongly entangled pairs is small*

All respect the Messiah-Greenberg super-selection rule!

Messiah-Greenberg super-selection rule:


Superposition of states with different symmetry are not allowed \rightarrow

Transition probability between two symmetry states is ZERO




VIP-2 Experiment: best limits on PEP violation of an elementary particle respecting the Messiah-Greenberg super-selection rule

Search for anomalous X-ray transitions performed by electrons introduced in a target *through a DC current (open system)*



Normal $2p \rightarrow 1s$ transition

~ 8.05 keV in Cu



$2p \rightarrow 1s$ transition violating Pauli principle

~ 7.7 keV in Cu

Paul Indelicato (Ecole Normale Supérieure et Université Pierre et Marie Curie)


Multiconfiguration Dirac-Fock approach

Accounts for the shielding of the two inner electrons


**Greenberg, O. W. & Mohapatra, R. N., Phys Rev Lett 59, (1987).
E. Ramberg and G. A. Snow, Phys Lett B 238, 438-441(1990)**

**Search for anomalous electronic transitions in Cu
induced by a circulating current**

**introduced electrons interact with the valence electrons
search transition from 2p to 1s already filled by 2 electrons
alternated to X-ray background measurements without current**





PEP Violation Signal:



From VIP to VIP-2

- copper ultrapure cylindrical foil
- surrounded by 16 Charge Coupled Devices (CCD) res. at 8 keV 320 eV (FWHM)
- inside a vacuum chamber: CCDs cooled to 168K by a cryogenic system
- amplifiers + read out ADC boards.



$$\beta^2/2 \leq 4.7 \times 10^{-29}$$





*improved the limit obtained by Ramberg & Snow
by a factor ~ 400*

(Foundation of Physics 41 (2011) 282+ other papers)

GOAL OF VIP-2 : improve the VIP result of 2 orders of magnitude

The VIP-2 Experiment

Silicon Drift Detectors (SDDs) higher resolution (190 eV FWHM at 8.0 \rightarrow keV), faster (triggerable) detectors. 4 arrays of 2 x 4 SDDs 8mm x 8mm each, liquid argon closed circuit cooling 170 °C





The VIP-2 Experiment


2 strip shaped Cu targets (25 μm x 7 cm x 2 cm) more compact target \rightarrow higher acceptance, thinner \rightarrow higher efficiency

DC current supply to Cu bars


Cu strips cooled by a closed Fryka chiller circuit \rightarrow higher current (100 A) @ 20 °C of Cu target implies 1 °K heating in SDDs

Sketch of the VIP2 Setup:







Upgrade concluded in April 2019:




1400 m rock coverage

Passive shielding → two layers, copper inside lead outside



Results of six months of data taking<https://doi.org/10.3390/sym14050893>

Symmetry 2022, 14(5), 893;

Results of six months of data taking<https://doi.org/10.3390/sym14050893>

Symmetry 2022, 14(5), 893;

Description spectrum with current

$$\mathcal{F}^{wc}(\theta, \mathbf{y}, \mathcal{S}) = y_1 \times \underline{\text{Ni}(\theta_1, \theta_2)} + y_2 \times \underline{\text{Cu}(\theta_3, \theta_4)} + y_3 \times \text{pol}_1(\theta_5) + \mathcal{S} \times \underline{\text{PEPV}(\theta_4)}$$

Description spectrum without current

$$\mathcal{F}^{woc}(\theta, \mathbf{y}) = y_1 \times \underline{\text{Ni}(\theta_1, \theta_2)} + y_2 \times \underline{\text{Cu}(\theta_3, \theta_4)} + y_3 \times \text{pol}_1(\theta_5)$$

Results of six months of data taking<https://doi.org/10.3390/sym14050893>

Likelihood

*Data with current**Data without current*

$$\mathcal{L}(\mathcal{D}^{wc}, \mathcal{D}^{woc} | \boldsymbol{\theta}, \mathbf{y}, \mathcal{S}) = \text{Poiss}(\mathcal{D}^{wc} | \mathcal{F}^{wc}(\boldsymbol{\theta}, \mathbf{y}, \mathcal{S})) \times \text{Poiss}(\mathcal{D}^{woc} | \mathcal{F}^{woc}(\boldsymbol{\theta}, \mathbf{y} \times \mathcal{R}))$$



Bayesian

Frequentist

$$p(\boldsymbol{\theta}, \mathbf{y}, \mathcal{S} | \mathcal{D}^{wc}, \mathcal{D}^{woc}) =$$

$$= \frac{\mathcal{L}(\mathcal{D}^{wc}, \mathcal{D}^{woc} | \boldsymbol{\theta}, \mathbf{y}, \mathcal{S}) p(\boldsymbol{\theta}, \mathbf{y}, \mathcal{S})}{\int d\boldsymbol{\theta} d\mathbf{y} \mathcal{L}(\mathcal{D}^{wc}, \mathcal{D}^{woc} | \boldsymbol{\theta}, \mathbf{y}, \mathcal{S}) p(\boldsymbol{\theta}, \mathbf{y}, \mathcal{S})}$$

$$t_{\mathcal{S}} = -2 \ln \Lambda(\mathcal{S}) = -2 \ln \frac{\mathcal{L}(\hat{\boldsymbol{\theta}}, \hat{\mathbf{y}}, \mathcal{S})}{\mathcal{L}(\hat{\boldsymbol{\theta}}, \hat{\mathbf{y}}, \hat{\mathcal{S}})}$$

$$p_{\mathcal{S}} = \int_{t_{\text{obs}}}^{\infty} f(t_{\mathcal{S}} | \mathcal{S}) dt_{\mathcal{S}}$$

Priors: gaussian distributions for parameters constrained within experimental uncertainties; flat distribution for signal

Experimental uncertainty (e.g. energy scale) included via additional penalty terms in Likelihood



Results of six months of data taking<https://doi.org/10.3390/sym14050893>

Likelihood

*Data with current**Data without current*

$$\mathcal{L}(\mathcal{D}^{wc}, \mathcal{D}^{woc} | \theta, \mathbf{y}, \mathcal{S}) = \text{Poiss}(\mathcal{D}^{wc} | \mathcal{F}^{wc}(\theta, \mathbf{y}, \mathcal{S})) \times \text{Poiss}(\mathcal{D}^{woc} | \mathcal{F}^{woc}(\theta, \mathbf{y} \times \mathcal{R}))$$

Bayesian

Frequentist

$$p(\theta, \mathbf{y}, \mathcal{S} | \mathcal{D}^{wc}, \mathcal{D}^{woc}) =$$

$$= \frac{\mathcal{L}(\mathcal{D}^{wc}, \mathcal{D}^{woc} | \theta, \mathbf{y}, \mathcal{S}) p(\theta, \mathbf{y}, \mathcal{S})}{\int d\theta d\mathbf{y} \mathcal{L}(\mathcal{D}^{wc}, \mathcal{D}^{woc} | \theta, \mathbf{y}, \mathcal{S}) p(\theta, \mathbf{y}, \mathcal{S})}$$

See Alessio's talk Thursday

$$-2 \ln \Lambda(\mathcal{S}) = -2 \ln \frac{\mathcal{L}(\hat{\theta}, \hat{\mathbf{y}}, \mathcal{S})}{\mathcal{L}(\hat{\theta}, \hat{\mathbf{y}}, \hat{\mathcal{S}})}$$

$$p_{\mathcal{S}} = \int_{t_{\text{obs}}}^{\infty} f(t_{\mathcal{S}} | \mathcal{S}) dt_{\mathcal{S}}$$




Priors: gaussian distributions for parameters constrained within experimental uncertainties; flat distribution for signal

Experimental uncertainty (e.g. energy scale) included via additional penalty terms in Likelihood


Symmetry **2022**, *14*(5), 893;

<https://doi.org/10.3390/sym14050893>

Bayesian



Frequentist



Small under-fluctuation in the ROI

$$\beta^2/2 \leq 8.6 \times 10^{-31} \text{ (Bayesian)}, \quad \beta^2/2 \leq 8.9 \times 10^{-31} \text{ (CL}_s\text{)}$$

New article in preparation with all the available statistics!

VIP-2 experimental upgrade: VIP-3

Scan the PEP violation probability as a function of Z (i.e. of Energy)

Okun, L.:

“The special place enjoyed by the Pauli principle in modern theoretical physics does not mean that this principle does not require further and exhaustive experimental tests. On the contrary, it is specifically the fundamental nature of the Pauli principle which would make such tests, over the entire periodic table, of special interest”



**L. Possible violation of the Pauli principle in atoms. JETP Lett.
1987, 46, 529532**

**“High sensitivity Pauli Exclusion Principle tests by the VIP experiment:
status and perspectives”**

Paper on the preparation of VIP-3 experiment accepted in APPA.



VIP-2 experimental upgrade: VIP-3

- *new vacuum chamber, increase the number of SDD detectors, increase the geometrical efficiency, higher current up to 400 A*
- *New thermal contact between cold finger and SDDs*
- *New target cooling system*
- *Higher quantum efficiency needed for the SDDs at higher Z: use 1 mm thick SDDs, allowing to scan e.g. Ag, Sn and Pd*




VIP-2 experimental upgrade: VIP-3


- 2x4 SDDs, 8x8 mm² each, in production with FBK & politecnico di Milano
- Improved charge collection at border of the active area
- Wider than previous chip
- Reduced charge sharing via focusing electrode on the window



Anode side



Window side



New paradigm for VIP-2

Are Quantum Gravity models experimentally testable?


A. Addazi (Chengdu Univ.) A. Marcianò (Fudan University)

VIP-2 underground experiment as a *Crash-Test* of Non-Commutative Quantum Gravity

Pauli Exclusion Principle (PEP) violations induced from non-commutative space-time can be searched VIP-2 experiment set-up. We show that the limit from VIP-2 experiments on non-commutative space-time scale Λ , related to energy dependent PEP violations, are severe: κ -Poincaré non-commutativity is ruled-out up to the Planck scale. In the next future θ -Poincaré will be probed until the Grand-Unification scale! This highly motivates Pauli Exclusion Principle tests from underground experiments as a test of quantum gravity and space-time microscopic structure.

See also A. Addazi et al., 2018 Chinese Phys. C 42 094001, arXiv:1712.08082 [hep-th]

The Pauli Exclusion Principle (PEP)



BSM theories embedding extra dimensions, non commutative and/or discrete spacetime could have effect on PEP

PEP violation in quantum gravity

Quantum gravity models can embed PEP violating transitions

PEP is a consequence of the spin statistics theorem based on: Lorentz/Poincaré and CPT symmetries; locality; unitarity and causality. Deeply related to the very same nature of space and time



Non-commutativity of space-time is common to several quantum gravity frameworks (e.g. k -Poincaré, θ -Poincaré)



non-commutativity induces a deformation of the Lorentz symmetry and of the locality → naturally encodes the violation of PEP not constrained by MG

PEP violation is suppressed with $\delta^2 (E, \Lambda)$

E is the characteristic transition energy, Λ is the scale of the space-time non-commutativity emergence.

A. P. Balachandran, G. Mangano, A. Pinzul and S. Vaidya, Int. J. Mod. Phys. A 21 (2006) 3111

A.P. Balachandran, T.R. Govindarajan, G. Mangano, A. Pinzul, B.A. Qureshi and S. Vaidya, Phys. Rev. D 75 (2007)

A. Addazi, P. Belli, R. Bernabei and A. Marciano, Chin. Phys. C 42 (2018) no.9

Theoretical prediction *Int.J.Mod.Phys.A* 35 (2020) 32, 2042003

specific calculation of atomic levels transitions probabilities for θ -Poincaré

$$W \simeq W_0 \phi_{PEPV}, \quad \phi_{PEPV} = \delta^2 \simeq \frac{D E_N \Delta E}{2 \Lambda \Lambda} \quad \phi_{PEPV} = \delta^2 \simeq \frac{C \bar{E}_1 \bar{E}_2}{2 \Lambda \Lambda}$$

for non-vanishing (vanishing) electric like components of the $\theta_{\mu\nu}$ tensor.

Connection with quon algebra (in the case of quon fields however the q factor does not show any energy dependence):

$$q(E) = -1 + 2\delta^2(E)$$

An experimental bound on the probability that PEP may be violated in atomic transition processes, straightforwardly translates into a bound on the new physics scale Λ , consistently with the choice of the θ_{0i} components.

Experimental Setup

High purity Ge detector measurement:

- **high purity co-axial p-type germanium detector (HPGe), diameter of 8.0 cm, length of 8.0 cm, surrounded by an inactive layer of lithium-doped germanium of 0.075 mm.**
- **The target material is composed of three cylindrical sections of radio-pure Roman lead, completely surrounding the detector.**




Fig. 1 Schematic representation of the Ge crystal (in green) and the surrounding lead target cylindrical sections (in grey)

Experimental Setup

- **Passive shielding: inner - electrolytic copper, outer - lead**
- **10B-polyethylene plates reduce the neutron flux towards the detector**
- **shield + cryostat enclosed in air tight steel housing flushed with nitrogen to avoid contact with external air (and thus radon).**

K. Piscicchia et al., Eur. Phys. J. C (2020) 80: 508

<https://doi.org/10.1140/epjc/s10052-020-8040-5>






Figure 1: Schematic representation of the experimental setup: 1 - Ge crystal, 2 - Electric contact, 3 - Plastic insulator, 4 - Copper cup, 5 - Copper end-cup, 6 - Copper block and plate, 7 - Inner Copper shield, 8 - Lead shield.

Results



Background

Signal from violating transitions
in Lead



First analysis which accounts for the predicted energy dependence of the PEP violation probability. Expected rate of $K_{\alpha 1}$ transitions:

$$\Gamma_{K_{\alpha 1}} = \frac{\delta^2(E_{K_{\alpha 1}})}{\tau_{K_{\alpha 1}}} \cdot \frac{BR_{K_{\alpha 1}}}{BR_{K_{\alpha 1}} + BR_{K_{\alpha 2}}} \cdot 6 \cdot N_{atom} \cdot \epsilon(E_{K_{\alpha 1}}).$$

- upper limit on the non-commutativity scale

$$\mu = \sum_{K=1}^{N_K} \mu_K = \frac{N}{\Lambda^k} < \bar{S}$$

Results



From which an upper limit on the non-commutativity scale is obtained (90% Probability):

θ_{0i}	\bar{S}	lower limit on Λ (Planck scales)
$\theta_{0i} = 0$	13.2990	$6.9 \cdot 10^{-2}$
$\theta_{0i} \neq 0$	18.1515	$2.6 \cdot 10^2$

accepted PRL: “Strongest atomic physics bounds on Non-Commutative Quantum Gravity Models”

PHYSICAL REVIEW LETTERS

[Highlights](#) [Recent](#) [Accepted](#) [Collections](#) [Authors](#) [Referees](#) [Search](#) [Press](#) [About](#) [Editorial Team](#) 

Access by INFN/Laboratory Nazionali Di Frascati

Strongest Atomic Physics Bounds on Noncommutative Quantum Gravity Models

Kristian Piscicchia, Andrea Addazi, Antonino Marcianò, Massimiliano Bazzi, Michael Cargnelli, Alberto Clozza, Luca De Paolis, Raffaele Del Grande, Carlo Guaraldo, Mihail Antoniu Iliescu, Matthias Laubenstein, Johann Marton, Marco Miliucci, Fabrizio Napolitano, Alessio Porcelli, Alessandro Scordo, Diana Laura Sirghi, Florin Sirghi, Oton Vazquez Doce, Johann Zmeskal, and Catalina Curceanu

Phys. Rev. Lett. **129**, 131301 – Published 19 September 2022

[Article](#)[References](#)[No Citing Articles](#)[PDF](#)[HTML](#)[Export Citation](#)

ABSTRACT

Investigations of possible violations of the Pauli exclusion principle represent critical tests of the microscopic space-time structure and properties. Space-time noncommutativity provides a class of universality for several quantum gravity models. In this context the VIP-2 lead experiment sets the strongest bounds, searching for the Pauli exclusion principle violating atomic transitions in lead, excluding the θ -Poincaré noncommutative quantum gravity models far above the Planck scale for nonvanishing $\theta_{\mu\nu}$ electriclike components, and up to 6.9×10^{-2} Planck scales if $\theta_{0i} = 0$.

[Issue](#)[Vol. 129, Iss. 13
2022](#)[Check](#)

Conclusions

- *VIP-2 Experiment in data taking: pushing the limit on Pauli exclusion principle violations*
- *BSM scenario motivate precision tests on PEP*
- *VIP-2 set the strongest limit in the MG super selection rule*
- *Publication of full statistics data soon!*
- *Quantum gravity model predict PEP violation in closed systems*
 - *Analyzed with high purity germanium detectors*
 - *Obtained upper limit on non-commutativity scale*

Thank you for your attention!
Questions?

Proof of spin-statistics theorem by Lüders and Zumino

Postulates:

- *The theory is invariant with respect to the proper inhomogeneous Lorentz group (includes translations, does not include reflections)*
- *Two operators of the same field at points separated by a spacelike interval either commute or anticommute (locality – microcausality)*
- *The vacuum is the state of lowest energy*
- *The metric of the Hilbert space is positive definite*
- *The vacuum is not identically annihilated by a field*

From these postulates it follows that (pseudo)scalar fields commute and spinor fields anticommute.

(G. Lüders and B. Zumino, Phys. Rev. 110 (1958) 1450)

Models of Pauli Exclusion Principle (PEP) Violations


Some more PEP Violating models:

Greenberg, O.W. Mohapatra, R.N. Physical Review Letters 1987, 59, 2507
Govorkov, A. Physica A: Statistical Mechanics and its Applications 1994, 203, 655
Rahal, V.; Campa, A. , Physical Review A (1988) 38, 3728

Messiah - Greenberg superselection rule

Superpositions of states with different symmetry are not allowed →
 transition probability between two symmetry states is ZERO

Messiah-Greenberg superselection rule :



**VIP-open systems sets the best limit on PEP violation
 for an elementary particle
 respecting the M-G superselection rule**

VIP-2 experiment goal










(Upper limit not using Close Encounters (CE) treatment)

As reference for past experiments

Experiment	Target	Upper limit of $\beta^2/2$	reference
Ramberg-Snow	Copper	1.7×10^{-26}	[5]
S.R. Elliott et al.	Lead	1.5×10^{-27}	[14]
VIP(2006)	Copper	4.5×10^{-28}	[12]
VIP(2012)	Copper	4.7×10^{-29}	[13]
VIP2(goal)	Copper	$\times 10^{-31}$	[15]

Bayesian Analysis Method

Check on all parameters



New paradigm for VIP-2

Quantum gravity models can embed PEP violating transitions!

PEP is a consequence of the spin statistics theorem based on: Lorentz/Poincaré and CPT symmetries; locality; unitarity and causality. Deeply related to the very same nature of space and time



most effective theories of QG foresee the non-commutativity of the space-time quantum operators (e.g. k -Poincaré, θ -Poincaré)



non-commutativity induces a deformation of the Lorentz symmetry and of the locality \rightarrow naturally encodes the violation of PEP

S. Majid, Hopf algebras for physics at the Planck scale, *Class. Quantum Grav.* 5 (1988) 1587.

S. Majid and H. Ruegg, Bicrossproduct structure of Kappa Poincare group and noncommutative geometry, *Phys. Lett. B* 334 (1994) 348, hep-th/9405107.

M. Arzano and A. Marciano, *Phys. Rev. D* 76, 125005 (2007) [arXiv:0707.1329].

G. Amelino-Camelia, G. Gubitosi, A. Marciano, P. Martinetti and F. Mercati, *Phys. Lett. B* 671, 298 (2009) [arXiv:0707.1863].

A. Addazi, A. Marcianò *International Journal of Modern Physics A* Vol. 35, No. 32, 2042003 (2020)



PEP violation is suppressed with $(E/\Lambda)^n$, n depends on the specific model, E is the energy of the PEP violating transition, Λ is the scale of the space-time non-commutativity emergence.