From Hadrons to Therapy: Fundamental Physics Driving New Medical Advances

Contribution ID: 1

Type: not specified

On the relative role of the physical mechanisms on complex biodamage induced by carbon irradiation

Monday 5 September 2022 16:15 (40 minutes)

The effective use of swift ion beams in cancer treatment (known as hadrontherapy) as well as an appropriate protection in manned space missions rely on the accurate understanding of energy delivery to cells damaging their genetic information [1]. The key ingredient characterizing the response of a medium to the perturbation induced by charged particles is its electronic excitation spectrum. By using linear response time-dependent density functional theory, we obtain the energy and momentum transfer excitation spectrum (the energyloss function, ELF) of liquid water which is the main constituent of biological tissue, in excellent agreement with experimental data [2,3]. The inelastic scattering cross sections obtained from this ELF, together with the elastic scattering cross sections derived considering the condensed phase nature of the medium, are used to perform accurate Monte Carlo simulations of the energy deposited by swift carbon ions in liquid water and carried away by the generated secondary electrons producing inelastic events (ionization, excitation, and dissociative electron attachment), strongly correlated with cellular death, which are scored in sensitive volumes having the size of two DNA convolutions [2,3,4]. The sizes of clusters of damaging events for a wide range of carbon ion energies, from those relevant to hadrontherapy up to cosmic radiation, predict with unprecedented statistical accuracy the nature and relative magnitude of the main inelastic processes contributing to radiation biodamage, confirming that ionization accounts for the vast majority of complex damage, while DEA only adds up for a minor contribution. Applications to the calculation of the ELF and REEL spectra in ceria [5] and of beta-decay will be shown in this talk [6,7]

REFERENCES

[1] M. Dapor, I. Abril, P. de Vera, R. Garcia-Molina, Physical Review B 96, 064113 (2017).

[2] S. Taioli et al., The Journal of Physical Chemistry Letters 12 (1), 487 (2020)

[3] P. de Vera et al., Energy Deposition around Swift Carbon-Ion Tracks in Liquid Water, International Journal of Molecular Sciences 23 (11), 6121 (2022)

[4] M. Dapor, Transport of Energetic Electrons in Solids, 3rd ed. (Springer Nature, 2020).

[5] A. Pedrielli, et al., Physical Chemistry Chemical Physics (accepted, 2020)

- [6] T. Morresi, S. Taioli, S. Simonucci (2018), Advanced Theory and Simulations 1, 1870030 (2018)
- [7] S. Taioli, S. Simonucci, Annual Reports in Computational Chemistry 17 (2021)

Author: TAIOLI, Simone (European Centre for Theoretical Studies in Nuclear Physics and Related Areas, Trento, Italy)

Co-authors: ABRIL, Isabel (Universitat d'Alacant); DAPOR, Maurizio; DE VERA GOMIS, Pablo (Universidad de Murcia, Murcia/E); TREVISANUTTO, Paolo E. (Università Campus Biomedico, Roma); GARCIA-MOLINA, Rafael (Centro de Investigación en Óptica y Nanofísica, Universidad de Murcia, Murcia/E); SIMONUCCI, Stefano (Università di Camerino)

Presenter: TAIOLI, Simone (European Centre for Theoretical Studies in Nuclear Physics and Related Areas, Trento, Italy)

Session Classification: New developments in the modelling of radiation propagation and effects

Track Classification: Modelling of radiation propagation, effects and radiobiology