
Advances in the Lattice QCD calculation of TMDs

ECT* Workshop: Opportunities with JLab
Energy and Luminosity Upgrade

Trento, Italy, Sep. 26—30, 2022


YONG ZHAO
SEP 28, 2022



Outline

- **TMDs in the non-perturbative region**
- **Lattice calculations from quasi-TMDs**
- **Outlook**


3D Tomography of the Proton



Jefferson Lab 12 GeV





+



The Electron-Ion Collider

3D Tomography of the Proton

Leading Quark TMDPDFs



		Quark Polarization		
		Un-Polarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \text{circle with red dot}$ Unpolarized		$h_1^\perp = \text{circle with red dot} - \text{circle with red dot}$ Boer-Mulders
	L		$g_1 = \text{circle with red dot} \rightarrow - \text{circle with red dot} \rightarrow$ Helicity	$h_{1L}^\perp = \text{circle with red dot} \rightarrow - \text{circle with red dot} \rightarrow$ Worm-gear
	T	$f_{1T}^\perp = \text{circle with red dot} \uparrow - \text{circle with red dot} \downarrow$ Sivers	$g_{1T}^\perp = \text{circle with red dot} \uparrow - \text{circle with red dot} \uparrow$ Worm-gear	$h_1 = \text{circle with red dot} \uparrow - \text{circle with red dot} \downarrow$ Transversity $h_{1T}^\perp = \text{circle with red dot} \uparrow - \text{circle with red dot} \uparrow$ Pretzelosity


From TMD Handbook, TMD Topical Collaboration, to appear soon.

TMDs from experiments

TMD processes:

Semi-Inclusive DIS


$$\sigma \sim f_{q/P}(x, k_T) D_{h/q}(x, k_T)$$



**HERMES, COMPASS,
JLab, EIC, ...**

Drell-Yan

$$\sigma \sim f_{q/P}(x, k_T) f_{q/P}(x, k_T)$$




$$q_T \ll Q$$

**Fermilab, RHIC,
LHC, ...**

Dihadron in e^+e^-

$$\sigma \sim D_{h_1/q}(x, k_T) D_{h_2/q}(x, k_T)$$



**Babar, Belle,
BESIII, ...**

TMDs from global analyses

Semi-inclusive deep inelastic scattering: $l + p \rightarrow l + h(P_h) + X$

$$\frac{d\sigma^W}{dxdydz_hd^2\mathbf{P}_{hT}} \sim \int d^2\mathbf{b}_T e^{i\mathbf{b}_T \cdot \mathbf{P}_{hT}/z}$$

$$\times f_{i/p}(x, \mathbf{b}_T, Q, Q^2) D_{h/i}(z_h, \mathbf{b}_T, Q, Q^2)$$


$$f_{i/p}(x, \mathbf{b}_T, \mu, \zeta) = f_{i/p}^{\text{pert}}(x, b^*(b_T), \mu, \zeta)$$

$$\times \left(\frac{\zeta}{Q_0^2} \right)^{g_K(b_T)/2}$$

$$Q_0 \sim 1 \text{ GeV}$$

$$f_{i/p}^{\text{NP}}(x, b_T) \longrightarrow \begin{array}{l} \text{Collins-Soper kernel (NP part)} \\ \text{Intrinsic TMD} \end{array}$$


Non-perturbative when $b_T \sim 1/\Lambda_{\text{QCD}}$!



Kang, Prokudin, Sun and Yuan, PRD
93, 014009 (2016)


TMDs from global analyses

Unpolarized quark TMD



Scimemi and Vladimirov, JHEP 06 (2020).


Quark Sivers function



Cammarota, Gamberg, Kang et al. (JAM Collaboration),
PRD 102 (2020).

TMDs from global analyses


Collins-Soper Kernel $K(b_T, \mu) = K^{\text{pert}}(b_T, \mu) + g_K(b_T)$



Bacchetta, Bertone, Bissolotti, et al., MAP Collaboration, 2206.07598

TMD definition


- Beam function:



Hadronic matrix element

$$n_b^2 = 0$$

- Soft function :



Vacuum matrix element

$$f_i(x, \mathbf{b}_T, \mu, \zeta) = \lim_{\epsilon \rightarrow 0} Z_{\text{UV}} \lim_{\tau \rightarrow 0} \frac{B_i}{\sqrt{S^q}}$$


Collins-Soper scale: $\zeta = 2(xP^+e^{-y_n})^2$

Rapidity divergence regulator

First principles calculation of TMDs from the above matrix elements would greatly complement global analyses!

TMD definition

- Beam function:




$$n_b^2 = 0$$

Rapidity divergences

Hadronic matrix element

- Soft function :



Vacuum matrix element

$$f_i(x, \mathbf{b}_T, \mu, \zeta) = \lim_{\epsilon \rightarrow 0} Z_{\text{UV}} \lim_{\tau \rightarrow 0} \frac{B_i}{\sqrt{S^q}}$$



Collins-Soper scale: $\zeta = 2(xP^+e^{-y_n})^2$

Rapidity divergence regulator

First principles calculation of TMDs from the above matrix elements would greatly complement global analyses!

Lattice QCD

Lattice gauge theory: a systematically improvable approach to solve non-perturbative QCD.



Imaginary time: $t \rightarrow i\tau$ $O(i\tau) \xrightarrow{?} O(t)$


Simulating real-time dynamics has been extremely difficult due to the issue of analytical continuation. 😞

Progress in the lattice study of TMDs


- **Lorentz invariant method**
 - Musch, Hägler, Engelhardt, Negele and Schäfer et al.
 - Primary efforts focused on **ratios of TMD x-moments (w/o soft function)** (2009–)
- **Quasi-TMDs**
 - Large-momentum effective theory (Ji, 2013, 2014; Ji, Liu, Liu, Zhang and YZ, 2021)
 - One-loop studies of quasi beam and soft functions (Ji, Yuan, Scäfer, Liu, Liu, Ebert, Stewart, YZ, Vladimirov, Wang, ..., 2015-2022)
 - Method to calculate the Collins-Soper kernel (Ji, Yuan et al., 2015; Ebert, Stewart and YZ, 2018)
 - Method to calculate the soft function, and thus the **x and b_T dependence of TMDs** (Ji, Liu and Liu, 2019)
 - Derivation of factorization formula (Ebert, Schindler, Stewart and YZ, 2022)
 - First lattice results (SWZ, LPC, ETMC/PKU, SVZES, 2020–)

Quasi TMD in the LaMET formalism

- Beam function in Collins scheme:



Lightcone direction
Lorentz boost and $L \rightarrow \infty$



Spacelike but close-to-lightcone
($y_B \rightarrow -\infty$) Wilson lines, not
calculable on the lattice 😞

Equal-time Wilson lines, directly
calculable on the lattice 😊

Related by Lorentz invariance, equivalent in the
large \tilde{P}^z or $(-y_B)$ expansion.

Ebert, Schindler, Stewart and
YZ, JHEP 04, 178 (2022).

TMDs from lattice QCD

$$\frac{\tilde{f}_{i/p}^{\text{naive}[s]}(x, \mathbf{b}_T, \mu, \tilde{P}^z)}{\sqrt{S_r^q(\mathbf{b}_T, \mu)}} = C(\mu, x\tilde{P}^z) \exp\left[\frac{1}{2}K(\mu, b_T)\ln\frac{(2x\tilde{P}^z)^2}{\zeta}\right] \\ \times f_{i/p}^{[s]}(x, \mathbf{b}_T, \mu, \zeta) \left\{ 1 + \mathcal{O}\left[\frac{1}{(x\tilde{P}^z b_T)^2}, \frac{\Lambda_{\text{QCD}}^2}{(x\tilde{P}^z)^2}\right] \right\}$$

Reduced soft function ✓

Ji, Liu and Liu, NPB 955 (2020),
PLB 811 (2020).

- Ji, Sun, Xiong and Yuan, PRD91 (2015);
- Ji, Jin, Yuan, Zhang and YZ, PRD99 (2019);
- Ebert, Stewart, YZ, PRD99 (2019), JHEP09 (2019) 037;
- Ji, Liu and Liu, NPB 955 (2020), PLB 811 (2020);
- Vladimirov and Schäfer, PRD 101 (2020);
- Ebert, Schindler, Stewart and YZ, JHEP 04, 178 (2022).

Matching coefficient:

- **Independent of spin;**
 - Vladimirov and Schäfer, PRD 101 (2020);
 - Ebert, Schindler, Stewart and YZ, JHEP 09 (2020);
 - Ji, Liu, Schäfer and Yuan, PRD 103 (2021).
- **No quark-gluon or flavor mixing, which makes gluon calculation much easier.**

One-loop matching for gluon TMDs:

Ebert, Schindler, Stewart and YZ, 2205.12369.

TMDs from lattice QCD

$$\frac{\tilde{f}_{i/p}^{\text{naive}[s]}(x, \mathbf{b}_T, \mu, \tilde{P}^z)}{\sqrt{S_r^q(\mathbf{b}_T, \mu)}} = C(\mu, x\tilde{P}^z) \exp\left[\frac{1}{2}K(\mu, \mathbf{b}_T)\ln\frac{(2x\tilde{P}^z)^2}{\zeta}\right] \\ \times f_{i/p}^{[s]}(x, \mathbf{b}_T, \mu, \zeta) \left\{ 1 + \mathcal{O}\left[\frac{1}{(x\tilde{P}^z b_T)^2}, \frac{\Lambda_{\text{QCD}}^2}{(x\tilde{P}^z)^2}\right] \right\}$$

* Collins-Soper kernel;

$$K(\mu, \mathbf{b}_T) = \frac{d}{d \ln \tilde{P}^z} \ln \frac{\tilde{f}_{i/p}^{\text{naive}[s]}(x, \mathbf{b}_T, \mu, \tilde{P}^z)}{C(\mu, x\tilde{P}^z)}$$

* Flavor separation;

$$\frac{f_{i/p}^{[s]}(x, \mathbf{b}_T)}{f_{j/p}^{[s']}(x, \mathbf{b}_T)} = \frac{\tilde{f}_{i/p}^{\text{naive}[s]}(x, \mathbf{b}_T)}{\tilde{f}_{j/p}^{\text{naive}[s']}(x, \mathbf{b}_T)}$$

* Spin-dependence, e.g., Sivers function (single-spin asymmetry);


* Full TMD kinematic dependence.

* Twist-3 PDFs from small b_T expansion of TMDs. Ji, Liu, Schäfer and Yuan, PRD 103 (2021).

* Higher-twist TMDs.

Rodini and Vladimirov, JHEP 08 (2022).

Collins-Soper (CS) kernel from lattice QCD




Current status for the Collins-Soper kernel

	Lattice setup	Renormalization	Operator mixing	Fourier transform	Matching	x-plateau search
SWZ20 PRD 102 (2020) Quenched	$a = 0.06 \text{ fm}$, $m_\pi = 1.2 \text{ GeV}$, $P_{\max}^z = 2.6 \text{ GeV}$	Yes	Yes	Yes	LO	Yes
LPC20 PRL 125 (2020)	$a = 0.10 \text{ fm}$, $m_\pi = 547 \text{ MeV}$, $P_{\max}^z = 2.11 \text{ GeV}$	N/A	No (small)	N/A	LO	N/A
SVZES 21 JHEP 08 (2021)	$a = 0.09 \text{ fm}$, $m_\pi = 422 \text{ MeV}$, $P_{\max}^+ = 2.27 \text{ GeV}$	N/A	No	N/A	NLO	N/A
PKU/ETMC 21 PRL 128 (2022)	$a = 0.09 \text{ fm}$, $m_\pi = 827 \text{ MeV}$, $P_{\max}^z = 3.3 \text{ GeV}$	N/A	No	N/A	LO	N/A
SWZ21 PRD 106 (2022)	$a = 0.12 \text{ fm}$, $m_\pi = 580 \text{ MeV}$, $P_{\max}^z = 1.5 \text{ GeV}$	Yes	Yes	Yes	NLO	Yes
LPC22 PRD 106 (2022)	$a = 0.12 \text{ fm}$, $m_\pi = 670 \text{ MeV}$, $P_{\max}^z = 2.58 \text{ GeV}$	Yes	No (small)	Yes	NLO	Yes

Collins Soper kernel

Comparison between lattice results and global fits



Approach	Collaboration
Quasi beam functions	P. Shanahan, M. Wagman and YZ (SWZ21), Phys. Rev.D 104 (2021)
Quasi TMD wavefunctions	Q.-A. Zhang, et al. (LPC20), Phys.Rev.Lett. 125 (2020).
Moments of quasi TMDs	Y. Li et al. (ETMC/PKU 21), Phys.Rev.Lett. 128 (2022).
	M.-H. Chu et al. (LPC22), Phys.Rev.D 106 (2022)
	Schäfer, Vladimirov et al. (SVZES21), JHEP 08 (2021)

MAP22: Bacchetta, Bertone, Bissolotti, et al., 2206.07598

SV19: I. Scimemi and A. Vladimirov, JHEP 06 (2020) 137

Pavia19: A. Bacchetta et al., JHEP 07 (2020) 117

Pavia 17: A. Bacchetta et al., JHEP 06 (2017) 081

CASCADE: Martinez and Vladimirov, 2206.01105

Improved calculation with TMD wave function

$\tilde{\Phi}$: Quasi-TMD wave function

$$\tilde{\Phi} = \langle 0 | \text{Diagram} | \pi(P) \rangle$$

Q.-A. Zhang, et al. (LPC), PRL 125 (2020);
Y. Li et al., PRL 128 (2022);
M.-H. Chu et al. (LPC22), Phys.Rev.D 106 (2022).



- **Physical pion mass** and reduced systematics from Fourier transform

- Better suppressed power correction
- More stable extraction of x-dependence

- Renormalization of nonlocal operator


- Systematic treatment of operator mixing using the RI-xMOM scheme
- Green, Jansen, and Steffens, Phys.Rev.Lett. 121 (2018) and PRD 101 (2020).
• Constantinou, Panagopoulos, and Spanoudes, PRD 99 (2019).

A. Avkhadiev, P. Shanahan, M. Wagman and YZ,
work in progress.



Reduced soft function from LaMET

Light-meson form factor: $F(b_T, P^z) = \langle \pi(-P) | j_1(b_T) j_2(0) | \pi(P) \rangle$



$$\begin{aligned} P^z \gg m_N &= S_q^r(b_T, \mu) \int dx dx' H(x, x', \mu) \\ &\times \Phi^\dagger(x, b_T, P^z) \Phi(x', b_T, P^z) \end{aligned}$$

Tree-level approximation:


- Ji, Liu and Liu, NPB 955 (2020), PLB 811 (2020);
- Ji and Liu, PRD 105, 076014 (2022);
- Deng, Wang and Zeng, 2207.07280.

$$H(x, x', \mu) = 1 + \mathcal{O}(\alpha_s)$$

$$\Rightarrow S_q^r(b_T) = \frac{F(b_T, P^z)}{[\tilde{\Phi}(b^z = 0, b_T, P^z)]^2}$$


First lattice results with tree-level matching

$$\begin{aligned}a &= 0.10 \text{ fm}, \\m_\pi &= 547 \text{ MeV}, \\P_{\max}^z &= 2.11 \text{ GeV}\end{aligned}$$



Q.-A. Zhang, et al. (LPC), PRL 125 (2020).

$$\begin{aligned}a &= 0.09 \text{ fm}, \\m_\pi &= 827 \text{ MeV}, \\P_{\max}^z &= 3.3 \text{ GeV}\end{aligned}$$



Y. Li et al., PRL 128 (2022).

Beyond tree-level, it is necessary to obtain the x -dependence to carry out the convolution.

Conclusion

- The quark and gluon quasi TMDs can be related to the new LR scheme, which can be factorized into the physical TMDs;
- There is no mixing between quarks of different flavors, quark and gluon channels, or different spin structures.
- The method for calculating all the leading-power TMDs is complete;
- Lattice results for the Collins-Soper kernel and soft function are promising, but systematics need to be under control.


Outlook

Targets for lattice QCD studies:

Observables	Status
Non-perturbative Collins-Soper kernel	✓, keep improving the systematics
Soft factor	✓, to be under systematic control
Info on spin-dependent TMDs (in ratios)	In progress
Proton v.s. pion TMDs, (x, b_T) (in ratios)	In progress
Flavor dependence of TMDs, (x, b_T) (in ratios)	to be studied
TMDs and TMD wave functions, (x, b_T)	In progress
Gluon TMDs (x, b_T)	to be studied
Wigner distributions/GTMDs (x, b_T)	to be studied

Backup slides


Data used by the MAP collaboration in 2206.07598



Bacchetta, Bertone, Bissolotti, et al., MAP Collaboration, 2206.07598

LaMET calculation of the collinear PDFs


A state-of-the-art calculation of the pion valence quark PDF with fine lattices, large momentum and NNLO matching:



Gao, Hanlon, Mukherjee, Petreczky, Scior, Syritsyn and YZ, PRL 128, 142003 (2022).

Factorization relation with the TMDs

Lattice



$$\tilde{f}_i(x, \mathbf{b}_T, \mu, \tilde{\zeta}, \tilde{P}^z) = \lim_{\tilde{P}^z \gg m_N} \lim_{a \rightarrow 0} \tilde{Z}_{\text{UV}} \frac{\tilde{B}_i}{\sqrt{S^q}}$$

Lorentz invariance

$$y_{\tilde{P}} = y_P - y_B$$

$$f_i^{\text{LR}}(x, \mathbf{b}_T, \mu, \zeta, y_P - y_B) = \lim_{-y_B \gg 1} \lim_{\epsilon \rightarrow 0} Z_{\text{UV}}^{\text{LR}} \frac{B_i}{\sqrt{S^q}}$$

Same matrix elements, but
different orders of UV limits

Perturbative matching in
LaMET!

$$f_i(x, \mathbf{b}_T, \mu, \zeta) = \lim_{\epsilon \rightarrow 0} Z_{\text{UV}} \lim_{y_B \rightarrow -\infty} \frac{B_i}{\sqrt{S^q}}$$


Continuum

Ebert, Schindler, Stewart and YZ, JHEP 04, 178 (2022).

Factorization relation with the TMDs

Lattice

Quasi



Large-rapidity
(LR)



Collins

Continuum

$$\tilde{f}_i(x, \mathbf{b}_T, \mu, \tilde{\zeta}, \tilde{P}^z) = \lim_{\tilde{P}^z \gg m_N} \lim_{a \rightarrow 0} \tilde{Z}_{\text{UV}} \frac{\tilde{B}_i}{\sqrt{S^q}}$$

Lorentz invariance

$$y_{\tilde{P}} = y_P - y_B$$

$$f_i^{\text{LR}}(x, \mathbf{b}_T, \mu, \zeta, y_P - y_B) = \lim_{-y_B \gg 1} \lim_{\epsilon \rightarrow 0} Z_{\text{UV}}^{\text{LR}} \frac{B_i}{\sqrt{S^q}}$$


Same matrix elements, but
different orders of UV limits

Perturbative matching in
LaMET!


$$f_i(x, \mathbf{b}_T, \mu, \zeta) = \lim_{\epsilon \rightarrow 0} Z_{\text{UV}} \lim_{y_B \rightarrow -\infty} \frac{B_i}{\sqrt{S^q}}$$

Ebert, Schindler, Stewart and YZ, JHEP 04, 178 (2022).

Backup slides



**i, j (including spinor indices)
remain intact**



$\propto \delta_{ij}$ **Can mix with singlet
channel and with gluons**

$$b^2 = -b_z^2 - b_T^2 < b_T^2 \sim 1/\Lambda_{\text{QCD}}^2$$

**Hard particles cannot propagate
that far!**