



A DYNAMICAL INFLATON COUPLED TO STRONGLY INTERACTING MATTER



HOLOGRAPHIC QFTS ON CURVED BACKGROUNDS

With Jorge Casalderrey-Solana, Christian Ecker, Elias Kiritsis and David Mateos

Reference: 2011.08194 (JHEP), 2109.10355 (JHEP) and 2302.06618

Wilke van der Schee

Holographic perspectives on chiral transport
ECT*, Trento, 14 March 2023

MOTIVATION

Strongly coupled QFT on de Sitter space

- Can be relevant for the early universe
- Many first principle questions arise in de Sitter: thermodynamics, entropy etc
- Can we understand **inflation/reheating**?

Evolution with dynamical boundary gravity and an inflaton

- Semi-classical Einstein Equations on the boundary
- A technical subtlety: how to extract the log terms?
- *Example* with inflation, running down of inflaton and reheating

Strongly coupled sector like in hidden sector *if* relevant for our own universe

Major caveat: this is *not quantum gravity on de Sitter*

INFLATION, PREHEATING AND REHEATING

$$S = \int d^4x\sqrt{-g}\left[-\frac{m_{\rm Pl}^2}{2}R + \frac{1}{2}\nabla_\mu\phi\nabla^\mu\phi - V(\phi)\right] + S_{\rm matter}$$

Inflation

- Inflaton vacuum energy leads to exponential expansion of the universe
- Solves several problems in cosmology
- Inflaton experiences friction and settles at bottom of potential
- After inflation one is left with an empty universe

$$\ddot{\bar{\phi}} + 3H\dot{\bar{\phi}} + \partial_{\bar{\phi}}V(\bar{\phi}) = 0$$

Inflaton potential



How to get a working universe

- The inflaton energy needs to be transferred to Standard Model

Perturbative Reheating

- Inflaton decays at an (approximately) constant rate (faster than H)
- Decaying particles thermalize → big bang

Preheating

- Bose enhancement leads to resonances: exponential transfer of energy
- Bose enhanced states thermalize/reheat → big bang



Lev Kofman, Andrei Linde and Alexei A. Starobinsky, Towards the Theory of Reheating After Inflation (1997)

INFLATION, REHEATING AND HOLOGRAPHY

Holographic set-up

- Just like standard EH + inflaton + coupling to scalar:

$$S_{hol} = S_* + \int \mathrm{d}^4x\sqrt{-\gamma}\phi_{(0)}\langle\mathcal{O}\rangle$$

- Inflaton acts as a source for the scalar operator

$$S = \int d^4x\sqrt{-g}\left[-\frac{m_{\rm Pl}^2}{2}R + \frac{1}{2}\nabla_\mu\phi\nabla^\mu\phi - V(\phi)\right] + S_{\rm matter}$$

$$\ddot{\phi}_0 = -U\phi_0\langle\mathcal{O}\rangle - 3H\dot{\phi}_0 - V'(\phi_0)$$

Inflaton potential



Two technical complications:

- Sources are now time-dependent numerical functions
- holographic renormalization becomes more complicated

Trial and error:

- Initial inflaton position, inflaton velocity, coupling strength, shape potential
- Balance between numerical stability, runtime and making a ‘realistic’ universe

$$ds_{\rm b}^2 = -{\rm d}t^2 + S_0(t)^2 {\rm d}\vec{x}^2$$

A DYNAMICAL BOUNDARY

Study dynamics including semi-classical gravity:

$$R_{\mu\nu} - \tfrac{1}{2} R\, g_{\mu\nu} = 8\pi G_{N,4} \langle T_{\mu\nu} \rangle$$

$$\ddot{\phi}_0 = -U \phi_0 \langle \mathcal{O} \rangle - 3H\dot{\phi}_0 - V'(\phi_0)$$

- Stress-tensor includes possible cosmological constant
- NB: renormalisation counterterms are now physical
- We treat the boundary Newton constant as a (small) parameter

Dynamics of scale factor $S_0(t)$ is now consequence of Friedmann equations

How to initialise the dynamics

- Start with thermal Minkowski profile and small boundary $G_{N,4}$

Jewel Ghosh, Elias Kiritsis, Francesco Nitti and Lukas Witkowski, Back-reaction in de Sitter QFTs: holography, gravitational DBI action and f(R) gravity (2020)
Paul Chesler and Abraham Loeb, Holographic duality and mode stability of de Sitter space in semiclassical gravity (2020)
Christian Ecker, WS, David Mateos and Jorge Casalderrey-Solana , Holographic evolution with dynamical boundary gravity (2021)

$$ds_{\rm b}^2 = -{\rm d}t^2 + S_0(t)^2 {\rm d}\vec{x}^2$$

SOME NUMERICAL DETAILS

Three bulk fields:

$$ds^2_{\rm bulk} = -A(r,t){\rm d}t^2 + 2{\rm d}r{\rm d}t + S(r,t)^2{\rm d}\vec{x}^2, \quad \Phi = \Phi(r,t)$$

Characteristic formulation leads to nested set of linear ODEs (spectral elements):

$$S'' = -\frac{2}{3}S\left(\Phi'\right)^2 ,$$
$$\dot{S}' = -\frac{2\dot{S}S'}{S} - \frac{2SV}{3} ,$$
$$\dot{\Phi}' = \frac{V'}{2} - \frac{3\dot{S}\Phi'}{2S} - \frac{3S'\dot{\Phi}}{2S} ,$$
$$A'' = \frac{12\dot{S}S'}{S^2} + \frac{4V}{3} - 4\dot{\Phi}\Phi'$$

Boundary 4D Einstein-inflaton equations:

$$H(t)^2 = \frac{\kappa_4}{3}\mathcal{E}(t) ,$$
$$\frac{a''(t)}{a(t)} = -\frac{1}{2}\left(\kappa_4\mathcal{P}(t) + H(t)^2\right) ,$$
$$\phi''(t) = \partial_\phi U(\phi(t))\,\mathcal{O}_{\rm QFT}(t) - 3H(t)\phi'(t) - \partial_\phi V_{\rm inf}(\phi(t))$$

S and Φ require boundary conditions: scale factor a(t) and inflaton ϕ(t)

- Normally sources are specified by hand; here: 4D Einstein-Inflaton equations
- Boundary conditions implemented through near-boundary expansion
- Subtract first few logs analytically (for stability in spectral elements)
- A requires initial boundary condition (energy density), then determined by SE-conservation

See also www.wilkevanderschee.nl for the full numerical code.

A DYNAMICAL BOUNDARY – CONSISTENT INITIAL CONDITIONS

$$ds_b^2 = -dt^2 + S_0(t)^2 d\vec{x}^2$$

Near-boundary expansion scalar (and metric):

$$\Phi = \frac{\phi(t)}{r} + \frac{\phi_2(t)}{r^3} + \frac{1}{r}\sum_{n\geq 3}\frac{\phi_n(t)}{r^n} + \frac{1}{r}\sum_{n\geq 2}\frac{\psi_n(t)\log r}{r^n} + \cdots$$

- Crucial subtlety: logarithmic terms ψ_n solely determined by sources $S_0(t)$
- Consistent IC not much of a problem with known source (ψ_n known analytically)

Dynamics of scale factor $S_0(t)$ is now consequence of Friedmann equations

Sufficiently smooth solution requires knowledge of sufficient # of log's

- Log's depend on time derivatives of $S_0(t)$
- Solution: extract $\partial_t^4 S_0(t)$ from near-boundary expansion of scalar (extract ϕ_n)
- Use time derivates to treat first few logs analytically

Final subtlety: VEV and energy density depend on $S_0''(t)$ and $\phi''(t)$

- Solve resulting 6th order polynomial equation numerically

Sebastian de Haro, Kostas Skenderis and Sergey N. Solodukhin, Holographic Reconstruction of Spacetime and Renormalization in the AdS/CFT Correspondence (2001)

BOUNDARY GRAVITY ONLY

$$ds_{\rm b}^2 = -{\rm d}t^2 + S_0(t)^2 {\rm d}\vec{x}^2$$

Stress-energy tensor for zero, positive and negative Λ



Positive Λ settles down to de Sitter state as studied before, includes (small) Casimir energy

Now also asymptotically flat + Big Crunch geometries

Christian Ecker, WS, David Mateos and Jorge Casalderrey-Solana, Holographic evolution with dynamical boundary gravity (2021)

BOUNDARY GRAVITY WITH AN INFLATON

Inflaton rolls down and undergoes damped oscillation

Rolls down relatively fast (helps numerical stability)

Inflaton potential



Inflaton dynamics



INFLATION AND REHEATING

Initially dominated by QFT energy

- Quickly dilutes due to expansion

Then dominated by inflaton

- Inflation, while slowly rolling down

Then inflaton reaches bottom: reheating

- Inflaton loses energy to QFT

Finally universe dominated by QFT in thermal equilibrium

Hubble rate



Energy densities (log-log)



Inflating phase (t ~ 5 till 12)

Reheating (around t=13)

TEMPERATURE FROM BULK AND QFT

Three ways to determine temperature:

- From energy density and EOS
- From apparent horizon
- From event horizon (agrees to apparent horizon)

Always `close to equilibrium'

- Somewhat curious Casimir shift by de Sitter temperature
- Also present in analytical solution of conformal case

$$\epsilon = \frac{3}{8}\pi^2 N^2 T^4 + \frac{3N^2(\dot{a})^4}{32\pi^2 a^4}$$

Temperature (log-log)



Alex Buchel, Michal P. Heller and Jorge Noronha, Entropy Production, Hydrodynamics, and Resurgence in the Primordial Quark-Gluon Plasma from Holography (2016)

HYDRODYNAMICS

$$\langle \hat{T}^{ij} \rangle = \epsilon\, u^i u^j + p_{\rm eq}(\epsilon)\Delta^{ij} - \eta(\epsilon)\,\sigma^{ij} - \zeta(\epsilon)\Delta^{ij}\overline{\nabla}^k u_k + O(\overline{\nabla}^2)\,,$$

$$\sigma^{ij} = \Delta^{ik}\Delta^{jl}(\overline{\nabla}_k u_l + \overline{\nabla}_l u_k) - \frac{2}{3}\Delta^{ij}\overline{\nabla}_k u^k\,,$$

Hydrodynamic constitutive relation (ideal+viscous):

$$\Delta\mathcal{P}^{\rm hydro}(t) \equiv \frac{N^2}{2\pi^2}\left\{\Delta p_{\rm eq}(\Delta\mathcal{E}(t)) - 3H\zeta(\Delta\mathcal{E}(t))\right\} + O(H^2)$$







- Quick initial hydrodynamisation (t~1)
- **Viscous** hydrodynamics during inflation (t ~ 5 – 13)
- Far-from-equilibrium during reheating (t~ 13 – 20)

DISCUSSION

Dynamical gravity and inflaton on the boundary

- Technical progress: how to deal with the logs?
- Several interesting evolution: flat space, Big Crunch

Inflation and reheating

- First principles computation of strongly coupled reheating
- Strongly coupled physics can thermalise after preheating
- Currently not a `realistic universe'

Future

- Realistic universe? Many e-foldings based on hydro?
- Do we understand entropy of de Sitter?





BACK-UP

NON-CONFORMAL MODEL ON DE SITTER$_4$

De Sitter is conformally flat: almost trivial for CFT

- Break scale invariance by V(Φ) with source M=1:

$$S = \frac{2}{8\pi G}\int_{\mathcal{M}} d^5x\sqrt{-g}\left(\frac{1}{4}R[g] - \frac{1}{2}(\partial\phi)^2 - V(\phi)\right)$$

$$L^2V(\phi) = -3 - \frac{3}{2}\phi^2 - \frac{1}{3}\phi^4 + \left(\frac{1}{3\phi_M^2} + \frac{1}{2\phi_M^4}\right)\phi^6 - \frac{1}{12\phi_M^4}\phi^8$$

- Leads to non-trivial EOS and bulk viscosity (no shear considered):

Entropy density



Trace anomaly



Bulk viscosity



Jorge Casalderrey-Solana, Christian Ecker, David Mateos and WS, Strong-coupling dynamics and entanglement in de Sitter space (2020)

$$ds_b^2 = -dt^2 + S_0(t)^2 d\vec{x}^2 \,. \quad S_0(t) = e^{Ht}$$

BULK GEOMETRY WITH dS BOUNDARY

Event, apparent and 'entanglement' horizons for bulk dual to empty de Sitter

- The apparent horizon is much beyond the event horizon
- Also shown: extremal surfaces
- `Temperature' of AH is negative; curiosity?
- Bulk confirms boundary intuition on Hawking temperature and boundary cosmological horizon



Apparent horizon: $T = -\frac{H}{2\pi}$

Entanglement horizon: $T = 0$

Event horizon: $T = \frac{H}{2\pi}$

Boundary



Cosmological horizon

Jorge Casalderrey, Christian Ecker, David Mateos and WS, Strong-coupling dynamics and entanglement in de Sitter space (2020)

TIME EVOLUTION OF THE PROTOCOL

Evolution of stress tensor for different Hubble constants

- Energy density decreases towards vacuum energy (VE) (afterwards renormalised to zero)
- Pressure decreases, changes sign and becomes –VE
- Enthalpy is scheme independent, decays due to expansion

Energy density



Pressure



Enthalpy



THE APPROACH TOWARDS HYDRODYNAMICS

$$\Pi = -\zeta\left(\nabla\cdot u\right) + \zeta\tau_{\Pi} D\left(\nabla\cdot u\right) + \xi_1\sigma^{\mu\nu}\sigma_{\mu\nu} + \xi_2\left(\nabla\cdot u\right)^2 + \xi_3\Omega^{\mu\nu}\Omega_{\mu\nu} + \xi_4\nabla_\mu^\perp \ln s \nabla_\perp^\mu \ln s + \xi_5 R + \xi_6 u^\alpha u^\beta R_{\alpha\beta}\,.$$

Non-trivial hydrodynamic prediction

$$\Delta\mathcal{P}^{\rm hydro}(t) \equiv \Delta p_{\rm eq}(\Delta\mathcal{E}(t)) - 3H\zeta(\Delta\mathcal{E}(t)) + O(H^2)\,,$$

- Δ captures our renormalisation scheme
- Conjecture: scheme ambiguities cancelled by higher order transport coefficient ξ_5

Results

- Viscous hydro works for small H (gradients). Negative `EOS' for large H.



Paul Romatschke, Relativistic Viscous Fluid Dynamics and Non-Equilibrium Entropy (2009)

$$ds^2 = -A(r,t)\mathrm{d}t^2 + 2\mathrm{d}r\mathrm{d}t + S(r,t)^2\mathrm{d}\vec{x}^2$$

BLACK HOLE THERMODYNAMICS



Keep track of bulk event and apparent horizons (EF coordinates)

- Dynamical setting: horizons not coincide at late times:
- Surface gravities can be shown analytically: $\kappa_{\mathrm{EH}} = -\kappa_{\mathrm{AH}} = H$
  EH confirms Hawking's temperature in de Sitter
- Area density apparent horizon vanishes for conformal theory

Surface gravity



Area densities



Willy Fischler, Sandipan Kundu and Juan Pedraza, Entanglement and out-of-equilibrium dynamics in holographic models of de Sitter QFTs (2013)
Alex Buchel, Entanglement entropy of N = 2* de Sitter vacuum (2019)

$$ds^2 = -A(r,t)\mathrm{d}t^2 + 2\mathrm{d}r\mathrm{d}t + S(r,t)^2\mathrm{d}\vec{x}^2$$

BLACK HOLE THERMODYNAMICS

Several interpretational issues

- Expanding space: mapping boundary to bulk horizon not clear
- Apparent horizon: time slicing dependent
- In general: no volume law entropy density expected

Resolution → entanglement entropy is well defined



Surface gravity



Area densities



Willy Fischler, Sandipan Kundu and Juan Pedraza, Entanglement and out-of-equilibrium dynamics in holographic models of de Sitter QFTs (2013)
Alex Buchel, Entanglement entropy of N = 2* de Sitter vacuum (2019)

ENTANGLEMENT IN DE SITTER

Extremal surfaces dual to spherical entangling regions:

- Surfaces probe beyond event horizon if and only if entangling region bigger than visible universe
- Only `superobservers' can see behind event horizon



Apparent horizon: $T = -\frac{H}{2\pi}$

Entanglement horizon: $T = 0$

Event horizon: $T = \frac{H}{2\pi}$

Boundary



Cosmological horizon

Jorge Casalderrey, Christian Ecker, David Mateos and WS, Strong-coupling dynamics and entanglement in de Sitter space (2020)

ENTANGLEMENT IN DE SITTER

Extremal surfaces go backward in time

- Time at the deepest point grows exactly as log(l) for large l
- Implies that `entanglement horizon' contribution has a **constant instead of volume law** contribution
- Standard `area law' divergence still applies





Cosmological horizon

COSMOLOGICAL HORIZON

Extremal surface dual to cosmological horizon:

- Separates points in the bulk from which light can reach the (boundary) origin
- *Boundary cosmological horizon → full bulk cosmological horizon*



Cosmological horizon

Jorge Casalderrey, Christian Ecker, David Mateos and WS, Strong-coupling dynamics and entanglement in de Sitter space (2020)

HOW WE SET UP A STATE

Non-trivial boundary metric: $ds_{\rm b}^2 = -{\rm d}t^2 + S_0(t)^2 {\rm d}\vec{x}^2 \,. \quad S_0(t) = e^{Ht}$

Start with thermal (high-temperature) state in flat space

- Quench system by suitable fast tanh to constant Hubble parameter
- Energy density decreases towards final `vacuum energy' (VE)
- Final (Bunch-Davis)-VE is ambiguous → chose scheme with zero VE

Hubble parameter



Energy density



$$ds_{\rm b}^2 = -{\rm d}t^2 + S_0(t)^2 {\rm d}\vec{x}^2 \,. \quad S_0(t) = e^{Ht}$$

THE NUMERICAL SCHEME

Eddington-Finkelstein metric ansatz: $ds^2 = -A(r,t){\rm d}t^2 + 2{\rm d}r{\rm d}t + S(r,t)^2{\rm d}\vec{x}^2$

- Advantage: Einstein-scalar equations factorise in nested set of linear equations

Perform near-boundary expansion of A **and** S

- Subtract first few divergent and log terms → treat them analytically
- Expand remainder on finite element mesh (6 domains with 12 spectral points)
- High precision time evolution (90 digit precision, 10000 time steps)
- Evolution only 1+1D, but still takes ~1 day on a laptop

THE APPROACH TOWARDS HYDRODYNAMICS

$$ds_{\rm b}^2 = -{\rm d}t^2 + S_0(t)^2 {\rm d}\vec{x}^2 \,. \quad S_0(t) = e^{Ht}$$

Comparing with the hydrodynamic constituent relations:

$$T_\perp^{\mu\nu} = P(\varepsilon)\Delta^{\mu\nu} - \eta(\varepsilon)\sigma^{\mu\nu} - \zeta(\varepsilon)\Delta^{\mu\nu}(\nabla\cdot u)\,,$$
$$\Delta^{\mu\nu} = g^{\mu\nu} + u^\mu u^\nu$$

Symmetric set-up: only non-trivial part is bulk viscosity:

$$\Delta\mathcal{P}^{\rm hydro}(t) \equiv \Delta p_{\rm eq}(\Delta\mathcal{E}(t)) - 3H\zeta(\Delta\mathcal{E}(t)) + O(H^2)\,,$$

A subtlety: EOS and viscosity computed in flat space;
what is the energy density in de Sitter space?

We decided to fix renormalisation freedom so that late time solution has zero energy density

(in any case: ambiguity is order H^2)
(also, note that scheme depends on H for our choice)

HOLOGRAPHIC RENORMALISATION

$$ds_{\rm b}^2 = -{\rm d}t^2 + S_0(t)^2 {\rm d}\vec{x}^2 \,. \quad S_0(t) = e^{Ht}$$

$$ds^2 = -A(r,t){\rm d}t^2 + 2{\rm d}r{\rm d}t + S(r,t)^2{\rm d}\vec{x}^2 \,, \quad \phi = \phi(r,t) \,,$$

Action needs (scheme-dependent) counter-terms:

$$S_{\rm ct} = \frac{1}{8\pi G}\int {\rm d}^4x\sqrt{-\gamma}\left[\left(-\frac{1}{8}R[\gamma] - \frac{3}{2} - \frac{1}{2}\phi^2\right) + \frac{1}{2}\left(\log\rho\right)\mathcal{A} + \left(\alpha\mathcal{A} + \beta\phi^4\right)\right],$$

$$\mathcal{A} = \mathcal{A}_g + \mathcal{A}_\phi \,, \qquad \mathcal{A}_g = \frac{1}{16}(R^{ij}R_{ij} - \frac{1}{3}R^2) \,, \qquad \mathcal{A}_\phi = -\frac{\phi^2}{12}R$$

Leads to an ambiguity in the stress-tensor:

$$\begin{aligned}\mathcal{E}(t) = & -\frac{3a_{(4)}(t)}{4} - M\bar{\phi}_{(2)}(t) + \frac{3S_0'(t)^4}{16S_0(t)^4} + M^2\left(\xi(t)^2 + \frac{S_0'(t)^2}{8S_0(t)^2} + \frac{2S_0''(t)}{3S_0(t)}\right) \\ & - M^2\alpha\frac{S_0'(t)^2}{2S_0(t)^2} - M^4\left(\beta - \frac{7}{36}\right),\end{aligned}$$

$$\begin{aligned}\mathcal{P}(t) = & -\frac{a_{(4)}(t)}{4} + \frac{1}{3}M\bar{\phi}_{(2)}(t) + \frac{S_0'(t)^2\left(S_0'(t)^2 - 4S_0(t)S_0''(t)\right)}{16S_0(t)^4} \\ & - \frac{M^2}{3}\left(\xi(t)^2 + \frac{S_0'(t)^2}{8S_0(t)^2} + \frac{13S_0''(t)}{12S_0(t)}\right) + M^2\alpha\left(\frac{S_0'(t)^2}{6S_0(t)^2} + \frac{S_0''(t)}{3S_0(t)}\right) + M^4\left(\beta - \frac{5}{108}\right)\end{aligned}$$

α and β encode scheme dependencies (cosmological constant); fixed such that late time solution has vanishing energy

Ambiguities come in at order H^2

BOUNDARY GRAVITY - TEMPERATURES

Temperatures extracted from surface gravity event and apparent horizons



De Sitter case again settles down with negative AH ‘temperature’

For the big crunch geometry the EH location is not obvious (no future timelike infinity)

HYDRODYNAMISATION AND BOUNDARY GRAVITY

Three different initial conditions for $\Lambda = 0$:

Stress-tensor, and hydrodynamisation:
all hydrodynamise within a time of ~1/T





HYDRODYNAMISATION AND BOUNDARY GRAVITY

AdS case: fast hydrodynamisation, hydro all the way till big crunch



dS case: hydrodynamisation, but falls out of equilibrium.
Casimir energy important.