

Color Glass Condensate at the Electron-Ion Collider
ECT*, Trento, May 18th 2023

Single inclusive particle production in pA collisions at forward rapidities: beyond the hybrid model

Néstor Armesto

IGFAE, Universidade de Santiago de Compostela

nestor.armesto@usc.es

with Tolga Altinoluk (NCBJ), Guillaume Beuf (NCBJ), Alex Kovner (UConn) and Michael Lublinsky (BGU)


Contents:

1. Introduction.
2. Our definition of TMD PDFs and FFs.
3. The $q \rightarrow q \rightarrow H$ channel.
4. Summary.

See the talks by Farid Salazar, Yossathorn Tawabutr, Yair Mulian, Leszek Motyka, Ian Balitsky,...


The hybrid model:

- Hybrid model proposed at LO in 2005 (hep-ph/0506308), NLO in 2011 (1112.1061): large x collinear parton which splits, rescatters with the target (eikonally) and fragments onto a hadron.



1405.6311



- Cross sections turned out to be **negative** at large transverse momentum, a problem alleviated at larger rapidities or energies.




The problem (I):

- Several solutions proposed along the years:
 - Kinematic constraints (1505.05183)/Ioffe time restriction (1411.2869) leading to new, BK-like terms.
 - Choice of rapidity scales (1403.5221, 1407.6314, 1608.05293).
 - Threshold (2004.11990) and Sudakov (2112.06975) resummation.

- They lead to a successful description of data but lack of understanding of what was or still is wrong, or of guidance on how to rectify it.




2004.11990




2112.06975

The problem (I):



- Several solutions proposed along the years:
 - Kinematic constraints (1505.05183)/Ioffe time restriction (1411.2869) leading to new, BK-like terms.
 - Choice of rapidity scales (1403.5221, 1407.6314, 1608.05293).
 - Threshold (2004.11990) and Sudakov (2112.06975) resummation.



• They lead to a successful description of data but lack of understanding of what was or still is wrong, or of guidance on how to rectify it.





2004.11990



2112.06975

The problem (II):

- Any eventual problem of negativity at NLO should not come from large transverse momentum: inelastic (real NLO) contribution squared (1102.5327), the elastic one (LO+virtual NLO) does not contribute unless the dipole has a large tail at $k_{\perp}^2 \gg Q_s^2$.



2112.06975


- The reason for the negativity is seemingly an over subtraction: the NLO is extracted collinear pieces that go to the DGLAP evolution of the collinear PDFs and FFs, and a soft piece (through the plus prescription) that goes into the BK evolution of the dipole scattering matrix. The remainder turns out to become negative at large transverse momentum.

- Here we conclude that the correct framework to resum all large logarithms is **not collinear factorization for the projectile (the hybrid model) but TMD factorization.**

Our setup:

- We work in a frame in which the target nucleus moves fast. We find a **TMD-factorized parton model expression**:

$$\int \frac{d\zeta}{\zeta^2} \int d^2k_{\perp} d^2q_{\perp} T\left(\frac{x_F}{\zeta}, k_{\perp}; \mu_T^2\right) P(k_{\perp}, q_{\perp}) F\left(\zeta; p_{\perp}, (k_{\perp} + q_{\perp}); \mu_F^2\right) + \mathcal{O}\left(\frac{p_{\perp}^2, k_{\perp}^2, Q_s^2, \mu^2}{s_0}\right)$$




- Our **scales** are

$$\mu_T^2 = \max\{k_{\perp}^2, q_{\perp}^2, Q_s^2\} \approx \max\{(k_{\perp} + q_{\perp})^2, Q_s^2\}, \mu_F^2 = ((q_{\perp} + k_{\perp}) - p_{\perp}/\zeta)^2 \approx \max\{(q_{\perp} + k_{\perp})^2, (p_{\perp}/\zeta)^2\}$$

Our TMD distributions: one flavor PDFs

- **TMD PDFs** (single parton species to start with) **are generated from collinear ones** (large k):

$$x\mathcal{T}_q(x, k^2; k^2; \xi_0) = \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{\xi_0}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \frac{x}{1 - \xi} f_{k^2}^q \left(\frac{x}{1 - \xi} \right) \frac{1}{k^2}$$


- **Evolution** (diagonal in parton species and momentum fraction; the second term corresponds to a loss due to the increase in resolution):

$$x\mathcal{T}_q(x, k^2; \mu^2; \xi_0) = \theta(\mu^2 - k^2) \left[x\mathcal{T}_q(x, k^2; k^2; \xi_0) - \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{k^2}^{\mu^2} \frac{\pi dl^2}{l^2} \int_{\xi_0}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} x\mathcal{T}_q(x, k^2; l^2; \xi_0) \right]$$

- **At $\mathcal{O}(\alpha_s)$:** $x\mathcal{T}_q(x, k^2; \mu^2; \xi_0) = \theta(\mu^2 - k^2) x\mathcal{T}_q(x, k^2; k^2; \xi_0) \left[1 - \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{k^2}^{\mu^2} \frac{\pi dl^2}{l^2} \int_{\xi_0}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \right]$

Our TMD distributions: one flavor PDFs

- With $xf_{\mu^2}^q(x) = \int_0^{\mu^2} \pi dk^2 x \mathcal{T}_q(x, k^2; \mu^2; \xi_0)$, we recover DGLAP for the collinear PDFs and their definition is independent of $\xi_0 \ll 1$.
- TMD FFs are defined analogously.
- These definitions can be generalised to n_f massless quarks and antiquarks, and gluons.
- It will turn out in the calculation that $\xi_0 \propto \mu^2/s_0$, with s_0 an energy scale that comes from the Ioffe time restriction ([1411.2869](#)).
- Relating these definitions and the corresponding evolution equations to more standard implementations ([see e.g. 2304.03302](#)) of the rapidity cut-off (s_0 in our case that acts as a longitudinal resolution) is yet to be done.

Our TMD distributions: one flavor FFs

- **TMD FFs** (single parton species):

$$\mathcal{F}_H^q(\zeta, k^2; k^2, \xi_0) = \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{\xi_0}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \frac{1}{1 - \xi} D_{H,k^2}^q \left(\frac{\zeta}{1 - \xi} \right) \frac{1}{k^2}$$

$$\mathcal{F}_H^q(x, k^2; \mu^2; \xi_0) = \theta(\mu^2 - k^2) \left[\mathcal{F}_H^q(x, k^2; k^2; \xi_0) - \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{k^2}^{\mu^2} \frac{\pi dl^2}{l^2} \int_{\xi_0}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \mathcal{F}_H^q(x, k^2; l^2; \xi_0) \right]$$

$$D_{H,\mu^2}^q(x) = \int_0^{\mu^2} \pi dk^2 \mathcal{F}_H^q(x, k^2; \mu^2; \xi_0)$$

Our TMD distributions: all flavor PDFs

- For n_f massless quarks and antiquarks, and gluons:

$$\begin{aligned}
 x\mathcal{T}_q(x, k^2; k^2; \xi_0) &= \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{\xi_0}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \frac{x}{1 - \xi} f_{k^2}^q \left(\frac{x}{1 - \xi} \right) \frac{1}{k^2} \\
 &+ \frac{g^2}{(2\pi)^3} \frac{1}{2} \int_{\xi_0}^1 d\xi [\xi^2 + (1 - \xi)^2] \frac{x}{1 - \xi} f_{k^2}^g \left(\frac{x}{1 - \xi} \right) \frac{1}{k^2}
 \end{aligned}
 \quad
 \begin{aligned}
 x\mathcal{T}_g(x, k^2; k^2; \xi_0) &= \frac{g^2}{(2\pi)^3} 2N_c \int_{\xi_0}^1 d\xi \left[\frac{1 - \xi}{\xi} + \frac{\xi}{1 - \xi} + \xi(1 - \xi) \right] \frac{x}{1 - \xi} f_{k^2}^g \left(\frac{x}{1 - \xi} \right) \frac{1}{k^2} \\
 &+ \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \sum_q \int_{\xi_0}^1 d\xi \frac{1 + \xi^2}{1 - \xi} \frac{x}{1 - \xi} \left[f_{k^2}^q \left(\frac{x}{1 - \xi} \right) + f_{k^2}^{\bar{q}} \left(\frac{x}{1 - \xi} \right) \right] \frac{1}{k^2}
 \end{aligned}$$

$$x\mathcal{T}_q(x, k^2; \mu^2; \xi_0) = \theta(\mu^2 - k^2) \left[x\mathcal{T}_q(x, k^2; k^2; \xi_0) - \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{k^2}^{\mu^2} \frac{\pi dl^2}{l^2} \int_{\xi_0}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} x\mathcal{T}_q(x, k^2; l^2; \xi_0) \right]$$

$$\begin{aligned}
 x\mathcal{T}_g(x, k^2; \mu^2; \xi_0) &= \theta(\mu^2 - k^2) \left[x\mathcal{T}_g(x, k^2; k^2; \xi_0) - \frac{g^2}{(2\pi)^3} N_c \int_{k^2}^{\mu^2} \frac{\pi dl^2}{l^2} \int_{\xi_0}^1 d\xi \left[\frac{1 - \xi}{\xi} + \frac{\xi}{1 - \xi} + \xi(1 - \xi) \right] x\mathcal{T}_g(x, k^2; l^2; \xi_0) \right. \\
 &\left. - \frac{g^2}{(2\pi)^3} \frac{n_f}{2} \int_{k^2}^{\mu^2} \frac{\pi dl^2}{l^2} \int_{\xi_0}^1 d\xi [\xi^2 + (1 - \xi)^2] x\mathcal{T}_g(x, k^2; l^2; \xi_0) \right]
 \end{aligned}$$

$$x f_{\mu^2}^g(x) = \int_0^{\mu^2} \pi dk^2 x\mathcal{T}_g(x, k^2; \mu^2; \xi_0)$$

- The collinear PDFs satisfy DGLAP.

Our TMD distributions: all flavor FFs

- For n_f massless quarks and antiquarks, and gluons:

$$\mathcal{F}_H^q(\zeta, k^2; k^2, \xi_0) = \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{\xi_0}^1 d\xi \frac{1 + (1-\xi)^2}{\xi} \frac{1}{1-\xi} D_{H,k^2}^q \left(\frac{\zeta}{1-\xi} \right) \frac{1}{k^2} + \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{\xi_0}^1 d\xi \frac{1 + \xi^2}{1-\xi} \frac{1}{1-\xi} D_{H,k^2}^g \left(\frac{\zeta}{1-\xi} \right) \frac{1}{k^2},$$

$$\mathcal{F}_H^g(x, k^2; k^2; \xi_0) = \frac{g^2}{(2\pi)^3} 2N_c \int_{\xi_0}^1 d\xi \left[\frac{1-\xi}{\xi} + \frac{\xi}{1-\xi} + \xi(1-\xi) \right] \frac{1}{1-\xi} D_{H,k^2}^g \left(\frac{x}{1-\xi} \right) \frac{1}{k^2} + \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \sum_q \int_{\xi_0}^1 d\xi [\xi^2 + (1-\xi)^2] \frac{1}{1-\xi} \left[D_{H,k^2}^q \left(\frac{x}{1-\xi} \right) + D_{H,k^2}^{\bar{q}} \left(\frac{x}{1-\xi} \right) \right] \frac{1}{k^2}$$

$$\mathcal{F}_H^q(x, k^2; \mu^2; \xi_0) = \theta(\mu^2 - k^2) \left[\mathcal{F}_H^q(x, k^2; k^2; \xi_0) - \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{k^2}^{\mu^2} \frac{\pi dl^2}{l^2} \int_{\xi_0}^1 d\xi \frac{1 + (1-\xi)^2}{\xi} \mathcal{F}_H^q(x, k^2; l^2; \xi_0) \right]$$

$$D_{H,\mu^2}^g(x) = \int_0^{\mu^2} \pi dk^2 \mathcal{F}_H^g(x, k^2; \mu^2; \xi_0) - \frac{g^2}{(2\pi)^3} N_c \int_{k^2}^{\mu^2} \frac{\pi dl^2}{l^2} \int_{\xi_0}^1 d\xi \left[\frac{1-\xi}{\xi} + \frac{\xi}{1-\xi} + \xi(1-\xi) \right] \mathcal{F}_H^g(x, k^2; l^2; \xi_0) - \frac{g^2}{(2\pi)^3} \frac{n_f}{2} \int_{k^2}^{\mu^2} \frac{\pi dl^2}{l^2} \int_{\xi_0}^1 d\xi [\xi^2 + (1-\xi)^2] \mathcal{F}_H^g(x, k^2; l^2; \xi_0).$$

- The collinear FFs satisfy DGLAP.

$q \rightarrow q \rightarrow H$: initial expressions

- We start from the expressions obtained in LCPT in [4] I.2869, before collinear subtraction, and we do not implement the plus prescription.
- We work at large N_c , and assume factorisation and translational invariance for the dipoles $s(r) \equiv s_F(r)$.

$$\frac{d\bar{\sigma}^{q \rightarrow q}}{d^2k d\eta}(k, x_p) = \frac{d\bar{\sigma}_0^{q \rightarrow q}}{d^2k d\eta}(k, x_p) + \frac{d\bar{\sigma}_{1,r}^{q \rightarrow q}}{d^2k d\eta}(k, x_p) + \frac{d\bar{\sigma}_{1,v}^{q \rightarrow q}}{d^2k d\eta}(k, x_p) \quad \frac{d\bar{\sigma}_0^{q \rightarrow q}}{d^2k d\eta}(k, x_p) = \frac{1}{(2\pi)^2} x_p f_{\mu_0^2}^q(x_p) \int_{y, \bar{y}} e^{ik \cdot (y - \bar{y})} s[y - \bar{y}]$$

- The WW factors with the Ioffe time restriction read

$$A_{\xi, x_p}^i(y - z) \equiv -i \int_{l^2 < \xi(1-\xi)x_p s_0} \frac{d^2l}{(2\pi)^2} \frac{l^i}{l^2} e^{-il \cdot (y-z)}, \quad A_{\xi}^i(y - z) \equiv A_{\xi, x_p/(1-\xi)}^i(y - z) \simeq A_{\xi, x_p}^i(y - z) \text{ for } k^2/(x_F s_0) \ll 1.$$

$$\begin{aligned} \frac{d\bar{\sigma}_{1,r}^{q \rightarrow q}}{d^2k d\eta}(k, x_p) &= \frac{g^2}{(2\pi)^3} \int_0^1 d\xi \int_{y, \bar{y}, z} e^{ik \cdot (y - \bar{y})} \frac{1 + (1 - \xi)^2}{\xi} \\ &\times \left[\frac{x_p}{1 - \xi} f_{\mu_0^2}^q \left(\frac{x_p}{1 - \xi} \right) A_{\xi}^i(y - z) A_{\xi}^i(\bar{y} - z) \left\{ C_F [s[y - \bar{y}] + s[(1 - \xi)(y - \bar{y})]] \right. \right. \\ &\quad \left. \left. - \frac{N_c}{2} [s[(1 - \xi)(y - z)] s[\bar{y} - z] + s[(1 - \xi)(\bar{y} - z)] s[y - z]] \right\} \right] \\ \frac{d\bar{\sigma}_{1,v}^{q \rightarrow q}}{d^2k d\eta}(k, x_p) &= -\frac{g^2}{(2\pi)^3} \int_0^1 d\xi \int_{y, \bar{y}, z} e^{ik \cdot (y - \bar{y})} \frac{1 + (1 - \xi)^2}{\xi} \\ &\times \left[x_p f_{\mu_0^2}^q(x_p) \left\{ C_F [A_{\xi, x_p}^i(y - z) A_{\xi, x_p}^i(y - z) + A_{\xi, x_p}^i(\bar{y} - z) A_{\xi, x_p}^i(\bar{y} - z)] s[y - \bar{y}] \right. \right. \\ &\quad - \left\{ A_{\xi, x_p}^i(y - z) A_{\xi, x_p}^i(y - z) \left[\frac{N_c}{2} s[y - z] s[(z - \bar{y}) + \xi(y - z)] \right] \right. \\ &\quad \left. \left. + A_{\xi, x_p}^i(\bar{y} - z) A_{\xi, x_p}^i(\bar{y} - z) \frac{N_c}{2} s[z - \bar{y}] s[(y - z) - \xi(\bar{y} - z)] \right\} \right] \end{aligned}$$

$q \rightarrow q \rightarrow H$: initial expressions

- Our dilute projectile contains quarks with transverse momentum smaller than $\mu_0 \sim \Lambda_{QCD}$.
- The dense target sits at some rapidity with no need of further evolution (no large rapidity logarithms found).
- We add fragmentation and Fourier transform to transverse momentum space:

$$\frac{d\bar{\sigma}^{q \rightarrow q}}{d^2k d\eta}(k, x_p) = \frac{d\bar{\sigma}_0^{q \rightarrow q}}{d^2k d\eta}(k, x_p) + \frac{d\bar{\sigma}_{1,r}^{q \rightarrow q}}{d^2k d\eta}(k, x_p) + \frac{d\bar{\sigma}_{1,v}^{q \rightarrow q}}{d^2k d\eta}(k, x_p) \quad \frac{d\sigma^{q \rightarrow q \rightarrow H}}{d^2p d\eta} = \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \frac{d\bar{\sigma}^{q \rightarrow q}}{d^2k d\eta} \left(\frac{p}{\zeta}, \frac{x_F}{\zeta} \right)$$

$$s(k) = \int_r \frac{1}{(2\pi)^2} e^{-ik \cdot r} s(r) \implies s(r) = \int_l e^{il \cdot r} s(l) \implies s(r=0) = 1 = \int_l s(l)$$

$$\frac{d\sigma_0^{q \rightarrow q \rightarrow H}}{d^2p d\eta} = S_{\perp} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \frac{x_F}{\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta} \right) s \left(\frac{p}{\zeta} \right)$$

$q \rightarrow q \rightarrow H$: real terms

$$\frac{d\sigma_{1,r}^{q \rightarrow q \rightarrow H}}{d^2pd\eta} = S_{\perp} \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \int_{k,q} \int_{\zeta k^2/(x_F s_0)}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \frac{x_F}{(1 - \xi)\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{(1 - \xi)\zeta} \right) s(k)s(q) \frac{[k - (1 - \xi)q]^2}{\left(\frac{p}{\zeta} - k\right)^2 \left(\frac{p}{\zeta} - (1 - \xi)q\right)^2}$$

- It can be written in terms of TMD distributions plus an NLO remainder:

$$\begin{aligned} \frac{d\sigma_{1,r}^{q \rightarrow q \rightarrow H}}{d^2pd\eta} = & S_{\perp} \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \int_{k,q} \int_{\zeta k^2/(x_F s_0)}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \frac{x_F}{(1 - \xi)\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{(1 - \xi)\zeta} \right) \\ & \times s(k)s(q) \left\{ \left[\frac{\frac{p}{\zeta} - k}{\left(\frac{p}{\zeta} - k\right)^2} - \frac{\frac{p}{\zeta} - (1 - \xi)k}{\left(\frac{p}{\zeta} - (1 - \xi)k\right)^2} \right] \left[\frac{\frac{p}{\zeta} - q}{\left(\frac{p}{\zeta} - q\right)^2} - \frac{\frac{p}{\zeta} - (1 - \xi)q}{\left(\frac{p}{\zeta} - (1 - \xi)q\right)^2} \right] \right. \\ & \left. + \frac{1}{2} \left[\frac{(k - q)^2}{\left(\frac{p}{\zeta} - k\right)^2 \left(\frac{p}{\zeta} - q\right)^2} + \frac{(1 - \xi)^2 (k - q)^2}{\left(\frac{p}{\zeta} - (1 - \xi)k\right)^2 \left(\frac{p}{\zeta} - (1 - \xi)q\right)^2} \right] \right\}. \end{aligned}$$

$q \rightarrow q \rightarrow H$: real terms

$$\frac{d\sigma_{1,r}^{q \rightarrow q \rightarrow H}}{d^2pd\eta} = S_{\perp} \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \int_{k,q} \int_{\zeta k^2/(x_F s_0)}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \frac{x_F}{(1 - \xi)\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{(1 - \xi)\zeta} \right) s(k)s(q) \frac{[k - (1 - \xi)q]^2}{\left(\frac{p}{\zeta} - k\right)^2 \left(\frac{p}{\zeta} - (1 - \xi)q\right)^2}$$

- It can be written in terms of TMD distributions plus an NLO remainder:

$$\begin{aligned} \frac{d\sigma_{1,r}^{q \rightarrow q \rightarrow H}}{d^2pd\eta} = & S_{\perp} \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \int_{k,q} \int_{\zeta k^2/(x_F s_0)}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \frac{x_F}{(1 - \xi)\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{(1 - \xi)\zeta} \right) \\ & \times s(k)s(q) \left\{ \left[\frac{\frac{p}{\zeta} - k}{\left(\frac{p}{\zeta} - k\right)^2} - \frac{\frac{p}{\zeta} - (1 - \xi)k}{\left(\frac{p}{\zeta} - (1 - \xi)k\right)^2} \right] \left[\frac{\frac{p}{\zeta} - q}{\left(\frac{p}{\zeta} - q\right)^2} - \frac{\frac{p}{\zeta} - (1 - \xi)q}{\left(\frac{p}{\zeta} - (1 - \xi)q\right)^2} \right] \right. \\ & \left. + \frac{1}{2} \left[\frac{(k - q)^2}{\left(\frac{p}{\zeta} - k\right)^2 \left(\frac{p}{\zeta} - q\right)^2} + \frac{(1 - \xi)^2 (k - q)^2}{\left(\frac{p}{\zeta} - (1 - \xi)k\right)^2 \left(\frac{p}{\zeta} - (1 - \xi)q\right)^2} \right] \right\}. \end{aligned}$$

$q \rightarrow q \rightarrow H$: real terms

$$\frac{d\sigma_{1,r}^{q \rightarrow q \rightarrow H}}{d^2pd\eta} = S_{\perp} \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \int_{k,q} \int_{\zeta k^2/(x_F s_0)}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \frac{x_F}{(1 - \xi)\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{(1 - \xi)\zeta} \right) s(k)s(q) \frac{[k - (1 - \xi)q]^2}{\left(\frac{p}{\zeta} - k\right)^2 \left(\frac{p}{\zeta} - (1 - \xi)q\right)^2}$$

- It can be written in terms of TMD distributions plus an NLO remainder:

$$x\mathcal{T}_q(x, k^2; k^2; \xi_0) = \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{\xi_0}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \frac{x}{1 - \xi} f_{k^2}^q \left(\frac{x}{1 - \xi} \right) \frac{1}{k^2}$$

$$\begin{aligned} & \frac{d\sigma_{1,r}^{q \rightarrow q \rightarrow H}}{d^2pd\eta} \\ &= S_{\perp} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} \int_{k^2 > \mu_0^2} \frac{x_F}{\zeta} \left\{ D_{H,\mu_0^2}^q(\zeta) \mathcal{T}_q \left(\frac{x_F}{\zeta}, k^2; k^2, \xi_0 = k^2\zeta/(x_F s_0) \right) + f_{\mu_0^2}^q \left(\frac{x_F}{\zeta} \right) \mathcal{F}_H^q(\zeta, k^2; k^2, \xi_0 = k^2\zeta/(x_F s_0)) \right\} \\ & \quad \times s(-k + p/\zeta) \left[1 - \int_q \frac{k \cdot q}{q^2} s(-q + p/\zeta) \right] \\ &+ \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \int_k \int_{k^2\zeta/(x_F s_0)}^1 d\xi \frac{x_F}{\zeta(1 - \xi)} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta(1 - \xi)} \right) \frac{1 + (1 - \xi)^2}{\xi} \\ & \times \int_q s(k)s(q) \left[\frac{p/\zeta - k}{(p/\zeta - k)^2} - \frac{p/\zeta - (1 - \xi)k}{(p/\zeta - (1 - \xi)k)^2} \right] \left[\frac{p/\zeta - q}{(p/\zeta - q)^2} - \frac{p/\zeta - (1 - \xi)q}{(p/\zeta - (1 - \xi)q)^2} \right]. \end{aligned}$$

$q \rightarrow q \rightarrow H$: virtual terms

$$\frac{d\sigma_{1,v}^{q \rightarrow q \rightarrow H}}{d^2pd\eta} = -2 \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \int_{k^2 > \mu_0^2} \int_{k^2 \zeta / (x_F s_0)}^1 d\xi \frac{x_F}{\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta} \right) \frac{1 + (1 - \xi)^2}{\xi} \int_q s \left(\frac{p}{\zeta} \right) s(q) \left[\frac{\frac{p}{\zeta}(1 - \xi) - q - k}{(\frac{p}{\zeta}(1 - \xi) - q - k)^2} \frac{k}{k^2} + \frac{1}{k^2} \right]$$

- It contains logarithmic divergencies that can be added and subtracted, to be combined with LO providing the evolution of TMDs from μ_0^2 to μ^2 :

$$\frac{d\sigma_{1,v}^{q \rightarrow q \rightarrow H}}{d^2pd\eta} = -2 \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \int_{\mu_0^2}^{\infty} d^2k \int_{k^2 \zeta / (x_F s_0)}^1 d\xi \frac{x_F}{\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta} \right) \frac{1 + (1 - \xi)^2}{\xi} \times \int_q s \left(\frac{p}{\zeta} \right) s(q) \left[\frac{\frac{p}{\zeta}(1 - \xi) - q - k}{(\frac{p}{\zeta}(1 - \xi) - q - k)^2} \frac{k}{k^2} + \frac{1}{k^2} \right]$$

$$\pm 2 \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \frac{x_F}{\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta} \right) \int_{\mu_0^2}^{\mu^2} \frac{d^2k}{k^2} \int_{k^2 \zeta / (x_F s_0)}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \times \int_m s \left(\frac{p}{\zeta} \right) s \left(-m + \frac{p}{\zeta} \right).$$

$q \rightarrow q \rightarrow H$: virtual terms

$$\frac{d\sigma_{1,v}^{q \rightarrow q \rightarrow H}}{d^2pd\eta} = -2 \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \int_{k^2 > \mu_0^2} \int_{k^2 \zeta / (x_F s_0)}^1 d\xi \frac{x_F}{\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta} \right) \frac{1 + (1 - \xi)^2}{\xi} \int_q s \left(\frac{p}{\zeta} \right) s(q) \left[\frac{\frac{p}{\zeta}(1 - \xi) - q - k}{(\frac{p}{\zeta}(1 - \xi) - q - k)^2} \frac{k}{k^2} + \frac{1}{k^2} \right]$$

- It contains logarithmic divergencies that can be added and subtracted, to be combined with LO providing the evolution of TMDs from μ_0^2 to μ^2 :

$$x\mathcal{T}_q(x, k^2; \mu^2; \xi_0) = \theta(\mu^2 - k^2) x\mathcal{T}_q(x, k^2; k^2; \xi_0) \left[1 - \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{k^2}^{\mu^2} \frac{\pi dl^2}{l^2} \int_{\xi_0}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \right]$$

$$\begin{aligned} \frac{d\sigma_0^{q \rightarrow q \rightarrow H}}{d^2pd\eta} + \frac{d\sigma_{1,v}^{q \rightarrow q \rightarrow H}}{d^2pd\eta} &= S_{\perp} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} \int_m \int_0^{\mu_0^2} d^2l \int_0^{\mu_0^2} d^2k \\ &\times \mathcal{F}_H^q \left(\zeta, l^2; \mu^2; \xi_0 = \frac{\zeta \mu^2}{x_F s_0} \right) \frac{x_F}{\zeta} \mathcal{T}_q \left(\frac{x_F}{\zeta}, k^2; \mu^2; \xi_0 = \frac{\zeta \mu^2}{x_F s_0} \right) s \left(-(k+l) + \frac{p}{\zeta} \right) \left[1 - \frac{(k+l) \cdot m}{m^2} \right] s \left(-m + \frac{p}{\zeta} \right) \\ &- 2 \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \int_{\mu_0^2}^{\infty} d^2k \int_{k^2 \zeta / (x_F s_0)}^1 d\xi \frac{x_F}{\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta} \right) \frac{1 + (1 - \xi)^2}{\xi} \\ &\times \int_q s \left(\frac{p}{\zeta} \right) s(q) \left[\frac{\frac{p}{\zeta}(1 - \xi) - q - k}{(\frac{p}{\zeta}(1 - \xi) - q - k)^2} \frac{k}{k^2} + \frac{1}{k^2} \right] \\ &+ 2 \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \frac{x_F}{\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta} \right) \int_{\mu_0^2}^{\mu^2} \frac{d^2k}{k^2} \int_{k^2 \zeta / (x_F s_0)}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \\ &\times \int_m s \left(\frac{p}{\zeta} \right) s \left(-m + \frac{p}{\zeta} \right). \end{aligned}$$

$q \rightarrow q \rightarrow H$: virtual terms

$$\frac{d\sigma_{1,v}^{q \rightarrow q \rightarrow H}}{d^2pd\eta} = -2 \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \int_{k^2 > \mu_0^2} \int_{k^2 \zeta / (x_F s_0)}^1 d\xi \frac{x_F}{\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta} \right) \frac{1 + (1 - \xi)^2}{\xi} \int_q s \left(\frac{p}{\zeta} \right) s(q) \left[\frac{\frac{p}{\zeta}(1 - \xi) - q - k}{(\frac{p}{\zeta}(1 - \xi) - q - k)^2} \frac{k}{k^2} + \frac{1}{k^2} \right]$$

- It contains logarithmic divergencies that can be added and subtracted, to be combined with LO providing the evolution of TMDs from μ_0^2 to μ^2 :

$$x\mathcal{T}_q(x, k^2; \mu^2; \xi_0) = \theta(\mu^2 - k^2) x\mathcal{T}_q(x, k^2; k^2; \xi_0) \left[1 - \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \int_{k^2}^{\mu^2} \frac{\pi dl^2}{l^2} \int_{\xi_0}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \right]$$


$$\begin{aligned} \frac{d\sigma_0^{q \rightarrow q \rightarrow H}}{d^2pd\eta} + \frac{d\sigma_{1,v}^{q \rightarrow q \rightarrow H}}{d^2pd\eta} &= S_{\perp} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} \int_m \int_0^{\mu_0^2} d^2l \int_0^{\mu_0^2} d^2k \\ &\times \mathcal{F}_H^q \left(\zeta, l^2; \mu^2; \xi_0 = \frac{\zeta \mu^2}{x_F s_0} \right) \frac{x_F}{\zeta} \mathcal{T}_q \left(\frac{x_F}{\zeta}, k^2; \mu^2; \xi_0 = \frac{\zeta \mu^2}{x_F s_0} \right) s \left(-(k+l) + \frac{p}{\zeta} \right) \left[1 - \frac{(k+l) \cdot m}{m^2} \right] s \left(-m + \frac{p}{\zeta} \right) \\ &- 2 \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \int_{\mu_0^2}^{\infty} d^2k \int_{k^2 \zeta / (x_F s_0)}^1 d\xi \frac{x_F}{\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta} \right) \frac{1 + (1 - \xi)^2}{\xi} \\ &\times \int_q s \left(\frac{p}{\zeta} \right) s(q) \left[\frac{\frac{p}{\zeta}(1 - \xi) - q - k}{(\frac{p}{\zeta}(1 - \xi) - q - k)^2} \frac{k}{k^2} + \frac{1}{k^2} \right] \\ &+ 2 \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H,\mu_0^2}^q(\zeta) \frac{x_F}{\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta} \right) \int_{\mu_0^2}^{\mu^2} \frac{d^2k}{k^2} \int_{k^2 \zeta / (x_F s_0)}^1 d\xi \frac{1 + (1 - \xi)^2}{\xi} \\ &\times \int_m s \left(\frac{p}{\zeta} \right) s \left(-m + \frac{p}{\zeta} \right). \end{aligned}$$

they can be combined

$q \rightarrow q \rightarrow H$: all terms


- Adding all terms we get, neglecting terms $\mathcal{O}\left(\frac{p^2, k^2, Q_s^2, \mu^2}{s_0}\right), \mathcal{O}(\alpha_s^2)$:

$$\begin{aligned}
 & \frac{d\sigma^{q \rightarrow q \rightarrow H}}{d^2 p d\eta} \\
 &= S_{\perp} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} \int_m \int d^2 l \int d^2 k \\
 & \quad \times \mathcal{F}_H^q \left(\zeta, l^2; \mu^2; \xi_0 = \frac{\zeta \mu^2}{x_F s_0} \right) \frac{x_F}{\zeta} \mathcal{T}_q \left(\frac{x_F}{\zeta}, k^2; \mu^2; \xi_0 = \frac{\zeta \mu^2}{x_F s_0} \right) s \left(-(k+l) + \frac{p}{\zeta} \right) \left[1 - \frac{(k+l) \cdot m}{m^2} \right] s \left(-m + \frac{p}{\zeta} \right) \\
 & - 2 \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H, \mu_0^2}^q(\zeta) \frac{x_F}{\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta} \right) \int_m \int_{\mu^2}^{(-m+\xi p/\zeta)^2} \frac{d^2 k}{k^2} \int_{k^2 \zeta / (x_F s_0)}^1 d\xi \frac{1 + (1-\xi)^2}{\xi} \\
 & \quad \times s \left(\frac{p}{\zeta} \right) s \left(-m + \frac{p}{\zeta} \right) \\
 & + \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H, \mu_0^2}^q(\zeta) \int_k \int_{k^2 \zeta / (x_F s_0)}^1 d\xi \frac{x_F}{\zeta(1-\xi)} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta(1-\xi)} \right) \frac{1 + (1-\xi)^2}{\xi} \\
 & \quad \times \int_q s(k) s(q) \left[\frac{p/\zeta - k}{(p/\zeta - k)^2} - \frac{p/\zeta - (1-\xi)k}{(p/\zeta - (1-\xi)k)^2} \right] \left[\frac{p/\zeta - q}{(p/\zeta - q)^2} - \frac{p/\zeta - (1-\xi)q}{(p/\zeta - (1-\xi)q)^2} \right].
 \end{aligned}$$



$q \rightarrow q \rightarrow H$: all terms

- Adding all terms we get, neglecting terms $\mathcal{O}\left(\frac{p^2, k^2, Q_s^2, \mu^2}{s_0}\right), \mathcal{O}(\alpha_s^2)$:




TMDs for PDFs and FFs

$$\begin{aligned}
 & \frac{d\sigma^{q \rightarrow q \rightarrow H}}{d^2 p d\eta} \\
 &= S_{\perp} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} \int_m \int d^2 l \int d^2 k \\
 & \quad \times \mathcal{F}_H^q \left(\zeta, l^2; \mu^2; \xi_0 = \frac{\zeta \mu^2}{x_F s_0} \right) \frac{x_F}{\zeta} \mathcal{T}_q \left(\frac{x_F}{\zeta}, k^2; \mu^2; \xi_0 = \frac{\zeta \mu^2}{x_F s_0} \right) s \left(-(k+l) + \frac{p}{\zeta} \right) \left[1 - \frac{(k+l) \cdot m}{m^2} \right] s \left(-m + \frac{p}{\zeta} \right) \\
 & - 2 \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H, \mu_0^2}^q(\zeta) \frac{x_F}{\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta} \right) \int_m \int_{\mu^2}^{(-m+\xi p/\zeta)^2} \frac{d^2 k}{k^2} \int_{k^2 \zeta / (x_F s_0)}^1 d\xi \frac{1 + (1-\xi)^2}{\xi} \\
 & \quad \times s \left(\frac{p}{\zeta} \right) s \left(-m + \frac{p}{\zeta} \right) \\
 & + \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H, \mu_0^2}^q(\zeta) \int_k \int_{k^2 \zeta / (x_F s_0)}^1 d\xi \frac{x_F}{\zeta(1-\xi)} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta(1-\xi)} \right) \frac{1 + (1-\xi)^2}{\xi} \\
 & \quad \times \int_q s(k) s(q) \left[\frac{p/\zeta - k}{(p/\zeta - k)^2} - \frac{p/\zeta - (1-\xi)k}{(p/\zeta - (1-\xi)k)^2} \right] \left[\frac{p/\zeta - q}{(p/\zeta - q)^2} - \frac{p/\zeta - (1-\xi)q}{(p/\zeta - (1-\xi)q)^2} \right].
 \end{aligned}$$

$q \rightarrow q \rightarrow H$: all terms

- Adding all terms we get, neglecting terms $\mathcal{O}\left(\frac{p^2, k^2, Q_s^2, \mu^2}{s_0}\right), \mathcal{O}(\alpha_s^2)$:



$$\frac{d\sigma^{q \rightarrow q \rightarrow H}}{d^2 p d\eta}$$

$$= S_{\perp} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} \int_m \int d^2 l \int d^2 k$$

TMDs for PDFs and FFs

$$\times \mathcal{F}_H^q \left(\zeta, l^2; \mu^2; \xi_0 = \frac{\zeta \mu^2}{x_F s_0} \right) \frac{x_F}{\zeta} \mathcal{T}_q \left(\frac{x_F}{\zeta}, k^2; \mu^2; \xi_0 = \frac{\zeta \mu^2}{x_F s_0} \right) s \left(-(k+l) + \frac{p}{\zeta} \right) \left[1 - \frac{(k+l) \cdot m}{m^2} \right] s \left(-m + \frac{p}{\zeta} \right)$$

$$-2 \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H, \mu_0^2}^q(\zeta) \frac{x_F}{\zeta} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta} \right) \int_m \int_{\mu^2}^{(-m+\xi p/\zeta)^2} \frac{d^2 k}{k^2} \int_{k^2 \zeta / (x_F s_0)}^1 d\xi \frac{1 + (1-\xi)^2}{\xi}$$

$$\times s \left(\frac{p}{\zeta} \right) s \left(-m + \frac{p}{\zeta} \right)$$

NLO remainders, no large logs for our choice of scales

$$+ \frac{g^2}{(2\pi)^3} S_{\perp} \frac{N_c}{2} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} D_{H, \mu_0^2}^q(\zeta) \int_k \int_{k^2 \zeta / (x_F s_0)}^1 d\xi \frac{x_F}{\zeta(1-\xi)} f_{\mu_0^2}^q \left(\frac{x_F}{\zeta(1-\xi)} \right) \frac{1 + (1-\xi)^2}{\xi}$$

$$\times \int_q s(k) s(q) \left[\frac{p/\zeta - k}{(p/\zeta - k)^2} - \frac{p/\zeta - (1-\xi)k}{(p/\zeta - (1-\xi)k)^2} \right] \left[\frac{p/\zeta - q}{(p/\zeta - q)^2} - \frac{p/\zeta - (1-\xi)q}{(p/\zeta - (1-\xi)q)^2} \right].$$

$q \rightarrow q \rightarrow H$: final expression

- Neglecting terms $\mathcal{O}\left(\frac{p^2, k^2, Q_s^2, \mu^2}{s_0}\right)$, $\mathcal{O}(\alpha_s^2)$, we get a parton model-like expression:

$$\frac{d\sigma^{q \rightarrow q \rightarrow H}}{d^2p d\eta} = S_{\perp} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} \int d\xi \int d^2l \int d^2k \mathcal{F}_H^q \left(\zeta, l^2; \mu^2; \xi_0 = \frac{\zeta \mu^2}{x_F s_0} \right) \times \frac{x_F}{\zeta(1-\xi)} \mathcal{T}_q \left(\frac{x_F}{\zeta(1-\xi)}, k^2; \mu^2; \xi_0 = \frac{\zeta \mu^2}{x_F s_0} \right) \mathcal{P}(\xi, \zeta; k+l; p, s_0, \mu^2, \mu_0^2),$$

$$\begin{aligned} \mathcal{P}(\xi, \zeta; k+l; p, s_0, \mu^2, \mu_0^2) = & \int d\lambda \int_m \left\{ \delta(\lambda) \delta(\xi - \lambda) s \left(-(k+l) + \frac{p}{\zeta} \right) \left[1 - \frac{(k+l) \cdot m}{m^2} \right] s \left(-m + \frac{p}{\zeta} \right) \right. \\ & + \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \frac{1 + (1-\lambda)^2}{\lambda} \theta(1-\lambda) \\ & \left[\delta(\lambda - \xi) \theta \left(\xi - \frac{m^2 \zeta}{x_F s_0} \right) \int_q s(m) s(q) \left[\frac{p/\zeta - m}{(p/\zeta - m)^2} - \frac{p/\zeta - (1-\xi)m}{(p/\zeta - (1-\xi)m)^2} \right] \left[\frac{p/\zeta - q}{(p/\zeta - q)^2} - \frac{p/\zeta - (1-\xi)q}{(p/\zeta - (1-\xi)q)^2} \right] \right. \\ & \left. \left. - 2\delta(\xi) \theta \left(\lambda - \frac{\mu^2 \zeta}{x_F s_0} \right) \theta(m^2 - \mu_0^2) s \left(\frac{p}{\zeta} \right) s \left(m + (1-\lambda) \frac{p}{\zeta} \right) \int_{\mu^2}^{\min[m^2, \lambda \bar{s}_0]} \frac{d^2q}{q^2} \right] \right\}. \end{aligned}$$

$q \rightarrow q \rightarrow H$: final expression

- Neglecting terms $\mathcal{O}\left(\frac{p^2, k^2, Q_s^2, \mu^2}{s_0}\right)$, $\mathcal{O}(\alpha_s^2)$, we get a parton model-like expression:

$$\frac{d\sigma^{q \rightarrow q \rightarrow H}}{d^2p d\eta} = S_{\perp} \int_{x_F}^1 \frac{d\zeta}{\zeta^2} \int d\xi \int d^2l \int d^2k \mathcal{F}_H^q \left(\zeta, l^2; \mu^2; \xi_0 = \frac{\zeta \mu^2}{x_F s_0} \right) \times \frac{x_F}{\zeta(1-\xi)} \mathcal{T}_q \left(\frac{x_F}{\zeta(1-\xi)}, k^2; \mu^2; \xi_0 = \frac{\zeta \mu^2}{x_F s_0} \right) \mathcal{P}(\xi, \zeta; k+l; p, s_0, \mu^2, \mu_0^2),$$

$$\mathcal{P}(\xi, \zeta; k+l; p, s_0, \mu^2, \mu_0^2) = \int d\lambda \int_m \left\{ \delta(\lambda) \delta(\xi - \lambda) s \left(-(k+l) + \frac{p}{\zeta} \right) \left[1 - \frac{(k+l) \cdot m}{m^2} \right] s \left(-m + \frac{p}{\zeta} \right) + \frac{g^2}{(2\pi)^3} \frac{N_c}{2} \frac{1 + (1-\lambda)^2}{\lambda} \theta(1-\lambda) \left[\delta(\lambda - \xi) \theta \left(\xi - \frac{m^2 \zeta}{x_F s_0} \right) \int_q s(m) s(q) \left[\frac{p/\zeta - m}{(p/\zeta - m)^2} - \frac{p/\zeta - (1-\xi)m}{(p/\zeta - (1-\xi)m)^2} \right] \left[\frac{p/\zeta - q}{(p/\zeta - q)^2} - \frac{p/\zeta - (1-\xi)q}{(p/\zeta - (1-\xi)q)^2} \right] - 2\delta(\xi) \theta \left(\lambda - \frac{\mu^2 \zeta}{x_F s_0} \right) \theta(m^2 - \mu_0^2) s \left(\frac{p}{\zeta} \right) s \left(m + (1-\lambda) \frac{p}{\zeta} \right) \int_{\mu^2}^{\min[m^2, \lambda \bar{s}_0]} \frac{d^2q}{q^2} \right] \right\}.$$

quark scattering
qg scattering due to $q \rightarrow qg$

The other channels:

- TMD PDFs:

- For quark: it gets contributions from $q \rightarrow q$ and $g \rightarrow q$.
- For antiquark: it gets contributions from $\bar{q} \rightarrow \bar{q}$ and $g \rightarrow \bar{q}$.
- For gluon: it gets contributions from $g \rightarrow g$, $q \rightarrow g$ and $\bar{q} \rightarrow g$.

- TMD FFs:

- For quark: it gets contributions from $q \rightarrow q \rightarrow H$ and $q \rightarrow g \rightarrow H$.
- For antiquark: it gets contributions from $\bar{q} \rightarrow \bar{q} \rightarrow H$ and $\bar{q} \rightarrow g \rightarrow H$.
- For gluon: it gets contributions from $g \rightarrow g \rightarrow H$, $g \rightarrow q \rightarrow H$ and $g \rightarrow \bar{q} \rightarrow H$.

- The **complete quark piece** of the parton-like formula keeps the form with additional NLO remainders.

- The **gluon piece of the parton-like formula contains 3 dipoles in the fundamental representation**, and additional NLO remainders (in progress).

Summary:

- In view of the negativity problem and the recent LHCb data, **we revisit the calculation of single inclusive particle production at NLO in the hybrid model**, avoiding collinear and soft (BK-like) subtractions. We assume large N_c , and factorisation/translational invariance for dipoles.
- We see that divergencies are absorbed into TMD PDFs and FFs defined from collinear ones.
- We get a parton model-like formula with a probability interpretation plus NLO, not log-enhanced remainders.

$$\int \frac{d\zeta}{\zeta^2} \int d^2k_{\perp} d^2q_{\perp} T\left(\frac{x_F}{\zeta}, k_{\perp}; \mu_T^2\right) P(k_{\perp}, q_{\perp}) F(\zeta; p_{\perp}, (k_{\perp} + q_{\perp}); \mu_F^2) + \text{NLO remainders} + \mathcal{O}\left(\frac{p_{\perp}^2, k_{\perp}^2, Q_s^2, \mu^2}{s_0}\right)$$

- **We conclude that the correct framework to resum all large logarithms is not collinear factorization for the projectile (the hybrid model) but TMD factorization.**
- **Outlook:** → Complete the calculation for all channels and provide the final expressions.
→ Relate our definition of TMD PDFs and FFs and their evolution equations with standard ones (relation of our rapidity cut-off with other choices).
→ Implement numerically our results.

Summary:

- In view of the ~~negativity problem and the recent LHCb data~~ **we revisit the calculation of single inclusive** *Thanks a lot to you for your attention and to the ECT* people for the organisation!!!* **and soft (BK-like) subtraction** or dipoles.

- We see that divergencies are absorbed into TMD PDFs and FFs defined from collinear ones.
- We get a parton model-like formula with a probability interpretation plus NLO, not log-enhanced remainders.

$$\int \frac{d\zeta}{\zeta^2} \int d^2k_{\perp} d^2q_{\perp} T\left(\frac{x_F}{\zeta}, k_{\perp}; \mu_T^2\right) P(k_{\perp}, q_{\perp}) F(\zeta; p_{\perp}, (k_{\perp} + q_{\perp}); \mu_F^2) + \text{NLO remainders} + \mathcal{O}\left(\frac{p_{\perp}^2, k_{\perp}^2, Q_s^2, \mu^2}{s_0}\right)$$

- **We conclude that the correct framework to resum all large logarithms is not collinear factorization for the projectile (the hybrid model) but TMD factorization.**
- **Outlook:** → Complete the calculation for all channels and provide the final expressions.
→ Relate our definition of TMD PDFs and FFs and their evolution equations with standard ones (relation of our rapidity cut-off with other choices).
→ Implement numerically our results.

Backup: