

RECENT UPDATES TO nCTEQ NUCLEAR PDFs

Pit Duwentäster



Short-Distance nuclear structure and PDFs, Trento, Italy


July 20, 2023

TABLE OF CONTENTS

- ▶ Basics of nuclear PDFs and nCTEQ15
- ▶ nCTEQ15WZ - Vector boson production
- ▶ nCTEQ15WZ+SIH - Single inclusive hadron production
- ▶ nCTEQ15HQ - Heavy quark production
- ▶ nCTEQ15HIX - High- x DIS data
- ▶ nCTEQ15 ν - Neutrino DIS data
- ▶ nCTEQ23 - Road to future nPDFs
- ▶ Conclusions

INTRODUCTION - NUCLEAR PDFs

Nuclear PDFs are more than the sum of their parts, i.e. not just the sum of Z proton PDFs and $(A - Z)$ neutron PDFs.



PDF FITS - PARAMETERIZATION

- ▶ The full nPDFs are parametrized in terms of the bound nucleon PDFs:

$$f_i^{A,Z}(x, Q) = \frac{Z}{A} f_i^{p/A}(x, Q) + \frac{A-Z}{A} f_i^{n/A}(x, Q)$$

PDF FITS - PARAMETERIZATION

- ▶ The full nPDFs are parametrized in terms of the bound nucleon PDFs:

$$f_i^{A,Z}(x, Q) = \frac{Z}{A} f_i^{p/A}(x, Q) + \frac{A-Z}{A} f_i^{n/A}(x, Q)$$

- ▶ Bound proton PDF parametrizations at $Q_0 = 1.3 \text{ GeV}$:

$$x f_i^{p/A}(x, Q_0) = c_0 x^{c_1} (1-x)^{c_2} e^{c_3 x} (1 + e^{c_4 x})^{c_5}, \quad i = u_v, d_v, g, \bar{u} + \bar{d}, s + \bar{s}$$
$$\frac{\bar{d}}{\bar{u}} = c_0 x^{c_1} (1-x)^{c_2} + (1 + c_3 x)(1-x)^{c_4}$$

PDF FITS - PARAMETERIZATION

- ▶ The full nPDFs are parametrized in terms of the bound nucleon PDFs:

$$f_i^{A,Z}(x, Q) = \frac{Z}{A} f_i^{p/A}(x, Q) + \frac{A-Z}{A} f_i^{n/A}(x, Q)$$

- ▶ Bound proton PDF parametrizations at $Q_0 = 1.3$ GeV :

$$x f_i^{p/A}(x, Q_0) = c_0 x^{c_1} (1-x)^{c_2} e^{c_3 x} (1 + e^{c_4 x})^{c_5}, \quad i = u_v, d_v, g, \bar{u} + \bar{d}, s + \bar{s}$$
$$\frac{\bar{d}}{\bar{u}} = c_0 x^{c_1} (1-x)^{c_2} + (1 + c_3 x)(1-x)^{c_4}$$

- ▶ The dependence of nPDFs on A, Z is parametrized as

$$c_k(A, Z) = p_k + a_k(1 - A^{-b_k})$$

PDF FITS - PARAMETERIZATION

- ▶ The full nPDFs are parametrized in terms of the bound nucleon PDFs:

$$f_i^{A,Z}(x, Q) = \frac{Z}{A} f_i^{p/A}(x, Q) + \frac{A-Z}{A} f_i^{n/A}(x, Q)$$

- ▶ Bound proton PDF parametrizations at $Q_0 = 1.3$ GeV :

$$x f_i^{p/A}(x, Q_0) = c_0 x^{c_1} (1-x)^{c_2} e^{c_3 x} (1 + e^{c_4 x})^{c_5}, \quad i = u_v, d_v, g, \bar{u} + \bar{d}, s + \bar{s}$$
$$\frac{\bar{d}}{\bar{u}} = c_0 x^{c_1} (1-x)^{c_2} + (1 + c_3 x)(1-x)^{c_4}$$

- ▶ The dependence of nPDFs on A, Z is parametrized as


$$c_k(A, Z) = p_k + a_k(1 - A^{-b_k})$$

- ▶ Open parameters:

$$\{a_1^{u_v}, a_2^{u_v}, a_4^{u_v}, a_5^{u_v}, a_1^{d_v}, a_2^{d_v}, a_5^{d_v}, a_1^{\bar{u}+\bar{d}}, a_5^{\bar{u}+\bar{d}}, a_1^g, a_4^g, a_5^g, b_0^g, b_1^g, b_4^g, b_5^g, a_0^{s+\bar{s}}, a_1^{s+\bar{s}}, a_2^{s+\bar{s}}\}$$


INTRODUCTION - THE nCTEQ15 BASELINE

Old main release: **nCTEQ15** [Kovarik et al., PRD 93 (2016) 085037]




INTRODUCTION - THE nCTEQ15 BASELINE

Old main release: **nCTEQ15** [Kovarik et al., PRD 93 (2016) 085037]



Many open problems:

- ▶ Gluon - huge uncertainties at low- x
- ▶ Strange quark - fixed to a constant fraction of $\bar{u} + \bar{d}$
- ▶ Considerable uncertainties even for valence quarks at high- x




nCTEQ15HIX - HIGH- x JLAB DATA

[Segarra et al., PRD 103 (2021) 114015]

Goal: Lower kinematic cuts to include more data and constrain high- x PDFs

- ▶ **Old:** $Q_{\min} = 2$ GeV and $W_{\min} = 3.5$ GeV
 - ▶ 708 DIS data points
 - ▶ JLab fully excluded




nCTEQ15HIX - HIGH- x JLAB DATA

[Segarra et al., PRD 103 (2021) 114015]

Goal: Lower kinematic cuts to include more data and constrain high- x PDFs


- ▶ **Old:** $Q_{\min} = 2$ GeV and $W_{\min} = 3.5$ GeV
 - ▶ 708 DIS data points
 - ▶ JLab fully excluded
- ▶ **New:** $Q_{\min} = 1.3$ GeV and $W_{\min} = 1.7$ GeV
 - ▶ 1564 DIS data points
 - ▶ JLab and HERMES contribute significant amounts
 - ▶ Requires low-energy corrections



Target mass corrections:

[Ruiz et al., arXiv:2301.07715]

- Revisited derivation for nuclear DIS




Target mass corrections:

[Ruiz et al., arXiv:2301.07715]

- ▶ Revisited derivation for nuclear DIS

Higher twist effects:

- ▶ Independent of the nucleus
- ▶ Using parameterization from the CJ15 analysis [Accardi et al, PRD 93 (2016) 114017]



Target mass corrections:

[Ruiz et al., arXiv:2301.07715]



- ▶ Revisited derivation for nuclear DIS

Higher twist effects:

- ▶ Independent of the nucleus
- ▶ Using parameterization from the CJ15 analysis [Accardi et al, PRD 93 (2016) 114017]


Deuteron corrections:

- ▶ Lightest bound nucleus
- ▶ Considerable differences to heavier targets
- ▶ Corrections have been extracted in the CJ15 analysis



nCTEQ15HIX - RESULTING PDFs

Iron PDF ratios to nCTEQ15 ($Q = 2$ GeV)




- ▶ **BASE**: nCTEQ15 with relaxed cuts + TMC
- ▶ **HT**: BASE + higher twist corrections
- ▶ **DEU**: BASE + deuteron corrections
- ▶ **nCTEQ15HIX**: BASE + higher twist and deuteron corrections

[Kusina et al., EPJC 80 (2020) 968]

Goal: Constrain the strange quark (and gluon) PDFs

- ▶ Open up strange quark parameters
- ▶ Strange quark channel contributes up to 30% of the total cross section
- ▶ 120 new data points

Data Overview						
			$\sqrt{s_{NN}}$ [TeV]	Norm σ	No Points	Ref.
ATLAS	Run I	W^\pm	5.02	2.7%	10+10	[49]
ATLAS	Run I	Z	5.02	2.7%	14	[50]
CMS	Run I	W^\pm	5.02	3.5%	10+10	[51]
CMS	Run I	Z	5.02	3.5%	12	[52]
CMS	Run II	W^\pm	8.16	3.5%	24+24	[53]
ALICE	Run I	W^\pm	5.02	2.0%	2+2	[54, 55]
LHCb	Run I	Z	5.02	2.0%	2	[56]



n CTEQ15WZ+SIH - SINGLE INCLUSIVE HADRON PRODUCTION

[Duwentäster et al., PRD 104 (2021) 094005]

Goal: Constrain the gluon PDFs



- ▶ Sensitivity to gluon PDFs not only from DGLAP evolution
- ▶ Precise new data from ALICE

NCTEQ15WZ+SIH - SINGLE INCLUSIVE HADRON PRODUCTION

[Duwentäster et al., PRD 104 (2021) 094005]

Goal: Constrain the gluon PDFs

- ▶ Sensitivity to gluon PDFs not only from DGLAP evolution
- ▶ Precise new data from ALICE





NCTEQ15WZ+SIH - SINGLE INCLUSIVE HADRON PRODUCTION

[Duwentäster et al., PRD 104 (2021) 094005]

Goal: Constrain the gluon PDFs


- ▶ Sensitivity to gluon PDFs not only from DGLAP evolution
- ▶ Precise new data from ALICE



Challenge: Fragmentation function dependence

- ▶ The most precise data lies in non-perturbative low p_T region
- ▶ Fragmentation Function dependence

- **FF dependence:** Boils down to normalization shift \rightarrow cancels in ratio





NCTEQ15WZ+SIH - FF DEPENDENCE

► **FF dependence:** Boils down to normalization shift \rightarrow cancels in ratio

► **Scale dependence:**

$\mu = \mu = \mu = \frac{1}{2}p_T$ gives optimal description





- ▶ **FF dependence:** Boils down to normalization shift \rightarrow cancels in ratio

- ▶ **Scale dependence:**

$\mu = \mu = \mu = \frac{1}{2}p_T$ gives optimal description



- ▶ **Caveat:** No combination of FF and scale choice can describe data with $p_T < 3$ GeV



nCTEQ15HQ - MOTIVATION

[Duwentäster et al., PRD 105 (2022) 114043]

Why is heavy-flavour production data interesting for nuclear PDFs?





nCTEQ15HQ - MOTIVATION

[Duwentäster et al., PRD 105 (2022) 114043]

Why is heavy-flavour production data interesting for nuclear PDFs?

- ▶ Sensitivity to gluon PDFs down to very low x values ($x \approx 10^{-5}$)
- ▶ Large available proton-lead data sets from multiple LHC experiments for D^0 , J/ψ , $\psi(2S)$ and $\Upsilon(1S)$
- ▶ Interesting data-driven approach [A. Kusina et al., PRL 121 (2018) 052004; PRD 104 (2021) 014010]





nCTEQ15HQ - MOTIVATION

[Duwentäster et al., PRD 105 (2022) 114043]

Why is heavy-flavour production data interesting for nuclear PDFs?

- ▶ Sensitivity to gluon PDFs down to very low x values ($x \approx 10^{-5}$)
- ▶ Large available proton-lead data sets from multiple LHC experiments for D^0 , J/ψ , $\psi(2S)$ and $\Upsilon(1S)$
- ▶ Interesting data-driven approach [A. Kusina et al., PRL 121 (2018) 052004; PRD 104 (2021) 014010]
 - ▶ Understanding of quarkonium production in pQCD is limited
 - ▶ Can quantify theory uncertainties
 - ▶ Potentially applicable for many single-inclusive particle production processes
 - ▶ Fast calculation



nCTEQ15HQ - DATA-DRIVEN APPROACH


Derive effective theory from p-p collision data to use with p-Pb data in the nPDF fit

$$\sigma(AB \rightarrow Q + X) = \int dx_1 dx_2 f_{1,g}(x_1) f_{2,g}(x_2) \frac{1}{2\hat{s}} \overline{|\mathcal{A}_{gg \rightarrow Q + X}|^2} dPS$$

NCTEQ15HQ - DATA-DRIVEN APPROACH

Derive effective theory from p-p collision data to use with p-Pb data in the nPDF fit


$$\sigma(AB \rightarrow Q + X) = \int dx_1 dx_2 f_{1,g}(x_1) f_{2,g}(x_2) \frac{1}{2\hat{s}} \overline{|\mathcal{A}_{gg \rightarrow Q+X}|^2} dPS$$



NCTEQ15HQ - DATA-DRIVEN APPROACH

Derive effective theory from p-p collision data to use with p-Pb data in the nPDF fit

$$\sigma(AB \rightarrow Q + X) = \int dx_1 dx_2 f_{1,g}(x_1) f_{2,g}(x_2) \frac{1}{2\hat{s}} \overline{|\mathcal{A}_{gg \rightarrow Q+X}|^2} \text{dPS}$$




$$\overline{|\mathcal{A}_{gg \rightarrow Q+X}|^2} = \frac{\lambda^2 \kappa \hat{s}}{M_Q^2} \begin{cases} e^{-\kappa \frac{p_T^2}{M_Q^2} + a|y|} & \text{if } p_T \leq \langle p_T \rangle \\ e^{-\kappa \frac{\langle p_T \rangle^2}{M_Q^2} + a|y|} \left(1 + \frac{\kappa}{n} \frac{p_T^2 - \langle p_T \rangle^2}{M_Q^2}\right)^{-n} & \text{if } p_T > \langle p_T \rangle \end{cases}$$

NCTEQ15HQ - DATA-DRIVEN APPROACH

Derive effective theory from p-p collision data to use with p-Pb data in the nPDF fit


$$\sigma(AB \rightarrow Q + X) = \int dx_1 dx_2 f_{1,g}(x_1) f_{2,g}(x_2) \frac{1}{2\hat{s}} \overline{|\mathcal{A}_{gg \rightarrow Q+X}|^2} \text{dPS}$$



$$\overline{|\mathcal{A}_{gg \rightarrow Q+X}|^2} = \frac{\lambda^2 \kappa \hat{s}}{M_Q^2} \begin{cases} e^{-\kappa \frac{p_T^2}{M_Q^2} + a|y|} & \text{if } p_T \leq \langle p_T \rangle \\ e^{-\kappa \frac{\langle p_T \rangle^2}{M_Q^2} + a|y|} \left(1 + \frac{\kappa}{n} \frac{p_T^2 - \langle p_T \rangle^2}{M_Q^2}\right)^{-n} & \text{if } p_T > \langle p_T \rangle \end{cases}$$

- **Fit parameters** → uncertainties can be determined just like those of PDFs
- 5 parameters per process → 6×5 parameters describe ≈ 1000 data points very well
- Agrees well with NRQCD predictions for J/ψ and GMVFNS predictions for D^0

nCTEQ15(WZ, WZ+SIH, HQ) - RESULTING PDFs



- ▶ Significant reduction in gluon and low- x uncertainties
- ▶ Strange quark still poorly constrained

NCTEQ15 ν - NEUTRINO DIS REVISITED

[Muzakka et al., PRD 106 (2022) 074004]

Motivation:

- ▶ Reduce the uncertainty of the strange PDF
- ▶ Improve flavour separation
- ▶ **HUGE** data sets

Neutrino Data used in this analysis:

Data set	Nucleus	#pts	Corr.sys.
CDHSW ν	Fe	465	No
CDHSW $\bar{\nu}$		464	
CCFR ν	Fe	1109	No
CCFR $\bar{\nu}$		1098	
NuTeV ν	Fe	1170	Yes
NuTeV $\bar{\nu}$		966	
Chorus ν	Pb	412	Yes
Chorus $\bar{\nu}$		412	
CCFR dimuon ν	Fe	40	No
CCFR dimuon $\bar{\nu}$		38	
NuTeV dimuon ν	Fe	38	No
NuTeV dimuon $\bar{\nu}$		34	

Motivation:

- ▶ Reduce the uncertainty of the strange PDF
- ▶ Improve flavour separation
- ▶ **HUGE** data sets

Challenge:

- ▶ Tensions between ν -DIS data sets
- ▶ Tensions within ν -DIS data sets
- ▶ Tensions between ν -DIS and other data

Neutrino Data used in this analysis:

Data set	Nucleus	#pts	Corr.sys.
CDHSW ν	Fe	465	No
CDHSW $\bar{\nu}$		464	
CCFR ν	Fe	1109	No
CCFR $\bar{\nu}$		1098	
NuTeV ν	Fe	1170	Yes
NuTeV $\bar{\nu}$		966	
Chorus ν	Pb	412	Yes
Chorus $\bar{\nu}$		412	
CCFR dimuon ν	Fe	40	No
CCFR dimuon $\bar{\nu}$		38	
NuTeV dimuon ν	Fe	38	No
NuTeV dimuon $\bar{\nu}$		34	

Motivation:

- ▶ Reduce the uncertainty of the strange PDF
- ▶ Improve flavour separation
- ▶ **HUGE** data sets

Challenge:

- ▶ Tensions between ν -DIS data sets
- ▶ Tensions within ν -DIS data sets
- ▶ Tensions between ν -DIS and other data


Improved methodology:

- ▶ Treatment of normalization uncertainties.
- ▶ Deuteron correction from CJ15 analysis.
- ▶ More PDF parameters to fit.
- ▶ New tools for compatibility analysis

Neutrino Data used in this analysis:

Data set	Nucleus	#pts	Corr.sys.
CDHSW ν	Fe	465	No
CDHSW $\bar{\nu}$		464	
CCFR ν	Fe	1109	No
CCFR $\bar{\nu}$		1098	
NuTeV ν	Fe	1170	Yes
NuTeV $\bar{\nu}$		966	
Chorus ν	Pb	412	Yes
Chorus $\bar{\nu}$		412	
CCFR dimuon ν	Fe	40	No
CCFR dimuon $\bar{\nu}$		38	
NuTeV dimuon ν	Fe	38	No
NuTeV dimuon $\bar{\nu}$		34	

NCTEQ15 ν - NEW FITS



- ▶ **nCTEQ15WZSIHdeut** = Base: l^\pm DIS+DY+SIH+WZ (940 pts)
- ▶ **BaseDimuNeuX**: Base + Dimuon + Neutrino, with $x \leq 0.1$ data are removed from the dimuon and neutrino data (5584 pts)
- ▶ **BaseDimuChorus**: Base + Dimuon + Chorus (1914 pts)

Combine nCTEQ15HIX, nCTEQ15HQ and nCTEQ15 ν

- ▶ Match cuts
- ▶ Match open parameters
- ▶ Check consistency between all data sets

Combine nCTEQ15HIX, nCTEQ15HQ and nCTEQ15 ν

- ▶ Match cuts
- ▶ Match open parameters
- ▶ Check consistency between all data sets

Update data

- ▶ New W/Z data from ALICE and LHCb
- ▶ New π^0 data from LHCb
- ▶ New Heavy quark data from all ALICE, CMS and LHCb

Combine nCTEQ15HIX, nCTEQ15HQ and nCTEQ15 ν

- ▶ Match cuts
- ▶ Match open parameters
- ▶ Check consistency between all data sets

Update data

- ▶ New W/Z data from ALICE and LHCb
- ▶ New π^0 data from LHCb
- ▶ New Heavy quark data from all ALICE, CMS and LHCb

New observables (under investigation)

- ▶ Dijet production
- ▶ Direct photon production

Updated parameterization

Based on the new CJ22 proton analysis [Accardi et al., arXiv:2303.11509]

$$xf_i(x, Q_0^2) = c_0 x^{c_1} (1-x)^{c_2} \left(1 + c_3 \sqrt{x} + c_5 \sqrt{x^3}\right) \quad i = u_v, g, \bar{u} + \bar{d}, \bar{s} + \bar{s}$$

$$xd_v(x, Q_0^2) = c_0 x^{c_1} (1-x)^{c_2} \left(1 + c_3 \sqrt{x} + c_5 \sqrt{x^3}\right) + c_6 x^{c_7} u_v(x, Q_0^2)$$

$$x(\bar{d} - \bar{u}) = c_0 x^{c_1} (1-x)^{c_2} (1 + c_3 x)$$

Updated parameterization

Based on the new CJ22 proton analysis [Accardi et al., arXiv:2303.11509]

$$xf_i(x, Q_0^2) = c_0 x^{c_1} (1-x)^{c_2} \left(1 + c_3 \sqrt{x} + c_5 \sqrt{x^3}\right) \quad i = u_v, g, \bar{u} + \bar{d}, \bar{s} + \bar{c}$$

$$xd_v(x, Q_0^2) = c_0 x^{c_1} (1-x)^{c_2} \left(1 + c_3 \sqrt{x} + c_5 \sqrt{x^3}\right) + c_6 x^{c_7} u_v(x, Q_0^2)$$

$$x(\bar{d} - \bar{u}) = c_0 x^{c_1} (1-x)^{c_2} (1 + c_3 x)$$

Changing the nuclear A dependence

$$\text{OLD:} \quad c_k \rightarrow c_k(A) \equiv p_k + a_k (1 - A^{-b_k})$$

$$\Downarrow$$

$$\text{NEW:} \quad c_k \rightarrow c_k(A) \equiv p_k + a_k \ln(A) + b_k \ln(A)^2$$

SUMMARY

Modern nuclear PDF analyses require a vast number of different components to come together to gain comprehensive understanding

DATA

- ▶ Vector bosons
- ▶ Inclusive hadrons
- ▶ Heavy quarks
- ▶ Neutrino DIS
- ▶ High- x DIS
- ▶ (Direct photons)
- ▶ (Dijets)

SUMMARY

Modern nuclear PDF analyses require a vast number of different components to come together to gain comprehensive understanding

DATA

- ▶ Vector bosons
- ▶ Inclusive hadrons
- ▶ Heavy quarks
- ▶ Neutrino DIS
- ▶ High- x DIS
- ▶ (Direct photons)
- ▶ (Dijets)

THEORY

- ▶ Target mass corrections
- ▶ Higher twist corrections
- ▶ Deuteron treatment
- ▶ FF dependence of single inclusive hadrons
- ▶ Data-driven approach for HQ

SUMMARY

Modern nuclear PDF analyses require a vast number of different components to come together to gain comprehensive understanding

DATA

- ▶ Vector bosons
- ▶ Inclusive hadrons
- ▶ Heavy quarks
- ▶ Neutrino DIS
- ▶ High- x DIS
- ▶ (Direct photons)
- ▶ (Dijets)

THEORY

- ▶ Target mass corrections
- ▶ Higher twist corrections
- ▶ Deuteron treatment
- ▶ FF dependence of single inclusive hadrons
- ▶ Data-driven approach for HQ


TECHNICAL

- ▶ CJ22 proton baseline
- ▶ A -dependence parameterization
- ▶ Compatibility criteria
- ▶ Increasing number of open parameters

Backup slides



CURRENT STATUS OF OTHER NPDF FITS

	N_{data}	N_{params}	Observables
nCTEQ15WZSIH	948	19	DIS, DY, SIH, WZ
EPPS21	2077	24	(ν) DIS, DY, SIH, WZ, dijet, D
nNNPDF3.0	2151	256	(ν) DIS, DY, WZ, dijet, γ , D





Many others exist: KSASG20, TUJU21, DSSZ,...

BACKUP - z DEPENDENCE











BACKUP - FF PARTON DEPENDENCE






HEAVY QUARKS - PROTON-PROTON BASELINE

► Impose cuts to remove data with $p_T < 3 \text{ GeV}$ and outside of $-4 \leq y_{\text{c.m.s.}} \leq 4$

	D^0	J/ψ	$B \rightarrow J/\psi$	$\Upsilon(1S)$	$\psi(2S)$	$B \rightarrow \psi(2S)$
N_{points}	34	501		375	55	
χ^2/N_{dof}	0.25	0.88		0.92	0.77	






BASELINE - COMPARISON WITH GMVFNS



- ▶ KKKS08 fragmentation functions
- ▶ Base scale $\mu_r = \mu_i = \mu_f = \sqrt{p_T^2 + 4m_c^2}$
- ▶ Uncertainties due to individual scale variations by factor 2 or $\frac{1}{2}$

BASELINE - COMPARISON WITH NRQCD

Calculations by Mathias Butenschoen, Bernd Kniehl [M. Butenschoen et al., Nucl.Phys.B Proc.Suppl. 222-224 (2012) 151-161]




- NRQCD Uncertainties due to scale variations:

$$1/2 < \mu_r/\mu_{r,0} = \mu_f/\mu_{f,0} = \mu_{NRQCD}/\mu_{NRQCD,0} < 2$$


- Base scale $\mu_{r,0} = \mu_{f,0} = \sqrt{p_T^2 + 4m_c^2}$ and $m_{NRQCD,0} = m_c$

PDF FITS - DATA SELECTION AND SETTINGS

- ▶ Include all data from nCTEQ15WZ+SIH
- ▶ Use the same open parameters as nCTEQ15WZ+SIH
- ▶ Cut HQ data below $p_T < 3.0$ GeV and outside $-4 \leq y_{\text{c.m.s.}} \leq 4$
 - ▶ Same as proton-proton baseline.
 - ▶ 548 new data points
 - ▶ Add Crystal Ball uncertainty to data systematics



nCTEQ15HQ PDFs - SCALE VARIATION



COMPATIBILITY ANALYSIS S vs \bar{S}

DEFINITION ($\Delta\chi_S^2$ -COMPATIBILITY)

$$\chi_S^2(\text{after}) - \chi_S^2(\text{before}) \leq \Delta\chi_S^2$$

DEFINITION (χ^2 -COMPATIBILITY)



$$p_S \leq 0.9 \quad \text{and} \quad p_{\bar{S}} \leq 0.9$$

$$p_S = \int_0^{\chi_S^2} P(\chi^2, N_S) d\chi^2$$

DEFINITION (S_E -COMPATIBILITY)

$$S_E = \sqrt{2\chi_E^2(N)} - \sqrt{2N-1} \sim \mathcal{N}(0, 1)$$

TREATMENT OF NORMALIZATION PARAMETERS



- ▶ The old way to include normalization uncertainties :

$$\chi^2 = (D - T)^T C_D^{-1} (D - T),$$

$$C_{D,ij} = C_{ij} + \sigma_{norm}^2 D_i D_j \text{ lead to d' Agostini bias.}$$

- ▶ Standard solution :

$$\chi^2 = (D - T/r)^T C^{-1} (D - T/r) + \frac{(1 - r)^2}{\sigma_{norm}^2}$$



- ▶ Drawbacks : (1) Additional fitting parameters. (2) Must be taken into account in the Hessian error analysis.

- ▶ Another solution :

$$\chi^2 = (D - rT)^T C_{-1} (D - rT) + \frac{(1 - r)^2}{\sigma_{norm}^2}$$

Advantages : (1) No additional open parameters as the optimal r can be calculated analytically. (2) The Hessian method automatically includes the contribution from r .

TREATMENT OF DEUTERON CORRECTION



- ▶ Without deuteron correction, F_2^D is computed as : $F_2^D = F_2^p + F_2^n \equiv F_2^N$.
- ▶ However, $F_2^D \neq F_2^N$.
- ▶ How we compute F_2^D :

$$F_2^D = F_2^{p,CTEQ} \times \frac{F_2^{D,CJ}}{F_2^{p,CJ}}$$

where $F_2^{D,CJ}$, $F_2^{p,CJ}$ comes from CJ15 analysis [Accardi *et al*, Phys.Rev.D93 11,(2016) 114017].

OPEN PARAMETERS IN NEUTRINO FITS

$$\begin{aligned} & a_1^{u\nu}, a_2^{u\nu}, a_4^{u\nu}, a_5^{u\nu}, b_1^{u\nu}, b_2^{u\nu}, \\ & a_1^{d\nu}, a_2^{d\nu}, a_4^{d\nu}, a_5^{d\nu}, b_1^{d\nu}, b_2^{d\nu}, \\ & a_1^{\bar{u}+\bar{d}}, a_2^{\bar{u}+\bar{d}}, a_5^{\bar{u}+\bar{d}}, \\ & a_1^g, a_4^g, a_5^g, b_0^g, b_1^g, b_4^g, b_5^g, \\ & a_0^{s+\bar{s}}, a_1^{s+\bar{s}}, a_2^{s+\bar{s}}, b_0^{s+\bar{s}}, b_2^{s+\bar{s}} \end{aligned}$$

COMBINED FIT?

- ▶ **Important question** : **issue with the data** or **incomplete understanding of neutrino interactions in nuclear environment?**
- ▶ Need new neutrino DIS data.
- ▶ In the mean time, three strategies :
 - ▶ Issue with experimental uncertainties → Ignore correlations, Weight variation, etc.
 - ▶ Issue with possibly different shadowing mechanism → Restricting Kinematic region → BaseDimuNeuX.
 - ▶ Select data sets that pass the compatibility criteria → BaseDimuChorus.