

First-order nucleon-to-quark phase transition: Thermodynamically-consistent constructions other than Maxwell and Gibbs

C. Constantinou

MICRA2023
Microphysics in Computational Relativistic Astrophysics
ECT*, 11 - 15 September 2023

Collaborators: M. Guerrini (UNIFE), S. Han (TDLI), T. Zhao (OU), M. Prakash (OU)




Motivation: Hybrid Stars (?)

- ▶ The size of nucleons (uncertain as it may be) implies that deconfined quark matter can exist in the cores of NSs.
- ▶ However, such a possibility lacks observational and theoretical support:
 - ▶ Measurements of M , R , Λ cannot differentiate normal and hybrid stars.
 - ▶ LQCD and PQCD not applicable to NS conditions.
- ▶ Possible solution: identify an observable with strong dependence on composition.
- ▶ Enter g-modes!

Hybrid Matter: 1st Order Transitions


▶ **Maxwell** (“strong”, “stiff”, ...)

- ▶ Infinite interface tension
- ▶ No phase mixing
- ▶ Local charge neutrality
- ▶ $\epsilon = f(\epsilon_H + \epsilon_{eH}) + (1 - f)(\epsilon_Q + \epsilon_{eQ})$




▶ **Gibbs** (“weak”, “soft”, ...)

- ▶ Zero surface tension
- ▶ Complete phase mixing
- ▶ Global charge neutrality
- ▶ $\epsilon = f \epsilon_H + (1 - f) \epsilon_Q + \epsilon_{eM}$



▶ **Intermediate case**

- ▶ Some phase mixing
- ▶ Charge neutrality is partially local and partially global
- ▶ $\epsilon = f(\epsilon_H + \eta \epsilon_{eH}) + (1 - f)(\epsilon_Q + \eta \epsilon_{eQ}) + (1 - \eta) \epsilon_{eM}$





Hybrid Matter: 1st Order Transitions (cont'd)

► Constraints

- Baryon number conservation
 $1 = f(y_n + y_p) + (1 - f)(y_u + y_d + y_s)/3$
- Lepton number conservation
 $0 = y_e - f\eta y_{eH} - (1 - f)\eta y_{eQ} - (1 - \eta)y_{eM}$
- Local charge neutrality
 $0 = (y_p - y_{eH}) = (2y_u - y_d - y_s)/3 - y_{eQ}$
- Global charge neutrality
 $0 = fy_p + (1 - f)(2y_u - y_d - y_s)/3 - y_{eM}$

► Equilibrium (= minimization of ε wrt f, y_i, η)

- Mechanical, $P_H + \eta P_{eH} = P_Q + \eta P_{eQ}$
- Quark weak, $\mu_d = \mu_s$
- Neutral strong, $\mu_n = \mu_u + 2\mu_d$
- Charged strong, $\mu_p = 2\mu_u + \mu_d - \eta(\mu_{eH} - \mu_{eQ})$
- Beta
 -or- $\mu_d = \mu_u + \eta\mu_{eQ} + (1 - \eta)\mu_{eM}$
 $\mu_p = \mu_n - \eta\mu_{eH} - (1 - \eta)\mu_{eM}$
- η optimization, $\varepsilon_{eM} = f\varepsilon_{eH} + (1 - f)\varepsilon_{eQ}$



Equation of State

- Nucleons: Zhao - Lattimer

$$\epsilon_B = \sum_{h=n,p} \frac{1}{\pi^2} \int_0^{k_{Fh}} k^2 \sqrt{M_B^2 + k^2} dk + n_B V(u, x)$$

$$V = 4x(1-x)(a_0 u + b_0 u^\gamma) + (1-2x)^2(a_1 u + b_1 u^{\gamma_1})$$

- Quarks: vMIT



$$\mathcal{L} = \sum_{q=u,d,s} [\bar{\psi}_q (i\not{\partial} - m_q - B) \psi_q + \mathcal{L}_{\text{int}}] \Theta$$

$$\mathcal{L}_{\text{int}} = -G_V \sum_q \bar{\psi} \gamma_\mu V^\mu \psi + (m_V^2/2) V_\mu V^\mu$$

$$\epsilon_Q = \sum_q \epsilon_{\text{FG},q} + \frac{1}{2} \left(\frac{G_V}{m_V} \right)^2 n_Q^2 + B$$

- Leptons: noninteracting, relativistic fermions

$$\epsilon_L = \sum_{l=e,\mu} \frac{1}{\pi^2} \int_0^{k_{Fl}} k^2 \sqrt{m_l^2 + k^2} dk$$



- ▶ Global, long-lived, nonradial fluid oscillations resulting from fluid-element perturbations in a stratified environment.
- ▶ Slow chemical equilibration generates buoyancy forces to oppose displacement.
- ▶ In stably-stratified systems the opposing force sets up oscillations with a characteristic frequency [Brunt-Väisälä, $N^2 = g^2 \Delta(c^{-2}) e^{\nu-\lambda}$] which depends on both the equilibrium and the adiabatic sound speeds [$\Delta(c^{-2}) = c_{\text{eq}}^{-2} - c_{\text{ad}}^{-2}$].
 - ▶ $c_{\text{eq}}^2(n_B) = \frac{dP}{d\varepsilon} = \frac{dP_\beta}{dn_B} \left(\frac{d\varepsilon_\beta}{dn_B} \right)^{-1}$
mechanical equilibrium restored instantaneously.
 - ▶ $c_{\text{ad}}^2(n_B, x) = \left(\frac{\partial P}{\partial \varepsilon} \right)_x = \left. \frac{\partial P}{\partial n_B} \right|_x \left(\left. \frac{\partial \varepsilon}{\partial n_B} \right|_x \right)^{-1}$
 $c_{\text{ad},\beta}^2(n_B) = c_{\text{ad}}^2[n_B, x_\beta(n_B)]$
slow restoration of chemical equilibrium because $\tau_\beta \gg \tau_{\text{oscillation}}$.
- ▶ g-mode oscillations couple to tidal forces; they can be excited in a NS merger and provide information on the interior composition.
- ▶ Detection remains a challenge; but within sensitivity of 3rd generation detectors.

g-mode Signals

- ▶ $\Delta(c^{-2}) = c_{\text{eq}}^{-2} - c_{\text{ad}}^{-2}$ drives the restoring force for g-mode oscillations. In *npe* matter,

$$c_{\text{ad}}^2 = c_{\text{eq}}^2 + \left[n_B \left(\frac{\partial \tilde{\mu}}{\partial n_B} \right)_x \right]^2 \left[\mu_n \left(\frac{\partial \tilde{\mu}}{\partial x} \right)_{n_B} \right]^{-1}$$



$$\tilde{\mu} = \mu_e + \mu_p - \mu_n \xrightarrow{\beta\text{-eq.}} 0$$

- ▶ g-modes in 1st-order hybrid matter have larger frequency range compared to pure nucleon matter $\hat{=}$ the behavior of $\Delta(c^{-2})$ in the mixed phase.

- ▶ Dramatic changes in ν_g require new DOFs not just a smooth change in composition.

Discontinuity g-modes

- ▶ Generated by the flatness of $P(n_B)$ in a Maxwell mixed phase that leads to a density jump in the core of a hybrid star.
- ▶ Characterized by discontinuous g-mode frequencies.
- ▶ A special case of a compositional g-mode in the limit $\eta \rightarrow 1$.



- ▶ Construction of a thermodynamically-consistent framework for the treatment of 1st-order phase transitions intermediate to Maxwell and Gibbs.
- ▶ Beware of the Maxwell construction!
- ▶ Calculation of g-mode properties for 1st-order phase transitions and for crossovers.
- ▶ g-modes can detect nonnucleonic matter in the cores of NS; assuming quark matter (by some other means), g-modes can distinguish between a first-order phase transition and a crossover.
- ▶ Discontinuity g-modes as a special case of compositional g-modes in the Maxwell limit.
- ▶ (Near) Future:
 - ▶ Extend 1st-order phase transition scheme to finite T .
 - ▶ Applications to protoneutron stars (cooling, superfluidity) and binary mergers.
 - ▶ Construct EOS that uses the same underlying description for quarks and hadrons.