



From COMPASS to AMBER

Trento, Italia, 18-22/09/2023



Spin crisis? It is over.. Mass “crisis”? Knocking in the door...
(how much we have learned so far about proton spin (selected topics), what is next science question to be addressed?)

Parton distribution functions at a crossroad

Outlook

1. Intro: Spin and Transverse Momentum Dependent PDFs
2. Polarised SIDIS:
 - Sivers function story
3. Crucial TMDs approach test:
 - SIDIS vs Drell-Yan
 - COMPASS results
4. Intro: EHM and pion – proton mass difference
5. CERN’s road map main focus and contribution by AMBER
7. Summary




Dr. Oleg Denisov, senior researcher INFN section of Turin, Italy

Materials/slides of Vincent Andrieux, Craig Roberts, Bakur Parsamyan, Alessandro Bacchetta, Stefan Wallner, Jan Friedrich, Stephan Paul and others have been used

On the one hand - Almost all visible matter of the universe we are able to observe consists of nucleons.

On the other hand - **SPIN is a fundamental quantum number** (Pauli principle), to some extent define a rules on how the atomic/nuclear matter is constructed.



Thus we better understand well how the spin of the nucleon (and hadron in general) is “constructed”.

$$\text{Nucleon spin } \frac{1}{2} = \frac{1}{2} \Delta\Sigma + \Delta G + L$$


quark gluon orbital mom.

$\Delta\Sigma$: sum over u, d, s, \bar{u} , \bar{d} , \bar{s}

Can take any value: superposition of several states


$$\Delta q = \vec{q} - \overleftarrow{q}$$

Parton spin parallel or anti parallel to nucleon spin




First two component were extensively studied in the SIDIS experiments with the longitudinally polarised target (collinear case approach): spin fraction carried by quarks and gluons is not sufficient to describe $\frac{1}{2}$ nucleon spin (**Spin Crisis, continued**):

- Quark spin contribution $\Delta\Sigma=0.24$ ($Q^2=10$ (GeV/c)² DSSV [arXiv:0804.0422](https://arxiv.org/abs/0804.0422))
- RHIC and COMPASS Open charm measurement and other direct measurements → $\Delta G/G$ is not sufficient →







In order to create Angular Momentum of partons spin-orbit correlation has to be taken into account → transverse momentum of the quark k_T appears → **3D structure of the Nucleon has to be studied**

Unified View of Nucleon Structure



Four probes to access transverse hadron structure (TMD PDFs)




18 structure functions
14 azimuthal modulations

$$\frac{d\sigma}{dx dy dz dp_T^2 d\phi_h d\phi_s} =$$

$$\left[\frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L})$$

$$\times \left\{ \begin{array}{l} 1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h + \varepsilon A_{UU}^{\cos 2\phi_h} \cos 2\phi_h \\ + \lambda \sqrt{2\varepsilon(1-\varepsilon)} A_{LU}^{\sin\phi_h} \sin\phi_h \\ + S_L \left[\sqrt{2\varepsilon(1+\varepsilon)} A_{UL}^{\sin\phi_h} \sin\phi_h + \varepsilon A_{UL}^{\sin 2\phi_h} \sin 2\phi_h \right] \\ + S_L \lambda \left[\sqrt{1-\varepsilon^2} A_{LL} + \sqrt{2\varepsilon(1-\varepsilon)} A_{LL}^{\cos\phi_h} \cos\phi_h \right] \\ + S_T \left[\begin{array}{l} A_{UT}^{\sin(\phi_h-\phi_s)} \sin(\phi_h-\phi_s) \\ + \varepsilon A_{UT}^{\sin(\phi_h+\phi_s)} \sin(\phi_h+\phi_s) \\ + \varepsilon A_{UT}^{\sin(3\phi_h-\phi_s)} \sin(3\phi_h-\phi_s) \\ + \sqrt{2\varepsilon(1+\varepsilon)} A_{UT}^{\sin\phi_s} \sin\phi_s \\ + \sqrt{2\varepsilon(1+\varepsilon)} A_{UT}^{\sin(2\phi_h-\phi_s)} \sin(2\phi_h-\phi_s) \end{array} \right] \\ + S_T \lambda \left[\begin{array}{l} \sqrt{(1-\varepsilon^2)} A_{LT}^{\cos(\phi_h-\phi_s)} \cos(\phi_h-\phi_s) \\ + \sqrt{2\varepsilon(1-\varepsilon)} A_{LT}^{\cos\phi_s} \cos\phi_s \\ + \sqrt{2\varepsilon(1-\varepsilon)} A_{LT}^{\cos(2\phi_h-\phi_s)} \cos(2\phi_h-\phi_s) \end{array} \right] \end{array} \right\}$$



Quark \ Nucleon	U	L	T
U	$f_1^q(x, k_T^2)$ number density		$h_1^{\perp q}(x, k_T^2)$ Boer-Mulders
L		$g_1^q(x, k_T^2)$ helicity	$h_{1L}^{\perp q}(x, k_T^2)$ worm-gear L
T	$f_{1T}^{\perp q}(x, k_T^2)$ Sivers	$g_{1T}^q(x, k_T^2)$ Kotzinian-Mulders worm-gear T	$h_1^q(x, k_T^2)$ transversity $h_{1T}^{\perp q}(x, k_T^2)$ pretzelosity


+ two FFs: $D_{1a}^h(z, P_1^2)$ and $H_{1a}^{\perp h}(z, P_1^2)$

At leading order, three PDFs are needed to describe the nucleon in the collinear case. If one admit a non-zero transverse quark momentum k_T in the nucleon five more PDFs (TMD PDFs) are needed. In this talk dedicated attention to non zero structure function Sivers function $f_{1T}^{\perp q}(x, k_T)$. It describes the influence of the transverse spin of the nucleon onto the quark transverse momentum distribution → provides model-dependent access to the orbital momentum

COMPASS Results of 2005

Hep-ex/0503002

Solid state ⁶LD polarised target




Full points – positive hadrons,
 Open points – negative hadrons

Hermes Results of 2004

hep-ph/0408013

Gaseous H₂ polarized target




DOUBTS.....

$$A_{UT}^{\sin(\phi_h - \phi_s)} \propto f_{1T}^{\perp q} \otimes D_{1q}^h$$

Joint data analysis form Hermes and COMPASS – no contradictions



As it was shown by Mauro Anselmino and Colleagues (second half of 2005) when first extraction of Sivers function has been performed from Hermes and COMPASS data (Transversity'2005, hep-ph/051101)) that the contributions from u- and d-quarks are opposite




Sivers 2009 – final results Hermes&COMPASS data perfectly fits together

COMPASS Final results on deuteron
(data 2002-2004) PLB 673 (2009)

Hermes Final results on proton
PRL 103 (2009)





Flavour separation is essential



Second round: COMPASS \leftrightarrow Hermes proton data

COMPASS final results on proton
(data 2007, 2010) PLB 744 (2015)



Hermes Final results on proton
PRL 103 (2009)



COMPASS \leftrightarrow Hermes proton data

COMPASS Sivvers is smaller – QCD evolution eff.?

Hint from the data: even if exist evolution has to be rather slow



The time-reversal odd character of the Sivers and Boer-Mulders PDFs lead to the prediction of a sign change when accessed from SIDIS or from Drell-Yan processes:

↪ Check the predictions:

$$f_{1T}^{\perp}(DY) = -f_{1T}^{\perp}(SIDIS)$$

$$h_1^{\perp}(DY) = -h_1^{\perp}(SIDIS)$$

Its experimental confirmation is considered a crucial test of non-perturbative QCD.

Universality test includes not only the sing-reversal character of the TMDs but also the comparison of the amplitude as well as the shape of the corresponding TMDs

Andreas Metz (Trento-TMD'2010):

Sign reversal of the Sivers function

- Prediction based on operator definition (Collins, 2002)

$$f_{1T}^\perp|_{DY} = - f_{1T}^\perp|_{DIS}$$


- What if sign reversal of f_{1T}^\perp is **not** confirmed by experiment?

- Would not imply that QCD is wrong
- Would imply that SSAs not understood in QCD
- Problem with TMD-factorization
- Problem with resummation of large logarithms
 - Resummation relevant if more than one scale present
 - CSS resummation in Drell-Yan (Collins, Soper, Sterman, 1985); resum logarithms of the type

$$\alpha_s^k \ln^{2k} \frac{\vec{Q}_T^2}{Q^2}$$


- Has also implications for Fermilab and LHC physics

2005 – Anatoly Efremov brings my attention for the first time to this effect (discussed in the famous paper by John Collins *Phys.Lett.B* 536 (2002) 43-48)



SIBIS

Drell-Van



$$s = (P_a + P_b)^2,$$

$$x_{a(b)} = q^2 / (2P_{a(b)} \cdot q),$$

$$x_F = x_a - x_b,$$

$$M_{\mu\mu}^2 = Q^2 = q^2 = s x_a x_b,$$

$$k_{T a(b)}$$

$$q_T = P_T = k_{T a} + k_{T b}$$


the momentum of the beam (target) hadron,
the total centre-of-mass energy squared,
the momentum fraction carried by a parton from $H_{a(b)}$,
the Feynman variable,
the invariant mass squared of the dimuon,
the transverse component of the quark momentum,
the transverse component of the momentum of the virtual photon.

SIDIS data:


- Global fits of available 1-D SIDIS data
- Different TMD evolution schemes
- Different predictions for Drell-Yan



- Extremely important to extract Sivers in SIDIS in Drell-Yan Q^2 range

M.G. Echevarria, A.Idilbi, Z.B. Kang and I. Vitev,
PRD 89 074013 (2014)




P. Sun and F. Yuan, **PRD 88 11, 114012 (2013)**







COMPASS PLB 770 (2017) 138





High mass Drell-Yan region: Kinematic coverage



- Valence region ($u\bar{u}$ annihilation).
- $\langle M_{\mu\mu} \rangle = 5.3 \text{ GeV}/c^2.$
- $q_T > 0.4 \text{ GeV}/c$ required.
- $\langle q_T \rangle = 1.17 \text{ GeV}/c.$



$$A_T^{\sin \varphi_S} \propto f_{1,\pi}^q \otimes f_{1T,p}^{\perp q} \quad (\text{number density} \otimes \text{Siverson function})$$




SIDIS in the corresponding Q^2 range.

$$A_{UT}^{\sin(\varphi_h - \varphi_S)} = f_{1T,p}^{\perp q} \otimes D_{1,q}^h$$


(Siverson \otimes unpolarised FF)

[Phys.Lett.B770 (2017) 138]



DGLAP (2016)
M. Anselmino et al., [arXiv:1612.06413](https://arxiv.org/abs/1612.06413)



TMD-1 (2014)
M. G. Echevarria et al. [PRD89,074013](https://arxiv.org/abs/1407.0740)



TMD-2 (2013)
P. Sun, F. Yuan, [PRD88, 114012](https://arxiv.org/abs/1303.1140)



**In 2018 – 2nd round of
polarized DY measurements
at COMPASS**



Integrated, compared to predictions.
Curves: [Bastami et al., JHEP 02 (2021) 166]

$$\frac{d\sigma}{dx dy dz dp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots \right.$$

$$+ S_T \left[\begin{array}{l} A_{UT}^{\sin(\phi_h - \phi_S)} \sin(\phi_h - \phi_S) \\ + \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) \\ + \varepsilon A_{UT}^{\sin(3\phi_h - \phi_S)} \sin(3\phi_h - \phi_S) \\ + \sqrt{2\varepsilon(1+\varepsilon)} A_{UT}^{\sin\phi_S} \sin\phi_S \\ + \sqrt{2\varepsilon(1+\varepsilon)} A_{UT}^{\sin(2\phi_h - \phi_S)} \sin(2\phi_h - \phi_S) \end{array} \right]$$

$$\frac{d\sigma^{LO}}{d\Omega} \propto F_U^1 (1 + \cos^2 \theta_{CS}) \left\{ 1 + \dots \right.$$

$$+ S_T \left[\begin{array}{l} A_T^{\sin\varphi_S} \sin\varphi_S \\ + D_{[\sin^2\theta_{CS}]} \left(\begin{array}{l} A_T^{\sin(2\varphi_{CS} - \varphi_S)} \sin(2\varphi_{CS} - \varphi_S) \\ + A_T^{\sin(2\varphi_{CS} + \varphi_S)} \sin(2\varphi_{CS} + \varphi_S) \end{array} \right) \\ + D_{[\sin 2\theta_{CS}]} \left(\begin{array}{l} A_T^{\sin(\varphi_{CS} - \varphi_S)} \sin(\varphi_{CS} - \varphi_S) \\ + A_T^{\sin(\varphi_{CS} + \varphi_S)} \sin(\varphi_{CS} + \varphi_S) \end{array} \right) \end{array} \right]$$



- There is a very clear recipe to fill up the missing part of the proton spin – angular momentum → 3D case → TMDs and GPDs
- TMDs study will provide essential input for 3-D structure of the hadron
- Experimental prove of the TMDs mechanism **validity is still missing**
- We found ourselves in Precision phase (Alessandro Bacchetta)
- More data to come in the next years from JLAB, COMPASS and later from EIC



GPS compass

Precision



• Exploration	2002
• parton-model theory	
• first measurements	
• Consolidation	2012
• TMD factorization	
• many consistent measurements	
• Precision	2022
• full-fledged global analysis	
• precision measurements	

from IWSS 2011



AMBER

more than 15 years-long effort



We have started to work on physics program of possible COMPASS successor > 15 years ago.

A Number of Workshops has been organized, for detail see AMBER web page:

<https://amber.web.cern.ch/>


EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH



CERN-SPSC-2019-003
SPSC-I-250
January 25, 2019

[LoI submitted in January 2019](http://arxiv.org/abs/1808.00848)
<http://arxiv.org/abs/1808.00848>

Apparatus for Meson and Baryon Experimental Research
> 270 authors



Welcome

Over the past four decades, measurements at the external beam lines of the CERN Super Proton Synchrotron (SPS) have received worldwide attention. The experimental results have been challenging Quantum Chromodynamics (QCD) as our theory of the strong interactions, thus serving as important input to develop improvements of the theory. As of today, these beam lines remain mostly unique and bear great potential for significant future advancements in our understanding of hadronic matter.

In the context of the Physics-beyond-colliders (PBC) initiative at CERN, the COMPASS++/AMBER (proto-) collaboration proposes to establish a "New QCD facility at the M2 beam line of the CERN SPS". Such an unrivalled installation would make the experimental hall EHN2 the site for a great variety of measurements to address fundamental issues of QCD. The proposed measurements cover a wide range in the squared four-momentum transfer Q^2 : from lowest values of Q^2 where we plan to measure the proton charge radius by elastic muon-proton scattering, over intermediate Q^2 where we plan to study the spectroscopy of mesons and baryons by using dedicated meson beams, to high Q^2 where we plan to study the structure of mesons and baryons via the Drell-Yan process and eventually address the fundamental quest on the emergence of hadronic mass [arxiv:1606.03909\[nucl-th\]](https://arxiv.org/abs/1606.03909), [arXiv:1905.05208\[nucl-th\]](https://arxiv.org/abs/1905.05208).

Letter of Intent:

A New QCD facility at the M2 beam line of the CERN SPS*

COMPASS++[†]/AMBER[‡]

B. Adams^{13,12}, C.A. Aidala¹, R. Akhunzyanov¹⁴, G.D. Alexeev¹⁴, M.G. Alexeev⁴¹, A. Amoroso^{41,42},



AMBER PHASE-1 (proposal submitted in Sep. 2019, approved in Dec. 2020)



Program	Physics Goals	Beam Energy [GeV]	Beam Intensity [s^{-1}]	Trigger Rate [kHz]	Beam Type	Target	Earliest start time, duration	Hardware additions
muon-proton elastic scattering	Precision proton-radius measurement	100	$4 \cdot 10^6$	100	μ^\pm	high-pressure H2	2022 1 year	active TPC, SciFi trigger, silicon veto,
Hard exclusive reactions	GPD E	160	$2 \cdot 10^7$	10	μ^\pm	NH_3^\dagger	2022 2 years	recoil silicon, modified polarised target magnet
Input for Dark Matter Search	\bar{p} production cross section	20-280	$5 \cdot 10^5$	25	p	LH2, LHe	2022 1 month	liquid helium target
\bar{p} -induced spectroscopy	Heavy quark exotics	12, 20	$5 \cdot 10^7$	25	\bar{p}	LH2	2022 2 years	target spectrometer: tracking, calorimetry
Drell-Yan	Pion PDFs	190	$7 \cdot 10^7$	25	π^\pm	C/W	2022 1-2 years	
Drell-Yan (RF)	Kaon PDFs & Nucleon TMDs	~ 100	10^8	25-50	K^\pm, \bar{p}	NH_3^\dagger , C/W	2026 2-3 years	"active absorber", vertex detector
Primakoff (RF)	Kaon polarisability & pion life time	~ 100	$5 \cdot 10^6$	> 10	K^-	Ni	non-exclusive 2026 1 year	
Prompt Photons (RF)	Meson gluon PDFs	≥ 100	$5 \cdot 10^6$	10-100	K^\pm π^\pm	LH2, Ni	non-exclusive 2026 1-2 years	hodoscope
K -induced Spectroscopy (RF)	High-precision strange-meson spectrum	50-100	$5 \cdot 10^6$	25	K^-	LH2	2026 1 year	recoil TOF, forward PID
Vector mesons (RF)	Spin Density Matrix Elements	50-100	$5 \cdot 10^6$	10-100	K^\pm, π^\pm	from H to Pb	2026 1 year	

PHASE-1

Conventional **hadron** and **muon** beams

2022 → 2029

PHASE-2

Improved conventional **Hadron/Hadron** and **muon** beam

2029 and beyond

Table 2: Requirements for future programmes at the M2 beam line after 2021. **Muon beams** are in blue, **conventional hadron beams** in green, and **RF-separated hadron beams** in red.

There are two bearing columns of the facility:

1. **Phenomenon of the Emergence of the Hadron Mass**
2. Proton spin? (largely addressed by COMPASS and others, Phase-2)

How does all the visible matter in the universe come about and what defines its mass scale?

Great discovery of the Higgs-boson unfortunately does not help to answer this question, because:

- ✓ The Higgs-boson mechanism produces only a small fraction of all visible mass
- ✓ The Higgs-generated mass scales explain neither the “huge” proton mass nor the ‘nearly-masslessness’ of the pion

As Higgs mechanism produces a few percent of visible mass, Where does the rest comes from (EHM phenomenon)?

Pion




- $M_{\pi} \sim 140\text{MeV}$
- Spin 0
- 2 light valence quarks

Kaon



- $M_K \sim 490\text{MeV}$
- Spin 0
- 1 light and 1 “heavy” valence quarks

Proton



- $M_p \sim 940\text{MeV}$
- Spin 1/2
- 3 light valence quarks

Higgs generated masses of the valence quarks:
 $M_{(u+d)} \sim 7 \text{ MeV}$ $M_{(u+s)} \sim 100 \text{ MeV}$ $M_{(u+u+d)} \sim 10 \text{ MeV}$




EHM phenomenon

What are the underlying mechanisms?

Intuitively one can expect that the answer to the question lies within SM, in particular within QCD.
Why? Because of the dynamical mass generation in continuum QCD.


As quark can emit and absorb gluons
It acquires its mass in infrared region
because of the gluon “self-mass-generation” mechanism, so the visible (or emergent) mass of hadrons must be dominated by gluon component



Truly “mass from nothing” phenomenon: Initially massless gluon produces dressed gluon fields which “generates” mass function that is large at infrared momenta



Dynamical mass generation in continuum quantum chromodynamics
J.M. Cornwall, Phys. Rev. D 26 (1981) 1453
 ... ~ 1000 citations



In order to “proof” that QCD underlies the EHM phenomenon we have to compare Lattice and Continuum QCD calculations with experimental data by measuring:

1. Quark and Gluon PDFs of the pion/kaon/proton
2. Hadron’s radii (confinement)
3. Excited-meson spectra

(Constituent quarks)

Dressed-quark mass function $M(p)$

The answer is obviously NOT (SM paradigm):


- proton is described by QCD ... 3 valence quarks
- pion is also described by QCD ... 1 valence quark and 1 valence antiquark
- expect $m_p \approx 1.5 \times m_\pi$... but, instead $m_p \approx 7 \times m_\pi$

Proton and pion/kaon difference:

- In the chiral limit the mass of the proton remains basically the same
- Chiral limit mass of pion and kaon is "0" by definition (Nambu-Goldstone bosons)
- Different gluon content expected for pion and kaon
- Contribution from interplay with Higgs mechanism is different

Thus it is equally important to study the internal structure and dynamics of pions, kaons and protons

Mass Budgets




Questions to be answered:

- Mass difference pion/proton/kaon
- Mass generation mechanism (emergent mass .vs. Higgs)
- Internal quark-gluon structure and dynamics, especially important pion/kaon/proton striking differences


A series of workshops entitled
 “Perceiving of the EHM through
 AMBER@CERN(SPS)”:
<https://indico.cern.ch/event/1021402/>

Methods:


Drell-Yan (compl. to Sullivan) and J/ψ





Prompt Photon Production



Diffractive scattering





Elastic scattering




Conventional muon/hadron M2 beams

Improved conventional hadron M2 beams



Phase-1 

Phase-2 

Proton Radius Measurement
Antimatter production cross section
Pion structure (PDFs) via DY and charmonia
Kaon and pion structure (PDFs and PDAs)


High precision strange-meson spectrum
Kaon and pion charge radius
Kaon induced Primakoff reaction

Phase-1 Proposal approved by RB on 02/12/2020

Phase-2 Proposal submission in the beginning of 2024


Pion induced Drell-Yan at AMBER

Status of the knowledge of the Pion structure



Chang et al, PRD 102, 054024 (2020)

From: E615, PRD 1989



Pion structure status:

- Scarce data, poor knowledge of valence, sea and glue basically unknown
- Mostly heavy nuclear targets: large nuclear effects
- For some experiments, no information on absolute cross sections
- Two experiments (E615, NA3) have measured so far with both pion beam sign, but only one (NA3) has used its data to separate sea-valence quark contributions
- Discrepancy between different experiments (i.e. NA10, E615)
- Old data, no way to reanalyse them using modern approaches




Probing valence and sea quark contents of pion at AMBER

Expected statistics 8 to 20 times higher than available





Apparatus for Meson and Baryon Experimental Research



Pion structure in pion induced DY
Expected accuracy as compared to NA3

- $\Sigma_V = \sigma^{\pi^-C} - \sigma^{\pi^+C}$: only valence-valence
- $\Sigma_S = 4\sigma^{\pi^+C} - \sigma^{\pi^-C}$: no valence-valence
- Collect at least a **factor 10 more statistics** than presently available
- Minimize nuclear effects on target side
 - Projection for 2×140 days of Drell-Yan d 3:1 king
 - π^+ to π^- 10:1 time sharing
 - 190 GeV beams on Carbon target ($1.9\lambda_{int}^\pi$)
 - Improvement of shielding to double the intensity is under investigation



Sea quark content of pion can be accurately measured at AMBER for the first time

Experiment	Target type	Beam energy (GeV)	Beam type	Beam intensity (part/sec)	DY mass (GeV/c ²)	DY events
E615	20 cm W	252	π^+	17.6×10^7	4.05 – 8.55	5000
			π^-	18.6×10^7		30000
NA3	30 cm H ₂	200	π^+	2.0×10^7	4.1 – 8.5	40
			π^-	3.0×10^7		121
	6 cm Pt	200	π^+	2.0×10^7	4.2 – 8.5	1767
			π^-	3.0×10^7		4961
NA10	120 cm D ₂	286	π^-	65×10^7	4.2 – 8.5	7800
		140			4.35 – 8.5	3200
	12 cm W	286	π^-	65×10^7	4.2 – 8.5	49600
		194			4.07 – 8.5	155000
		140			4.35 – 8.5	29300
COMPASS 2015	110 cm NH ₃	190	π^-	7.0×10^7	4.3 – 8.5	35000
COMPASS 2018						52000
	75 cm C	190	π^+	1.7×10^7	4.3 – 8.5	21700
						4.0 – 8.5
AMBER	12 cm W	190	π^-	6.8×10^7	4.3 – 8.5	67000
						4.0 – 8.5
	12 cm W	190	π^+	0.4×10^7	4.3 – 8.5	8300
						4.0 – 8.5
	12 cm W	190	π^-	1.6×10^7	4.3 – 8.5	24100
						4.0 – 8.5

AMBER


Isoscalar target + Both positive and negative beams + High statistics



Pion induced J/ψ at AMBER



Apparatus for Meson and Baryon
Experimental Research




Collected simultaneously with DY data, with large counting rates

Physics objectives:

- Study of the J/ψ (charmonia) production mechanisms (gg -fusion vs $q\bar{q}$ -annihilation), comparison of **CEM** and **NRQCD**
- Probe gluon and quark PDFs of pion (arXiv:2103.11660v1 [hep-ph] 22 Mar 2021)
- $\Psi(2S)$ signal study, free of feed-down effect from χ_{c1} χ_{c2}


Cheung and Vogt, priv. comm.



Improved CEM, CT10 + GRS99 global fit for proton/pion


Experiment	Target type	Beam energy (GeV)	Beam type	J/ψ events
NA3 [76]	Pt	150	π^-	601000
		280	π^-	511000
		200	π^+ π^-	131000 105000
E789 [129, 130]	Cu	800	p	200000
	Au			110000
	Be			45000
E866 [131]	Be	800	p	3000000
	Fe			124700
NA50 [132]	Al	450	p	100700
	Cu			130600
	Ag			132100
	W			78100
NA51 [133]	P d	450	p	301000 312000
HERA-B [134]	C	920	p	152000
COMPASS 2015 COMPASS 2018	110 cm NH_3	190	π^-	1000000 1500000
AMBER	75 cm C	190	π^+	1200000
			π^-	1800000
		p	1500000	
	12 cm W	190	π^+	500000
			π^-	700000
	p	700000		

Cross section (ICEM)



Polarization (ICEM)

CHEUNG AND VOGT,
PRIV. COMM., 2020



Both x_F -distribution and polarization depend on the relative amount of valence and glue




Huge statistics: π^+ , π^- , p : 1.2 – 1.8 M J/ψ and 20 – 30 k ψ'

Extremely important to compare the gluon content of kaon and pion (emergent mass)

- Identify the kaon component with the CEDARs
 - positive beam (K = 1.5%)
 - negative beam (K = 2.4%)
- Expected statistics
 - 210 days of positive beam (K+)
 - 70 days of negative beam (K-)
 - CEDARs efficiency: 60%


Nb of events: 25 000 K⁻

32 000 K⁺



Projected statistical errors after 280 days of running, compared to NA3 stat. errors


Proton Radius Measurement at AMBER (confinement)



Spectroscopy

ep	μp
<p>New measurements with</p> <ul style="list-style-type: none"> lower systematics new transitions 	
<p>New measurements with</p> <ul style="list-style-type: none"> lower systematics reaching lower Q^2 <p>ProRAD, ULQ2, ISR @ MESA, PRad</p>	<p>No data yet.</p> <p>MUSE at PSI coming soon</p> <p>AMBER</p>

Scattering



statistical precision of the proposed measurement, down to $Q^2 = 0,001 \text{ GeV}^2/c^2$, Cross section is normalised to the G_D - dipole form factor

$$\langle r_p^2 \rangle = -6\hbar^2 \cdot \left. \frac{dG_E(Q^2)}{dQ^2} \right|_{Q^2 \rightarrow 0}$$

$$\frac{d\sigma^{\mu p \rightarrow \mu p}}{dQ^2} = \frac{4\pi\alpha^2}{Q^4} R (\epsilon G_E^2 + \tau G_M^2) \quad \epsilon = \frac{E_\mu^2 - \tau(s - m_\mu^2)}{\vec{p}_\mu^2 - \tau(s - 2m_p^2(1 + \tau))} \quad \tau = \frac{Q^2}{(4m_p^2)}$$

- Suppress magnetic form factor G_M^2
 - Requires $\tau \rightarrow 0$
 - Measurement at low- Q^2 values of $\mathcal{O}(<10^{-2})$
- Measurement at high-energy $\mathcal{O}(10 - 100 \text{ GeV})$
 - Results in $\epsilon \rightarrow 1$
 - Cross-section directly proportional to G_E^2



Proton Radius Experiment at Jefferson Lab

PRoton
Radius



MESA
Mainz
Energy-recovering
Superconducting
Accelerator




Proton Radius Measurement at AMBER (confinement)





- A number of experiments is on the way in different laboratories
- There is a synergy between PRES at MAMI ($E_e = 720 \text{ MeV}$) and AMBER ($E_\mu = 100 \text{ GeV}$):
 - The same type of active target (hydrogen filled TPC) will be used for both experiment
 - The same Q^2 range will be covered ($10^{-3} - 4 \times 10^{-2} \text{ GeV}^2$)
 - Mutual calibration of the transferred momentum
- Significant advantage of the AMBER measurement is much lower radiative corrections: for soft bremsstrahlung photon energy $E_\gamma/E_{\text{beam}} \sim 0.01$ QED corrections amount to $\sim 15\text{-}20\%$ for electrons and to $\sim 1.5\%$ for muons (AMBER will be able to make a control measurement with Electromagnetic Calorimeters).

If compared to the muon scattering experiment at PSI (MUSE):

- Much cleaner experimental conditions (pure muon beam with less than 10^{-6} admixture of hadrons)
- Much higher beam momentum, thus contribution from magnetic form factor is suppressed ($0.1\text{-}0.2 \text{ GeV}/c$ vs $100 \text{ GeV}/c$)
- Small statistical errors achievable with the proposed running time



Precise measurements of pion and kaon radii will reveal the compositeness (confinement) scale for (near) Nambu-Goldstone bosons. At the moment there is basically no precise experimental information on kaon charge radius.



$$s = 2E_b m_e + m_b^2 + m_e^2$$

$$Q_{max}^2 = \frac{4p_b^2 m_e^2}{s}$$

Beam	E_b [GeV]	Q_{max}^2 [GeV ²]	$E'_{b,min}$ [GeV]	Relative charge-radius effect on c.s. at Q_{max}^2
π	190	0.176	17.3	~40%
K	190	0.086	105.7	~20%
	80	0.066	59.9	~15%
	50	0.037	41.3	~8%

For **kaons**, a significant increase of the form factor knowledge in the range $0.001 < Q^2 < 0.07$ appears in reach with AMBER using an **80 GeV rf-separated kaon beam**

S. R. Amendolia, et al., Phys. Lett. B 178, 435 (1986)





Fig. 3. The measured kaon form factor squared. The line corresponds to the pole fit with $\langle r^2 \rangle = 0.34 \text{ fm}^2$.

PDG lists 25 strange mesons



- ▶ 16 established states, 9 need further confirmation
- ▶ Missing states with respect to quark-model predictions
- ▶ Many measurements performed more than 30 years ago



Stefan Wallner's talk at HADRON'23

Strange-Meson Spectroscopy with COMPASS

The $K^- \pi^- \pi^+$ Data Sample




- ▶ World's largest data set of about 720 k events
- ▶ Rich spectrum of overlapping and interfering X^-
 - ▶ Dominant well known states
 - ▶ States with lower intensity are "hidden"

AMBER QCD Facility, goal for Kaon induced Spectroscopy to Collect $10\text{-}20 \times 10^6$ $K^- \pi^+ \pi^-$ events using high-intensity high-energy kaon beam:


- Optimised Conventional Hadron beam line
- Higher wrt COMPASS beam intensity
- Better pion/kaon beam particles separation
- Much more powerful pid in the final state



Primakoff at AMBER: Chiral Anomaly and Polarizabilities (kaon enriched beam)




- Data
- Primakoff MC
- 3 π MC
- Sum of MC contributions




Polarizabilities


Interaction between **hadron** and **external electromagnetic field** described by parameters α , β (LO), encoding information about its internal structure



PRL 114, 062002 (2015)



hep-ex 1808.00848v6




Dominik Ecker's talk at HADRON'23



AMBER (Prompt Photons)








Apparatus for Meson and Baryon
Experimental Research



Prompt photons probe – direct access to the gluon content of the kaon.
At the moment there is no experimental information about gluon contribution in kaon.

Pythia-based MC simulation for prompt photons production was used for preliminary estimation of kinematic range accessible at COMPASS. It was compared with corresponding ranges accessible by previous experiments with pion beams.

Possibilities to identify signal and reject background were tested. Some optimization of the setup from point of the material budget was tested.





AMBER Phase-1 running plan




Milestones:

1. May 1st 2023 – Antimatter production Run (Std. DAQ)
2. Sep. 1st 2023 – PRM pilot (FreeDAQ, very limited setup)
3. May 1st 2024 – PRM Run (FreeDAQ, limited setup)
4. Sep. 1st 2025 – DY Pilot (FreeDAQ, all trackers + mu id)
5. May 1st 2028 – DY Run (Full Spectr. Ex. RICH, Calorimeters)



- Title
- 1) Proton Radius
 - 1.1) 2021 TEST Run
 - 1.2) 2022 TEST Run
 - 1.3) 2023 TEST Run
 - 1.4) 2024 TEST Run
 - 1.5) 2025 Run
- 2) Anti-Matter production
 - 2.1) Test measurement
 - 2.2) Commissioning
 - 2.3) Data Taking 2023
 - 2.4) Change-over to PRM
 - 2.5) Data Taking 2024
- 3) Drell-Yan
 - 3.1) First test Run
 - 3.2) First RUN






Summary: AMBER at CERN SPS



- A wide and extremely competitive physics program brought together, strong interest in the hadron physics community
- 33 Institutions and 13 countries, ~200 members
- Main goal of the AMBER Phase-1: high precision study of the pion structure as well as first study of the kaon structure via Drell-Yan and J/Psi production
- Improved hadron beam for Phase-2 → unique new opportunities in Hadron Physics



Spares



factorization

TMD Fragmentation Functions (FF)

TMD Distribution Functions (DF)

$$\sigma^{\ell p \rightarrow \ell h X} = \sum_q (\text{DF} \otimes \sigma^{\ell q \rightarrow \ell q} \otimes \text{FF})$$

(Un)polarized SIDIS process allows to probes both TMD PDFs and FFs







STAR: W -Boson Production in $p\uparrow+p$: $p+p \rightarrow W_{\pm} \rightarrow e_{\pm} + \nu$



Apparatus for Meson and Baryon
Experimental Research

- Very important STAR (RHIC) result:
- First experimental investigation of Sivers-non-universality in pp collision (W/Z production)
- Very different hard scale (Q^2) compared to the available SIDIS (FT) data
- QCD evolution effects may play a substantial role


Phys. Rev. Lett. 116, 132301 (2016)
Comparison with Phys. Rev. Lett. 103, 172001



Bacchetta et al., Phys. Lett. B : Lett. B 827 (2022) 136961
Comparison with PRL116(2016) 13201




$$A_T^{\sin(2\varphi_{CS} + \varphi_S)} \propto h_{1,\pi}^{\perp q} \otimes h_{1T,p}^{\perp q} \quad (\text{Boer-Mulders} \otimes \text{pretzelosity})$$



Compatible with zero, no significant kinematic dependence visible.


The error bars are statistical, the color bands show systematic uncertainty.

An additional scale uncertainty of 5% is not shown (dilution factor, λ , polarization).




Integrated, compared to predictions.

Curves: [Bastami et al., JHEP 02 (2021) 166]




SIDIS in the corresponding Q^2 range.


$$A_{UT}^{\sin(3\varphi_h - \varphi_S)} \propto h_{1T,p}^{\perp q} \otimes H_{1,q}^{\perp h} \quad (\text{pretzelosity} \otimes \text{Collins FF})$$



Model dependence of the J/ψ production cross section




Relative contribution
From quarks and gluons
Very uncertain




SMRS vs JAM fits: strong dependence on the PDFs



$$A_T^{\sin(2\varphi_{CS}-\varphi_S)} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^q \quad (\text{Boer-Mulders function} \otimes \text{transversity})$$




Negative (about 1.5σ significance), kinematic dependence not really significant.
The error bars are statistical, the color bands show systematic uncertainty.
An additional scale uncertainty of 5% is not shown (dilution factor, λ , polarization).



Integrated, compared to predictions.

Curves: [Bastami et al., JHEP 02 (2021) 166]





SIDIS in the corresponding Q^2 range.

$$A_{UT}^{\sin(\varphi_h + \varphi_S - \pi)} \propto h_{1,p}^q \otimes H_{1,q}^{\perp h} \quad (\text{transversity} \otimes \text{Collins FF})$$

Drell-Yan process is a low cross-section process:

- High intensity hadron beam
- Hadron absorber to protect Spectrometer from a very high secondary flux
- Vertex Detector to compensate losses in resolution because of the absorber in order to improve mass and space resolution



Drell-Yan experiment preparation II

Proposal by LANL group to reuse PHENIX Silicon Vertex Detector






Figure 7 (a) A completed half FVTX detector, with sensors, frontend electronics, supporting structures, and cooling system. Two half FVTX endcaps are shown on either end. The overall length is about 80 cm. (b) A structural illustration of one endcap of the FVTX. One small disk and three large disks are included in one endcap. (c) A segment (wedge) of the FVTX sensor. Each wedge holds two columns of the silicon strips as shown in the zoomed-in portion.

Table 1 Summary of the FVTX specifications.

Silicon sensor thickness (μm)	320
Strip pitch (μm)	75
Number of strips per column	1664
Inner radius of silicon (mm)	44
Outer radius of silicon (mm)	168.8
Strip length at inner radius (mm)	3.4
Strip length at outer radius (mm)	11.5
Pulse timing (ns)	30
Number of wedges per disk	48



Active silicons mini-strip sensors plus front-end ASIC,
the FPHX chip bonded directly on sensors

- Time resolution: \sim ns
- Spatial resolution: $\sim 20\mu\text{m}$

Simulations and optimisation of the
apparatus and reconstruction ongoing

Preliminary:

$$\rightarrow \sigma_{\mu\mu} \sim 110 \text{ MeV}/c^2$$

$$M_{\mu\mu} > 4.3 \text{ GeV}/c^2 \rightarrow M_{\mu\mu} > 4.0 \text{ GeV}/c^2:$$


$\Rightarrow \sim 50\%$ gain in DY statistics

Study and optimisation of the shielding to:


- Contain the radiation
- Minimise the environmental impact
- Comply with regulations

⇒ **Compatible with 2×current Intensities**

⇒ **ECR to be submitted**




Area	Annual dose limit (year)	Ambient dose equivalent rate		Sign
		permanent occupancy	low occupancy	
Non-designated	1 mSv	0.5 μSv/h	2.5 μSv/h	
Supervised	6 mSv	3 μSv/h	15 μSv/h	
Simple Controlled	20 mSv	10 μSv/h	50 μSv/h	
Limited Stay	20 mSv	-	2 mSv/h	
High Radiation	20 mSv	-	100 mSv/h	
Prohibited	20 mSv	-	> 100 mSv/h	





AMBER Phase-1 Torino construction plan



← Prototipo mec. + consulting

Unified Tracking Station

