

Contribution ID: 35

Type: **not specified**

## **E57 Kaonic Deuterium at J-PARC**

*Wednesday 11 October 2023 11:30 (30 minutes)*

Exotic hadronic atoms are a valuable tool to experimentally investigate the strong interaction described by Quantum Chromodynamics (QCD). With kaonic hydrogen/deuterium atoms one can directly measure the interaction at threshold and will be able to extract the antikaon-nucleon scattering lengths at zero energy (no interpolation to zero energy is necessary as in scattering experiments). The antikaon-nucleon interaction close to threshold provides crucial information on the interplay between spontaneous and explicit chiral symmetry breaking in low-energy QCD. Kaonic hydrogen was measured successfully with SIDDHARTA and DEAR at DAFNE (LNF, Italy) and with KpX at KEK (Japan). A measurement of kaonic deuterium is still missing and just now ongoing at DAFNE. A second measurement with different systematic corrections is prepared at J-PARC. The kaonic deuterium data will allow for the first time, together with the already existing kaonic hydrogen data, the determination of the isospin dependent antikaon-nucleon scattering lengths  $a_0$  and  $a_1$ , eagerly awaited by theory.

An overview of the status of the proposed kaonic deuterium measurement at J-PARC will be given.

**Primary author:** ZMESKAL, Johann (Stefan Meyer Institute)

**Presenter:** ZMESKAL, Johann (Stefan Meyer Institute)

**Session Classification:** Session V