

# Polarization of jet partons in an anisotropic plasma

**Siggi Hauksson**

IPhT, Saclay

ECT\*, Trento


February 14th 2024

In collaboration with Edmond Iancu

JHEP 08 (2023) 027, arXiv:2303.03914

# Early stages of heavy-ion collisions

- How does far-from-equilibrium QCD medium become a fluid?
- Different stages of collisions:  
Glasma  $\rightarrow$  Kinetic theory  $\rightarrow$  Hydrodynamics  
[See e.g. Berges, Heller, Mazeliauskas, Venugopalan (2020)]
- Medium can quench jets and change jet substructure.
- How important are the initial stages for jets?
- Are there some specific signatures on jets from the initial stages?




[Chun Shen, 2014]

# Jet physics in homogeneous, equilibrated medium


- Transverse momentum broadening:

$$\hat{q} = \frac{d\langle \mathbf{p}_{\perp}^2 \rangle}{dt}$$

- Allows for medium-induced gluon emission.
- Wavepackets overlap for a long time during emission.




- Schematic estimate:  $\tau \sim \sqrt{E/\hat{q}}$
- Get rate  $\Gamma \sim \alpha_s P(z)/\tau \sim \alpha_s P(z) \frac{\sqrt{\hat{q}}}{\sqrt{E}}$



- This process determines whole jet structure.

[For vacuum-like emission see e.g. Majumder (2018); Wang, Guo (2001)]



# What changes out of equilibrium?

- Before hydro  $\hat{q}$  is big.
- Big pressure anisotropy at early times.




- Expect to lead to anisotropy in momentum broadening:

$$\hat{q}_z \neq \hat{q}_y \text{ with } \hat{q}_y = \frac{d\langle p_y^2 \rangle}{dt}, \hat{q}_z = \frac{d\langle p_z^2 \rangle}{dt}$$


[Hauksson, Jeon, Gale (2022)]

- Has been seen in glasma stage and kinetic theory stage calculations.

[See also: Ipp, Muller, Schuh (2020); Carrington, Czajka, Mrowczynski (2022)]




[Avramescu, Bäran, Greco, Ipp, Müller, Ruggieri]



[Boguslavski, Kurkela, Lappi, Lindenbauer, Peuron]

# This work


- How does anisotropy in broadening affect jet evolution?




- Jet partons get spin polarization.

- Calculate  $\frac{dP_{a \rightarrow bc}}{d\zeta dt}$  where  $abc \in \{y, z\}$ ,  $\zeta = E_b/E_a$
- Look at how polarization propagates in jet evolution.

- How could this be measured?




# Single gluon emission in an anisotropic medium



- Hard splitting sensitive to different components of  $\mathbf{P}$ . E.g.
  - $|\Gamma_{y \rightarrow z, z}|^2 = 4P_y \bar{P}_y$
  - $|\Gamma_{y \rightarrow z, y}|^2 = 4P_z \bar{P}_z / \zeta^2$
- An anisotropic medium enhances certain polarization states.
  - E.g. if  $\hat{q}_z > \hat{q}_y$ , second case more likely.

$$\frac{d\mathcal{P}_{a \rightarrow bc}}{d\zeta dt} \sim \bar{\alpha}_s \frac{(\hat{q}_y \hat{q}_z)^{1/4}}{\sqrt{2E}} A_{a \rightarrow bc}(\zeta, \hat{q}_z / \hat{q}_y)$$

# Single gluon emission in an anisotropic medium



$$\frac{d\mathcal{P}_{a \rightarrow bc}}{d\zeta dt} = \frac{g^2 N_c}{8\pi\omega^2 \zeta(1-\zeta)} \text{Re} \int_0^L d\Delta t \int \frac{d^2 \mathbf{P}}{(2\pi)^2} \int \frac{d^2 \bar{\mathbf{P}}}{(2\pi)^2} \times \Gamma_{a \rightarrow bc}(\mathbf{P}, \zeta) \Gamma_{a \rightarrow bc}(\bar{\mathbf{P}}, \zeta) \tilde{S}^{(3)}(\Delta t, \mathbf{P}, \bar{\mathbf{P}})$$

[E.g. Baier, Dokshitzer, Mueller, Peigne, Schiff (1996); Zakharov (1996); Wiedemann, Gyulassy (1999); Blaizot, Dominguez, Iancu, Mehtar-Tani (2013)]

- Use polarized vacuum splitting functions:  $\Gamma_{a \rightarrow bc}$
- $\tilde{S}^{(3)}$  describes the momentum broadening of three partons.
- Use harmonic oscillator approximation:  $\hat{q}_z r_z^2 + \hat{q}_y r_y^2$

# Single gluon emission in an anisotropic medium

- Ensemble of gluons: Probability  $p$  of polarization in beam direction.


$$p' = \frac{p \mathcal{P}_{z \rightarrow z} + (1-p) \mathcal{P}_{y \rightarrow z}}{p (\mathcal{P}_{z \rightarrow z} + \mathcal{P}_{z \rightarrow y}) + (1-p) (\mathcal{P}_{y \rightarrow z} + \mathcal{P}_{y \rightarrow y})}$$

- Daughter parton has ( $\zeta = E_b/E_a$ )

$$p' - \frac{1}{2} = f(\zeta) \left(p - \frac{1}{2}\right) + g(\zeta) G(\hat{q}_z/\hat{q}_y)$$

$$f(\zeta) = \frac{\zeta^2}{(1-\zeta)^2 + \zeta^2 + \zeta^2(1-\zeta)^2}, \quad g(\zeta) = \frac{(1-\zeta)^2}{(1-\zeta)^2 + \zeta^2(1-\zeta)^2 + \zeta^2}$$



- Isotropic medium:  
Polarization reduced at each splitting.
- Anisotropic:  
Unpolarized mother radiates polarized daughter!





# Single gluon emission in an anisotropic medium

- Two intuitive limits ( $\zeta = E_b/E_a$ ):
  - $\zeta \rightarrow 1$  :  
$$p' - \frac{1}{2} = (p - \frac{1}{2}) + (1 - \zeta)^2 G(\hat{q}_z/\hat{q}_y)$$
Hard daughter parton gets polarization of parent parton.
  - $\zeta \rightarrow 0$  :  
$$p' - \frac{1}{2} = \zeta^2 (p - \frac{1}{2}) + G(\hat{q}_z/\hat{q}_y)$$
Soft daughter parton is always polarized.
- Size of effect given by  $G(\hat{q}_z/\hat{q}_y)$ .
  - Maximal value is around  $G \sim 0.3$ .
  - Plot as a function of  $\frac{\Delta\hat{q}}{\hat{q}} = \frac{\hat{q}_z - \hat{q}_y}{\hat{q}_z + \hat{q}_y}$ .
- Expected branching is democratic ( $\zeta \sim \frac{1}{2}$ ).
  - Not clear which wins out in the end.
  - Need evolution of jet as a whole



# Evolution of polarization



- Consider total evolution of jet in glasma brick with constant  $G(\hat{q}_z/\hat{q}_y)$ .

- $\tau = \frac{\alpha_s N_c}{\pi} \sqrt{\frac{\hat{q}}{E}} t$

- $$\frac{dD_{\text{tot}}(\xi, \tau)}{d\tau} = \int_{\xi}^1 dz \mathcal{K}_0(\zeta) \sqrt{\frac{\zeta}{\xi}} D_{\text{tot}}\left(\frac{\xi}{\zeta}, \tau\right) - \int_0^1 d\zeta \mathcal{K}_0(\zeta) \frac{\zeta}{\sqrt{\xi}} D_{\text{tot}}(\xi, \tau)$$

- $$\frac{d\tilde{D}(\xi, \tau)}{d\tau} = \int_{\xi}^1 d\zeta \mathcal{M}_0(\zeta) \sqrt{\frac{\zeta}{\xi}} \tilde{D}\left(\frac{\xi}{\zeta}, \tau\right) - \int_0^1 d\zeta \mathcal{K}_0(\zeta) \frac{\zeta}{\sqrt{\xi}} \tilde{D}(\xi, \tau)$$

- $$+ \int_{\xi}^1 d\zeta \mathcal{L}_0(\zeta) \sqrt{\frac{\zeta}{\xi}} D_{\text{tot}}\left(\frac{\xi}{\zeta}, \tau\right).$$


$$\mathcal{K}_0(\zeta) \approx \frac{1}{\zeta^{3/2}(1-\zeta)^{3/2}}, \quad \mathcal{M}_0(\zeta) \approx \zeta^2 \mathcal{K}_0(\zeta), \quad \mathcal{L}_0(\zeta) \approx G(\hat{q}_z/\hat{q}_y)(1-\zeta)^2 \mathcal{K}_0(\zeta)$$

- $D_{\text{tot}} = \xi \frac{d(N_z + N_y)}{d\xi}$  is energy spectrum,  $\tilde{D} = \xi \frac{d(N_z - N_y)}{d\xi}$  is polarization.

[Equation for  $D_{\text{tot}}$ : Blaizot, Iancu, Mehtar-Tani (2013); Blaizot, Mehtar-Tani (2015); Fister, Iancu (2014); Iancu, Wu (2015); Escobedo, Iancu (2016). See also e.g. Mehtar-Tani, Schlichting (2018),

# Propagation of polarization in isotropic medium



- Have solved new equation numerically (and analytically for small  $\xi$ ).
- In an isotropic medium, polarization eventually goes away.
- Use  $\tilde{D}(\xi, \tau = 0) = 1$ : get  $\tilde{D}(\xi, \tau) \sim \xi^{3/2}$  for small  $\xi$ .



- Since  $\mathcal{M}_0(\zeta) = \zeta^2 \mathcal{K}_0(\zeta)$  get  $\tilde{D} = \xi^2 D_{\text{tot}} \sim \xi^{3/2}$

# Propagation of polarization in anisotropic medium

- Anisotropic medium: Use  $D_{\text{tot}}(\xi, \tau = 0) = \delta(1 - \xi)$   $\tilde{D}(\xi, \tau = 0) = 0$ .





- Get that  $\tilde{D} \sim 1/\sqrt{\xi}$  and thus  $\tilde{D}/D_{\text{tot}}$  is constant at small  $\xi$ .
- Confirm by analytical solution at  $\xi \ll \tau$ :
  - Get

$$\tilde{D}(\xi, \tau) \approx \frac{G(\hat{q}_z/\hat{q}_y)}{3} \frac{\tau e^{-\pi\tau^2}}{\sqrt{\xi}}$$

- Nearly constant fraction of jet partons is polarized across different energies.

# How could this be measured?



- Need to measure spin of  $\Lambda$  baryons in jet!
- Measurements have been done at lower  $p_T$ . See e.g. [STAR, Nature 548, 62 (2017)]
  - Motivation to study vorticity of QGP.
  - Weak decay  $\Lambda \rightarrow p + \pi$  and measure angle of  $p$ .
  - Has been measured at tenth of percent.
- Need to connect our theoretical calculation to polarization of  $\Lambda$ .
  - Evolution of medium.
  - Fragmentation.



[Mohanty, 2020]

# How could this be measured?




- Need spin-dependent fragmentation function for  $\Lambda$ .
  - Has been constrained by LEP data in unpolarized  $e^+e^-$ .  
[de Florian, Stratmann, Vogelsang (1997)]
  - Recently used for correlation of spin of two  $\Lambda$ s in heavy-ion collisions.  
[Li, Chen, Cao, Wei (2023)]
- Need to fold our calculation with evolution of medium.
  - Evolution of  $\Delta\hat{q}$  particularly important.
  - How well does spin survive hydro stage?
- Theoretical calculation has to include quarks.



[Boguslavski, Kurkela, Lappi,  
Lindenbauer, Peuron]


# Conclusions

- Early stages have anisotropic momentum broadening,  $\hat{q}_y \neq \hat{q}_z$ .
- Leads to polarization of partons in jet.
  - Have evaluated for gluons in BDMPS-Z framework.
- Competing effect: Abundant sourcing of polarization vs. washing out.
  - Derived and solved evolution equations for jets.
- Can potentially measure through polarization of  $\Lambda$  in jet.




# Jet broadening in glasma

- Jet partons traverse heavily occupied gluon fields. [Ipp, Muller, Schuh (2020); Carrington, Czajka, Mrowczynski (2022); Avramescu, Băran, Greco, Ipp, Müller, Ruggieri (2023)]
- Deflected by chromomagnetic and chromoelectric forces.
- Claim that as much broadening as during hydro stage.
  - $\Delta p_{\perp}^2|_{\text{glasma}}/\Delta p_{\perp}^2|_{\text{hydro}} \approx 0.9$  [Carrington, Czajka, Mrowczynski (2022)]
- Heavily anisotropic broadening,  $\hat{q}_z \approx 2\hat{q}_y$



[Carrington, Czajka, Mrowczynski (2022)]




[Ipp, Muller, Schuh (2020)]



# Jet broadening in kinetic theory

- Anisotropic broadening recently evaluated in kinetic theory.


Boguslavski, Kurkela, Lappi, Lindenbauer, Peuron (2023)



- Longitudinal expansion squeezes momentum distribution.

- Get  $\hat{q}_z > \hat{q}_y$

- Calculation uncertainties: Cutoff dependence, matching with hydro  $\hat{q}$ , strength of coupling.



# Single gluon emission in an anisotropic medium

- $\tilde{S}^{(3)}$  describes the momentum broadening of the three partons during emission.
- $\tilde{S}^{(3)} \sim \langle GGG \rangle$  with propagator

$$G \sim \int \mathcal{D}\mathbf{r} e^{i\frac{E}{2} \int dt \dot{\mathbf{r}}^2} P e^{ig \int dt A^-}$$

- Medium average gives

$$\langle A^-(x^+, \mathbf{x}) A^-(y^+, \mathbf{y}) \rangle \sim \delta(x^+ - y^+) \gamma(\mathbf{x} - \mathbf{y})$$

- Use harmonic oscillator approximation but in anisotropic medium

$$\gamma(0) - \gamma(\mathbf{r}) \sim \hat{q}_z r_z^2 + \hat{q}_y r_y^2$$

- Get that

$$S^{(3)}(\Delta t, \mathbf{u}, \mathbf{v}) = \int_{\mathbf{r}(t=0)=\mathbf{u}}^{\mathbf{r}(t=\Delta t)=\mathbf{v}} \mathcal{D}\mathbf{r} \exp \left\{ iM \int_0^{\Delta t} dt \left[ \frac{\dot{\mathbf{r}}^2}{2} + \frac{\Omega_z^2 r_z^2 + \Omega_y^2 r_y^2}{2} \right] \right\}$$

# Single gluon emission in an anisotropic medium

Get e.g. for  $\frac{d\mathcal{P}_{z \rightarrow z}}{d\zeta dt} = \sum_c \frac{d\mathcal{P}_{z \rightarrow zc}}{d\zeta dt}$

$$\frac{d\mathcal{P}_{z \rightarrow z}}{d\zeta dt} = \bar{\alpha}_s \frac{(\hat{q}_y \hat{q}_z)^{1/4}}{\sqrt{2\omega}} A(\hat{q}_z/\hat{q}_y) \gamma(\zeta) [\mathcal{F}_{z \rightarrow z}^0(\zeta) + G(\hat{q}_z/\hat{q}_y) \mathcal{F}_{z \rightarrow z}^1(\zeta)] .$$

where

$$A(\hat{q}_z/\hat{q}_y) \equiv \frac{1}{2} [f(a) + f(1/a)] , \quad G(\hat{q}_z/\hat{q}_y) \equiv \frac{f(1/a) - f(a)}{f(a) + f(1/a)}$$

$$f(a) \equiv \int_0^\infty dx \left[ \frac{1}{a^{1/2} x^2} - \frac{1}{\sinh^{1/2} ax \sinh^{3/2} x} \right] .$$

and

$$\gamma(\zeta) = \frac{[1 - \zeta(1 - \zeta)]^{1/2}}{\zeta^{1/2}(1 - \zeta)^{1/2}} ,$$

$$\mathcal{F}_{z \rightarrow z}^0(\zeta) = \frac{1}{2} \left( \frac{1 - \zeta}{\zeta} + \frac{2\zeta}{1 - \zeta} + \zeta(1 - \zeta) \right) , \quad \mathcal{F}_{z \rightarrow z}^1(\zeta) = \frac{1}{2} \left( \frac{1 - \zeta}{\zeta} + \zeta(1 - \zeta) \right)$$

# Single gluon emission in an anisotropic medium

- Total unpolarized rate is




$$\frac{d\mathcal{P}}{d\zeta dt} = \frac{1}{2} \sum_{a,b,c} \frac{d\mathcal{P}_{a \rightarrow bc}}{d\zeta dt}$$

- It is nearly unaffected by anisotropy

$$(\zeta = E_b/E_a; \hat{q} = \hat{q}_z + \hat{q}_y)$$

$$\begin{aligned} \frac{d\mathcal{P}}{d\zeta dt} &= \frac{\alpha_s}{2\pi} P_{g \rightarrow g}(\zeta) \frac{\sqrt{1 - \zeta(1 - \zeta)}}{\sqrt{\zeta(1 - \zeta)} E_a} (4\hat{q}_z \hat{q}_y)^{1/4} \\ &\times \frac{1}{2} \left[ f\left(\sqrt{\frac{\hat{q}_z}{\hat{q}_y}}\right) + f\left(\sqrt{\frac{\hat{q}_y}{\hat{q}_z}}\right) \right] \end{aligned}$$

- Plot  $(d\mathcal{P})_{\text{aniso}} / (d\mathcal{P})_{\text{iso}}$  at fixed  $\hat{q}$  with  $\frac{\hat{q}_z - \hat{q}_y}{\hat{q}_z + \hat{q}_y}$  varying.



# Interpretation

- Green's function is


$$G(\xi, \xi_1, \tau) = \frac{1}{\xi_1} \tilde{D}_0 \left( \frac{\xi}{\xi_1}, \frac{\Delta\tau}{\sqrt{\xi_1}} \right) \approx \left( \frac{\xi}{\xi_1(\xi_1 - \xi)} \right)^{3/2} \Delta\tau e^{-\pi\Delta\tau^2/(\xi_1 - \xi)}$$

- Get  $\xi_1 - \xi \sim \Delta\tau^2$
- In general also have  $\Delta\tau \sim \sqrt{\xi}$
- Thus
  - Democratic branching  $\xi_1 \sim 2\xi$ , quasilocal in energy.
  - Also quasilocal in time  $\Delta\tau \sim \sqrt{\xi} \ll 1$ .
- Polarization is produced abundantly.
- Quickly goes away after a few branchings.
- Thus a good approximation is

$$\tilde{D}(\xi, \tau) \propto G(\hat{q}_z/\hat{q}_y) D_{\text{tot}}(\xi, \tau)$$

# Propagation of polarization in isotropic medium

- Have solved new equation numerically (and analytically for small  $\xi$ ).
- In an isotropic medium, polarization quickly goes away.
- Use  $\tilde{D}(\xi, \tau = 0) = 1$ : get  $\tilde{D}(\xi, \tau) \sim \xi^{3/2}$  for small  $\xi$  after time  $\tau \sim \sqrt{\xi}$ .





- Well known that for  $D_{\text{tot}}(\xi, \tau = 0) = \delta(1 - \xi)$

$$D_{\text{tot}}(\xi, \tau) = \frac{\tau}{\sqrt{\xi}(1 - \xi)^{3/2}} e^{-\pi\tau^2/(1-\xi)} \sim \frac{\tau e^{-\pi\tau^2}}{\sqrt{\xi}}$$

- Physics of democratic branching and turbulence.
- Since  $\mathcal{M}_0(\zeta) = \zeta^2 \mathcal{K}_0(\zeta)$  get  $\tilde{D} = \xi^2 D_{\text{tot}}$

# Propagation of polarization in anisotropic medium

- Anisotropic medium: Use  $D_{\text{tot}}(\xi, \tau = 0) = \delta(1 - \xi)$   $\tilde{D}(\xi, \tau = 0) = 0$ .



- Get that  $\tilde{D} \sim 1/\sqrt{\xi}$  and thus  $\tilde{D}/D_{\text{tot}}$  is constant at small  $\xi$ .
- Confirm by analytical solution at  $\xi \ll \tau$ :
  - Use method of Green's functions:

$$\tilde{D}(\xi, \tau) = \int_{\xi}^1 d\xi_1 \int_0^{\tau} d\tau_1 G(\xi, \xi_1, \tau - \tau_1) I(\xi_1, \tau_1),$$

- Get

$$\tilde{D}(\xi, \tau) \approx \frac{G(\hat{q}_z / \hat{q}_y)}{3} \frac{\tau e^{-\pi\tau^2}}{\sqrt{\xi}}$$