

Lévy α -stable generalization of the ReBB model

based on [Universe 2023, 9\(8\), 361](#) & [Universe 2024, 10\(3\), 127](#)

T. Csörgő^{1,2}, S. Hegyi², [I. Szanyi](#)^{1,2,3}

¹MATE KRC, Gyöngyös, Hungary

²Wigner RCP, Budapest, Hungary

³Eötvös University, Budapest, Hungary

Diffraction and gluon saturation at the LHC and the EIC
10-14 June 2024, Trento, Italy

Outline

- **preliminaries**
- **the $p=(q,d)$ Bialas-Bzdak model and its extended version**
- **motivation for Lévy α -stable generalization**
- **Lévy α -stable generalization of the Bialas-Bzdak model**
- **an approximate simple Lévy α -stable model and fits to data**
- **relation between the parameters of the simple Lévy α -stable model and the full generalized model**

Preliminaries: ReBB model analysis of pp and p \bar{p} data

- the Real extended Bialas-Bzdak (ReBB) model describes elastic pp and p \bar{p} $d\sigma/dt$ data in a statistically acceptable way (CL \geq 0.1%) in the kinematic region:


$$546 \text{ GeV} \leq \sqrt{s} \leq 8 \text{ TeV}$$

$$0.38 \text{ GeV}^2 \leq -t \leq 1.2 \text{ GeV}^2$$

- significant model dependent odderon signal is observed
- main goal:** to improve the ReBB model to have a statistically acceptable (CL \geq 0.1%) description to elastic pp and p \bar{p} data in a wider kinematic range

T. Csörgő, I. Szanyi, *Eur. Phys. J. C* **81**, 611 (2021)

I. Szanyi, T. Csörgő, *Eur. Phys. J. C* **82**, 827 (2022)



ReBB model description to the 8 TeV pp data

Unitarity and the elastic scattering amplitude

- the unitarity of the S -matrix expresses the conservation of probability

$$SS^\dagger = I$$

- the unitarity constraint in impact parameter (b) representation at high energies is

$$2 \operatorname{Im} t_{el}(s, b) = |t_{el}(s, b)|^2 + \tilde{\sigma}_{in}(s, b) \quad (\sqrt{s} \text{ is the CM energy})$$

- the elastic scattering amplitude $t_{el}(s, b)$ can be written as a solution of the unitarity equation in terms of the inelastic cross section $\tilde{\sigma}_{in}(s, b)$

$$0 \leq \tilde{\sigma}_{in}(s, b) \leq 1$$

- at a given energy $\tilde{\sigma}_{in}(s, b)$ is the probability of inelastic scattering as a function of b and it can be calculated by using probability calculus and R. J. Glauber's multiple diffractive scattering theory


The Bialas-Bzdak (BB) $p=(q,d)$ model

A. Bialas, A. Bzdak, *Acta Phys.Polon. B 38, 159-168 (2007)*

- in the **Bialas-Bzdak (BB) $p=(q,d)$ model** the proton is a **bound state of a constituent quark and constituent a diquark**
- the inelastic scattering probability of two protons at a fixed impact parameter vector (\vec{b}) and at fixed constituent transverse position vectors ($\vec{s}_q, \vec{s}_d, \vec{s}'_q, \vec{s}'_d$) is given by a **Glauber expansion**:

$$\sigma(\vec{s}_q, \vec{s}_d; \vec{s}'_q, \vec{s}'_d; \vec{b}) = 1 - \prod_{a \in \{q,d\}} \prod_{b \in \{q,d\}} [1 - \sigma_{ab}(\vec{b} + \vec{s}'_b - \vec{s}_a)]$$

- $\sigma_{ab}(\vec{x}) \equiv \frac{d^2\sigma_{ab}(\vec{x})}{dx^2}$ is the inelastic differential cross section (inelastic scattering probability) for the collision of two constituents at a fixed relative transverse position \vec{x} of the constituents
- the **Glauber expansion sums the probabilities of all possible single and multiple binary inelastic collisions of the constituents (back scattering is prohibited)**
- the collision of two protons is inelastic if at least one constituent-constituent collision is inelastic



Proton-proton collision in the quark-diquark model

- the **probability of inelastic scattering** of protons at a fixed impact parameter vector (\vec{b}) is given by averaging over the constituent positions inside the protons:

$$\tilde{\sigma}_{in}(\vec{b}) = \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} d^2 s_q d^2 s'_q d^2 s_d d^2 s'_d D(\vec{s}_q, \vec{s}_d) D(\vec{s}'_q, \vec{s}'_d) \sigma(\vec{s}_q, \vec{s}_d; \vec{s}'_q, \vec{s}'_d; \vec{b})$$

- $D(\vec{s}_q, \vec{s}_d)$ is the (transverse) distribution of constituents inside a proton
- the scattering amplitude in the original BB model is considered to be completely imaginary by neglecting its relatively small real part

$$\tilde{t}_{el}(s, b) = i \left(1 - \sqrt{1 - \tilde{\sigma}_{in}(s, b)} \right)$$

$$b = |\vec{b}|$$

$$T(s, t) = 2\pi \int_0^{\infty} \tilde{t}_{el}(s, b) J_0(qb) b db$$

$$q = \sqrt{-t}$$

- the s -dependence of the amplitude happens through the s -dependencies of the model parameters

The Bialas-Bzdak (BB) $p=(q,d)$ model

A. Bialas, A. Bzdak, *Acta Phys.Polon.*
B 38, 159-168 (2007)

- in the original BB model the distribution of constituents inside a proton is given in terms of products of Gaussians

$$D(\vec{s}_q, \vec{s}_d) = \frac{1 + \lambda^2}{R_{qd}^2 \pi} e^{-\frac{\vec{s}_q^2}{R_{qd}^2}} e^{-\frac{\vec{s}_d^2}{R_{qd}^2}} \delta^2(\vec{s}_d + \lambda \vec{s}_q)$$

$$\lambda = \frac{m_q}{m_d}$$

$$\vec{s}_d = -\lambda \vec{s}_q$$

$$\vec{s}'_d = -\lambda \vec{s}'_q$$

- the constituent-constituent inelastic differential cross sections have also Gaussian shapes

$$\sigma_{ab}(\vec{x}) = A_{ab} e^{-\frac{\vec{x}^2}{R_a^2 + R_b^2}}$$

$$a, b \in \{q, d\}$$

- the constituent-constituent inelastic integrated cross sections are

$$\sigma_{ab}^{int} = \iint \sigma_{ab}(\vec{x}) d^2x = \pi A_{ab} (R_a^2 + R_b^2)$$

- assuming that the diquark contains twice as many partons than the quark and the colliding constituents do not shadow each other, $\sigma_{qq}^{int} : \sigma_{qd}^{int} : \sigma_{dd}^{int} = 1 : 2 : 4$ and the number of free parameters reduces to five: A_{qq} , λ , R_q , R_d , and R_{qd}

Real extended Bialas-Bzdak (ReBB) model

- in the original BB model the differential cross section is zero around the position of the diffractive minimum
- as a solution, the elastic scattering amplitude was extended in a unitary manner leading to the Real extended Bialas-Bzdak (ReBB) model

F. Nemes, T. Csörgő, M. Csanád, Int. J. Mod. Phys. A Vol. 30, 1550076 (2015)

$$\begin{array}{l} \tilde{t}_{el}(s, b) = i[1 - e^{-\Omega(s, b)}] \\ \text{Re}\Omega(s, b) = -1/2 \ln[1 - \tilde{\sigma}_{in}(s, b)] \\ \text{Im}\Omega(s, b) = 0 \end{array} \quad \longrightarrow \quad \begin{array}{l} \text{new free parameter} \\ \text{Im}\Omega(s, \vec{b}) = -\alpha_R \tilde{\sigma}_{in}(s, \vec{b}) \end{array}$$

$$\tilde{t}_{el}(s, b) = i \left(1 - \sqrt{1 - \tilde{\sigma}_{in}(s, b)} \right) \quad \longrightarrow \quad \tilde{t}_{el}(s, b) = i \left(1 - e^{i \alpha_R \tilde{\sigma}_{in}(s, b)} \sqrt{1 - \tilde{\sigma}_{in}(s, b)} \right)$$

- the ReBB model gives a statistically acceptable description (CL \geq 0.1%) to elastic pp and p \bar{p} scattering in the kinematic region:

$$0.546 \text{ TeV} \leq \sqrt{s} \leq 8 \text{ TeV} \quad \& \quad 0.38 \text{ GeV}^2 \leq -t \leq 1.2 \text{ GeV}^2$$

T. Csörgő, I. Szanyi, Eur. Phys. J. C 81, 611 (2021)

I. Szanyi, T. Csörgő, Eur. Phys. J. C 82, 827 (2022)

- the main goal is to have a statistically acceptable description in a wider kinematic range
- the TOTEM measurement at LHC at $\sqrt{s} = 8$ TeV excluded a purely exponential pp differential cross-section in the range of four-momentum transfer squared $0.027 \text{ GeV}^2 \leq -t \leq 0.2 \text{ GeV}^2$ with a significance greater than 7σ TOTEM Collab., *Nucl. Phys. B*, 899, 527 (2015)
- a simple model with Gaussian impact parameter amplitude yields a purely exponential t-distribution while a simple model with Levy α -stable impact parameter amplitude and $\alpha_L < 2$ yields a non-exponential t-distribution

$$\tilde{T}_{el}(s, b) = \frac{i + \rho_0(s)}{2} \sigma_{tot}(s) \frac{1}{2\pi B_0(s)} e^{-\frac{1}{2} \frac{b^2}{B_0(s)}} \quad \rightarrow \quad \tilde{T}_{el}(s, b) = \frac{i + \rho_0(s)}{2} \sigma_{tot}(s) \frac{1}{4\pi^2} \int d^2 q e^{-i\vec{q} \cdot \vec{b}} e^{-\frac{1}{2} |q^2 B_L(s)|^{\alpha_L(s)/2}}$$

$$T_{el}(s, t) = \int d^2 b e^{i\vec{q} \cdot \vec{b}} \tilde{T}_{el}(s, b) \quad \frac{d\sigma_{el}}{dt}(s, t) = \frac{1}{4\pi} |T_{el}(s, t)|^2 \quad a(s) = \frac{1 + \rho_0^2(s)}{16\pi} \sigma_{tot}^2(s)$$

$$\frac{d\sigma_{el}}{dt}(s, -t) = a(s) e^{-t B_0(s)} \quad \rightarrow \quad \frac{d\sigma_{el}}{dt}(s, -t) = a(s) e^{-|t B_L(s)|^{\alpha_L(s)/2}}$$


Lévy α -stable distributions in HEP

- the application of Lévy α -stable distributions is not new in the field of high-energy physics
- the Cauchy-Lorentz or Breit-Wigner distribution ($\alpha_L = 1$ case) is used to model unstable particles
- the Lévy expansion technique was applied to describe elastic pp scattering also at 13 TeV

T. Csörgő, R. Pasechnik, A. Ster, *Eur. Phys. J. C* 79, 62 (2019)

- the application of stable distributions is widespread in heavy ion physics

M. Csanád, D. Kincses, *Universe* 10 (2024) 2, 54



Description to pp elastic differential cross section data at 13 TeV using the Lévy expansion technique

Gaussian vs Lévy α -stable distribution

- the bivariate Gaussian distribution centered at $\vec{0}$ is

$$G(\vec{x}|R_G) = \frac{1}{2\pi R_G^2} e^{-\frac{\vec{x}^2}{2R_G^2}}$$


- the bivariate symmetric Levy α -stable distribution centered at $\vec{0}$ is

$$L(\vec{x}|\alpha_L, R_L) = \frac{1}{(2\pi)^2} \int d^2 q e^{-i\vec{q}\cdot\vec{x}} e^{-|\vec{q}^2 R_L^2|^{\alpha_L/2}}$$

$$0 < \alpha_L \leq 2$$

- for $\alpha_L = 2$ the the Lévy α -stable distribution is the Gaussian distribution

$$L(\vec{x}|\alpha_L = 2, R_L = R_G/\sqrt{2}) \equiv G(\vec{x}|R_G)$$



The bivariate symmetric Levy α -stable distribution for $R_L = 1$ as a function of $x = |x|$

Lévy α -stable distributions with $\alpha_L < 2$ have tails behaving asymptotically as a power law (infinite variance)

- the inelastic differential cross section for the collision of two constituents can be written **in terms of a convolution of their parton distributions**
- in the original BB model the parton distributions of the constituents are Gaussian distributions


$$\sigma_{ab}(\vec{x}) = A_{ab} \pi S_{ab}^2 \int d^2 r_a G(\vec{r}_a | R_a / \sqrt{2}) G(\vec{x} - \vec{r}_a | R_b / \sqrt{2})$$

$$\equiv A_{ab} \pi S_{ab}^2 G(\vec{x} | S_{ab} / \sqrt{2})$$

$$\vec{x} = \vec{b} + \vec{s}'_b - \vec{s}_a$$

$$S_{ab}^2 = R_a^2 + R_b^2$$

$$a, b \in \{q, d\}$$



The picture of the proton in the quark-diquark model


The quark-diquark distribution

- in the original BB model $D(\vec{s}_q, \vec{s}_d)$, the distribution of constituents inside a proton is given in terms of products of Gaussians
- **considering the relative distance** between the quark and diquark ($\vec{s}_q - \vec{s}_d$) one can write $D(\vec{s}_q, \vec{s}_d)$ in terms of a single Gaussian distribution:

$$D(\vec{s}_q, \vec{s}_d) = (1 + \lambda)^2 G(\vec{s}_q - \vec{s}_d | R_{qd}/\sqrt{2}) \delta^2(\vec{s}_q + \lambda \vec{s}_d)$$

$$\lambda = m_q/m_d$$

- the Dirac δ fixes the center of the mass of the proton making the calculations easier
- $D(\vec{s}_q, \vec{s}_d)$ is normalized as $\int d^2 s_q d^2 s_d D(\vec{s}_q, \vec{s}_d) = 1$



The picture of the proton in the quark-diquark model

Lévy α -stable generalized Bialas-Bzdak (LBB) model

Universe 2023, 9(8), 361

- **the parton distributions of the constituent quark and diquark are now Levy α -stable distributions** and the inelastic differential cross section for the collision of two constituents is:

$$\sigma_{ab}(\vec{x}) = A_{ab}\pi S_{ab}^2 \int d^2r_a L(\vec{r}_a|\alpha_L, R_a/2)L(\vec{x} - \vec{r}_a|\alpha_L, R_b/2) \equiv A_{ab}\pi S_{ab}^2 L(\vec{x}|\alpha_L, S_{ab}/2)$$

$$S_{ab}^{\alpha_L} = R_a^{\alpha_L} + R_b^{\alpha_L}$$

- **the distribution of the constituents inside the proton is now given in terms of a Levy α -stable distribution:**

$$D(\vec{s}_q, \vec{s}_d) = (1 + \lambda)^2 L(\vec{s}_q - \vec{s}_d|\alpha_L, R_{qd}/2)\delta^2(\vec{s}_q + \lambda\vec{s}_d)$$

$$\int d^2s_q d^2s_d D(\vec{s}_q, \vec{s}_d) = 1$$

α_L is a new free parameter of the model and if $\alpha_L = 2$ the BB model with Gaussian distributions is recovered

Difficulties with LBB model

- $\tilde{\sigma}_{in}(\vec{b})$ can be written as sum of 11 different terms that are integrals of products of Lévy α -stable distributions

$$\tilde{\sigma}_{in}(\vec{b}) = \tilde{\sigma}_{in}^{qq}(\vec{b}) + 2\tilde{\sigma}_{in}^{qd}(\vec{b}) + \tilde{\sigma}_{in}^{dd}(\vec{b}) - [2\tilde{\sigma}_{in}^{qq,qd}(\vec{b}) + \tilde{\sigma}_{in}^{qd,dq}(\vec{b}) + \tilde{\sigma}_{in}^{qq,dd}(\vec{b}) + 2\tilde{\sigma}_{in}^{qd,dd}(\vec{b})] \\ + [\tilde{\sigma}_{in}^{qq,qd,dq}(\vec{b}) + 2\tilde{\sigma}_{in}^{qq,qd,dd}(\vec{b}) + \tilde{\sigma}_{in}^{dd,qd,dq}(\vec{b})] - \tilde{\sigma}_{in}^{qq,qd,dq,dd}(\vec{b})$$

- difficulties with the calculation of integrals of products of Lévy α -stable distributions
- the calculation is easy only if the integral can be written in a convolution form as in case of the leading order terms in $\tilde{\sigma}_{in}(s, \vec{b})$

Leading order terms in $\tilde{\sigma}_{in}$ in the LBB model

Universe 2023, 9(8), 361

$$\begin{aligned}\tilde{\sigma}_{in}^{qq}(\vec{b}) &= \pi A_{qq} (2R_q^{\alpha_L})^{2/\alpha_L} \times \\ &\times \int d^2 s_q d^2 s'_q L(\vec{s}_q | \alpha_L, R_{qd^*}/2) L(\vec{s}'_q | R_{qd^*}/2) L(\vec{b} + \vec{s}'_q - \vec{s}_q | (2R_q^{\alpha_L})^{1/\alpha_L}/2) \\ &= \pi A_{qq} (2R_q^{\alpha_L})^{2/\alpha_L} L(\vec{b} | \alpha_L, (2R_{qd^*}^{\alpha_L} + 2R_q^{\alpha_L})^{1/\alpha_L}/2),\end{aligned}$$

$$\begin{aligned}\tilde{\sigma}_{in}^{qd}(\vec{b}) &= 2\pi A_{qq} (2R_q^{\alpha_L})^{2/\alpha_L} \times \\ &\times \int d^2 s_q d^2 s'_q L(\vec{s}_q | R_{qd^*}/2) L(\vec{s}'_q | R_{qd^*}/2) L(\vec{b} - \lambda \vec{s}'_q - \vec{s}_q | \alpha_L, (R_q^{\alpha_L} + R_d^{\alpha_L})^{1/\alpha_L}/2) \\ &= 2\pi A_{qq} (2R_q^{\alpha_L})^{2/\alpha_L} L(\vec{b} | \alpha_L, ((1 + \lambda^{\alpha_L})R_{qd^*}^{\alpha_L} + R_q^{\alpha_L} + R_d^{\alpha_L})^{1/\alpha_L}/2),\end{aligned}$$

$$\begin{aligned}\tilde{\sigma}_{in}^{dd}(\vec{b}) &= 4\pi A_{qq} (2R_q^{\alpha_L})^{2/\alpha_L} \times \\ &\times \int d^2 s_q d^2 s'_q L(\vec{s}_q | R_{qd^*}/2) L(\vec{s}'_q | R_{qd^*}/2) L(\vec{b} + \lambda(\vec{s}_q - \vec{s}'_q) | \alpha_L, (2R_d^{\alpha_L})^{1/\alpha_L}/2) \\ &= 4\pi A_{qq} (2R_q^{\alpha_L})^{2/\alpha_L} L(\vec{b} | \alpha_L, (2\lambda^{\alpha_L} R_{qd^*}^{\alpha_L} + 2R_d^{\alpha_L})^{1/\alpha_L}/2).\end{aligned}$$

Difficulties with LBB model fits to the data

- since multivariate Lévy α -stable distributions can be given only in terms of special functions, it is hard to perform a numerical fitting procedure
- a relatively high computing capacity and improved analytic insight is needed to proceed with the full model
- **quick solution:** approximations that are valid at the low $-t$ domain
- at low $-t$ values, the original ReBB model had difficulties to describe the strongly non-exponential features of the experimental data on $d\sigma/dt$
- a simple model which is valid at the low $-t$ domain easily illustrates the power of the Lévy α -stable generalization

Simple Lévy α -stable model for low- $|t|$ pp $d\sigma/dt$

Universe 2023, 9(8), 361

- low- $|t|$ scattering corresponds to high- b scattering and at high b values $\tilde{\sigma}_{in}(s, b)$ is small
- leading order term in the Taylor expansion of the amplitude in $\tilde{\sigma}_{in}(s, b)$ dominates at low $-t$ values if α_R is small too

$$\tilde{t}_{el}(s, b) = i \left(1 - e^{i \alpha_R(s) \tilde{\sigma}_{in}(s, b)} \sqrt{1 - \tilde{\sigma}_{in}(s, b)} \right) \longrightarrow \tilde{t}_{el}(s, b) = \left(\alpha_R(s) + \frac{i}{2} \right) \tilde{\sigma}_{in}(s, b)$$

- motivated by the fact that the leading order terms in $\tilde{\sigma}_{in}(s, \vec{b})$ have Lévy α -stable shapes in the LBB model, $\tilde{\sigma}_{in}(s, \vec{b})$ is approximated with a single Lévy α -stable shape

$$\tilde{\sigma}_{in}(s, \vec{b}) = \tilde{c}(s) L(\vec{b} | \alpha_L(s), r(s))$$



- **a simple Lévy α -stable model model for low- $|t|$ pp $d\sigma/dt$ arises**

$$t_{el}(s, t) = \int d^2 b e^{i\vec{q}\cdot\vec{b}} \tilde{t}_{el}(s, \vec{b}), |\vec{\Delta}| = \sqrt{-t} \longrightarrow \frac{d\sigma}{dt}(s, t) = \frac{1}{4\pi} |t_{el}(s, t)|^2 = a(s) e^{-|tb(s)|^{\alpha_L(s)/2}}$$

- the model has three adjustable parameters, α_L , a , and b , to be determined at a given energy

Simple Lévy α -stable model and the data

Universe 2023, 9(8), 361



- the non-exponential Lévy α -stable model with $\alpha_L = 1.953 \pm 0.004$ represents the LHC TOTEM $\sqrt{s} = 8 \text{ TeV}$ low- $|t|$ differential cross section data with a confidence level of 55% (published)
- similarly good description is obtained to all the LHC data on low- $|t|$ pp (and $p\bar{p}$) $d\sigma/dt$

Fits with simple Lévy α -stable model

Universe 2024, 10(3), 127

- fits to the existing pp and p \bar{p} $d\sigma/dt$ data in the kinematic range:

$$546 \text{ GeV} \leq \sqrt{s} \leq 13 \text{ TeV}$$

$$0.02 \text{ GeV}^2 \leq -t \leq 0.15 \text{ GeV}^2$$



- the CL values of the fits range between 8.8% and 96%.
- statistical, systematic and normalization errors are taken into account using the χ^2 definition developed by PHENIX Collab.

\sqrt{s} , GeV	α_L	a , mb/GeV ²	b , GeV ⁻²	CL, %
546	1.93 \pm 0.09	209 \pm 15	15.8 \pm 0.9	18.1
1800	2.0 \pm 1.5	270 \pm 24	16.2 \pm 0.2	77.1
2760	1.600 \pm 0.3	637 \pm 252	28 \pm 11	20.5
7000 (T)	1.95 \pm 0.01	535 \pm 30	20.5 \pm 0.2	8.8
7000 (A)	1.97 \pm 0.01	463 \pm 13	19.8 \pm 0.2	96.0
8000 (T1)	1.955 \pm 0.005	566 \pm 31	20.09 \pm 0.08	43.86
8000 (T2)	1.90 \pm 0.03	582 \pm 33	20.9 \pm 0.4	19.6
8000 (A)	1.97 \pm 0.01	480 \pm 11	19.9 \pm 0.1	55.8
13000 (T1)	1.959 \pm 0.006	677 \pm 36	20.99 \pm 0.08	76.5
13000 (T2)	1.958 \pm 0.003	648 \pm 10	21.06 \pm 0.05	89.1
13000 (A)	1.968 \pm 0.006	569 \pm 17	20.84 \pm 0.07	29.7

Values of the fitted parameters of the simple Lévy- α stable model at different energies

A. Adare *et al.* (PHENIX Collab.) *Phys. Rev. C* 77, 064907

$\alpha_L = 2$ versus $\alpha_L < 2$ results @ 13 TeV




Energy dependence of the α_L parameter

Universe 2024, 10(3), 127

- the value of the α_L parameter does not depend on energy
- its value is 1.959 ± 0.002 , i.e., slightly but in a statistical sense significantly different from 2

→ strong non-exponential behavior at low $-t$ in the differential cross section, power law tail at high- \vec{b} in $\tilde{\sigma}_{in}(s, \vec{b})$



Energy dependence of the α_L parameter of the simple Lévy- α stable model

Energy dependence of the optical point parameter


Universe 2024, 10(3), 127

- the energy dependence of the a parameter is quadratically logarithmic:

$$a(s) = p_0 + p_1 \ln \frac{s}{1 \text{ GeV}^2} + p_2 \ln^2 \frac{s}{1 \text{ GeV}^2}$$

- ATLAS and TOTEM data result slightly different energy dependencies
- reason: ATLAS and TOTEM use different methods to obtain the absolute normalization of the measurements

ATLAS Collab., *Eur. Phys. J. C* 83 (2023) 441



Energy dependence of the a parameter of the simple Lévy- α stable model

Energy dependence of the slope parameter


Universe 2024, 10(3), 127

- the energy dependence of the b parameter for TOTEM and ATLAS data together, and for $p\bar{p}$ data alone are linearly logarithmic:

$$b(s) = p_0 + p_1 \ln \frac{s}{1 \text{ GeV}^2}$$

- the LHC pp and the lower energy $p\bar{p}$ data do not lie on the same curve
- reason: the slope parameter data have a jump in the energy dependence around 3-4 GeV

TOTEM Collab., *Eur. Phys. J. C* (2019) 79:103



Energy dependence of the b parameter of the simple Lévy- α stable model

Simple Lévy α -stable & LBB model parameters

- parameters of the simple Levy α -stable model and the measurable quantities at $t \rightarrow 0$ can be approximately expressed in terms of the parameters of the LBB model *Universe 2023, 9(8), 361*
- only leading order terms in $\tilde{\sigma}_{in}(s, \vec{b})$ are considered; $A_{qq} = 1$ and $\lambda = 1/2$ are fixed

$$\frac{d\sigma}{dt}(s, t = 0) = a(s) = \frac{81}{16} \pi \left(2R_q^{\alpha_L(s)}(s) \right)^{4/\alpha_L} (1 + 4\alpha_R^2(s))$$

$$b(s) = \frac{1}{36} \left(\frac{4}{3} \right)^{2/\alpha_L(s)} \left((2 + 2^{\alpha_L(s)}) R_{qd}^{\alpha_L(s)}(s) + 3^{\alpha_L(s)} \left(2R_d^{\alpha_L(s)}(s) + R_q^{\alpha_L(s)}(s) \right) \right)^{2/\alpha_L(s)}$$

(obtained after a Taylor expansion in $t^{\alpha_L/2}$)

$$\sigma_{tot}(s) = 9\pi \left(2R_q^{\alpha_L(s)}(s) \right)^{2/\alpha_L(s)}$$

$$\rho_0(s) = \frac{Ret_{el}(s, t = 0)}{Imt_{el}(s, t = 0)} = 2\alpha_R$$

$$\sigma_{el}(s) = \frac{a(s)}{b(s)} \Gamma \left(\frac{2 + \alpha_L(s)}{\alpha_L(s)} \right)$$

- according to the analysis of elastic pp and $p\bar{p}$ data in the energy region $0.5 \text{ TeV} \leq \sqrt{s} \leq 8 \text{ TeV}$ only α_R is different for pp and $p\bar{p}$ scattering (T. Csörgő, I. Szanyi, *Eur. Phys. J. C* **81**, 611 (2021))
- in the low- $|t|$ approximation, difference between pp and $p\bar{p}$ scattering could be seen in the data on $d\sigma/dt$, ρ_0 , a (optical point), and σ_{el} , no difference in the data on σ_{tot} and b

Summary

- the formal Lévy α -stable generalization of the Bialas-Bzdak model is done, the $\alpha_L = 2$ limit corresponds to the original model
- solution of difficult and complex technical (mathematical and computational) problems is needed to fit the experimental data with the generalized model
- based on approximations a highly simplified Levy α -stable model of the pp (and $p\bar{p}$) differential cross section is deduced and successfully fitted to the data in the low- $|t|$ region
- the energy dependences of the parameters of the simple model are determined; the parameters of the simple model are related to the parameters of the Lévy α -stable generalized real extended Bialas-Bzdak (LBB) model
- final conclusion: the successful fit results indicate promising prospect for the future utility of the LBB model in describing experimental data



Thank you for your attention!

Backup slides

ρ_0 & α_R : connection between $t = 0$ and $t \neq 0$ data

- there is a connection between the ρ_0 parameter and the α_R parameter of the ReBB model regulating the real part of the scattering amplitude and the minimum-maximum structure of the $d\sigma/dt$
- α_R is determined by the $d\sigma/dt$ data at the minimum-maximum region but at the same time specifies the value of the ρ_0 in the ReBB model

T. Csörgő, I. Szanyi, *Eur. Phys. J. C* **81**, 611 (2021)



Most general term in $\tilde{\sigma}_{in}$

$$\tilde{\sigma}_{in}^{qq,qd,dq,dd}(\vec{b}) = \int d^2 s_q d^2 s'_q L(\vec{s}_q | R_{qd^*}/2) L(\vec{s}'_q | R_{qd^*}/2) \times \sigma_{qq}(\vec{s}_q, \vec{s}'_q; \vec{b}) \sigma_{qd}(\vec{s}_q, -\lambda \vec{s}'_q; \vec{b}) \sigma_{dq}(\vec{s}'_q, -\lambda \vec{s}_q; \vec{b}) \sigma_{dd}(-\lambda \vec{s}_q, -\lambda \vec{s}'_q; \vec{b})$$

$$\sigma_{qq}(\vec{s}_q, \vec{s}'_q; \vec{b}) = \pi A_{qq} (2R_q^\alpha)^{2/\alpha} \times L(\vec{b} + \vec{s}'_q - \vec{s}_q | \alpha, (2R_q^\alpha)^{1/\alpha} / \sqrt{2})$$



$$\sigma_{qd}(\vec{s}_q, \vec{s}'_d; \vec{b}) = 2\pi A_{qq} (2R_q^\alpha)^{2/\alpha} \times L(\vec{b} + \vec{s}'_d - \vec{s}_q | \alpha, (R_q^\alpha + R_d^\alpha)^{1/\alpha} / \sqrt{2})$$

$$\sigma_{dd}(\vec{s}_d, \vec{s}'_d; \vec{b}) = 4\pi A_{qq} (2R_q^\alpha)^{2/\alpha} \times L(\vec{b} + \vec{s}'_d - \vec{s}_d | \alpha, (2R_d^\alpha)^{1/\alpha} / 2)$$

$$\sigma_{dq}(\vec{s}_d, \vec{s}'_q; \vec{b}) = 2\pi A_{qq} (2R_q^\alpha)^{2/\alpha} \times L(\vec{b} + \vec{s}'_q - \vec{s}_d | \alpha, (R_q^\alpha + R_d^\alpha)^{1/\alpha} / 2)$$

Energy dependences of the ReBB model parameters

T. Csörgő, I. Szanyi, *Eur. Phys. J. C* 81, 611 (2021)





The energy dependences of the scale parameters, $R_q(s)$, $R_d(s)$, and $R_{qd}(s)$ are **linear logarithmic** and the **same** for pp and p \bar{p} processes!

The energy dependence of the α parameter, $\alpha(s)$ is **linear logarithmic** too, but **not** the same for pp and p \bar{p} processes!

Energy dependences of the ReBB model parameters

T. Csörgő, I. Szanyi, *Eur. Phys. J. C* 81, 611 (2021)



The energy dependences of the scale parameters, $R_q(s)$, $R_d(s)$, and $R_{qd}(s)$ are linear logarithmic and the same for pp and $p\bar{p}$ processes!

The energy dependence of the α parameter, $\alpha(s)$ is linear logarithmic too, but not the same for pp and $p\bar{p}$ processes!

Fit method

- least squares fitting with the method developed by the PHENIX collaboration
- this method is **equivalent to the diagonalization of the covariance matrix** if the experimental errors are separated into three different types:
 - type A: point-to-point varying uncorrelated statistical and systematic errors
 - type B: point-to-point varying 100% correlated systematic errors
 - type C: point-independent, overall systematic uncertainties
- i.e least squares fitting with:

[A. Adare et al. \(PHENIX Collab.\)
Phys. Rev. C 77, 064907](#)

$$\chi^2 = \left(\sum_{j=1}^M \left(\sum_{i=1}^{n_j} \frac{(d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj} - th_{ij})^2}{\tilde{\sigma}_{ij}^2} \right) + \epsilon_{bj}^2 + \epsilon_{cj}^2 \right) + \left(\frac{d_{\sigma_{tot}} - th_{\sigma_{tot}}}{\delta\sigma_{tot}} \right)^2 + \left(\frac{d_{\rho_0} - th_{\rho_0}}{\delta\rho_0} \right)^2$$

$$\tilde{\sigma}_{ij}^2 = \tilde{\sigma}_{aij} \left(\frac{d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj}}{d_{ij}} \right)$$

$$\tilde{\sigma}_{kij} = \sqrt{\sigma_{kij}^2 + (d'_{ij}\delta_k t_{ij})^2}, \quad k \in \{a, b\}$$

- minimization with **CERN Root MINUIT**, parameter error estimation by **MINOS**.

Fit method

- the method takes into account (in M separately measured t ranges):
 - the t -dependent statistical (**type A**) and systematic (**type B**) errors (both vertical σ_k and horizontal $\delta_k t$) $\rightarrow \epsilon_b$ parameters;
 - the t -independent σ_c normalization uncertainties (**type C**) $\rightarrow \epsilon_c$ parameters;
 - the measured total cross-section $d_{\sigma_{tot}}$ and ratio d_{ρ_0} and their total uncertainties $\delta\sigma_{tot}$ and $\delta\rho_0$.

[A. Adare et al. \(PHENIX Collab.\)
Phys. Rev. C 77, 064907](#)

- i.e least squares fitting with:**

$$\chi^2 = \left(\sum_{j=1}^M \left(\sum_{i=1}^{n_j} \frac{(d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj} - th_{ij})^2}{\tilde{\sigma}_{ij}^2} \right) + \epsilon_{bj}^2 + \epsilon_{cj}^2 \right) + \left(\frac{d_{\sigma_{tot}} - th_{\sigma_{tot}}}{\delta\sigma_{tot}} \right)^2 + \left(\frac{d_{\rho_0} - th_{\rho_0}}{\delta\rho_0} \right)^2$$

$$\tilde{\sigma}_{ij}^2 = \tilde{\sigma}_{aij} \left(\frac{d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj}}{d_{ij}} \right)$$

$$\tilde{\sigma}_{kij} = \sqrt{\sigma_{kij}^2 + (d'_{ij}\delta_k t_{ij})^2}, \quad k \in \{a, b\}$$

- minimization with **CERN Root MINUIT**, parameter error estimation by **MINOS**.

Fit method

- the method takes into account (in M separately measured t ranges):
 - the ϵ_i -s must be considered as both measurements and fit parameters not effecting the NDF (since they have known central value of zero and known standard deviation of one)
 - the measured total cross-section $d_{\sigma_{tot}}$ and ratio d_{ρ_0} and their total uncertainties $\delta\sigma_{tot}$ and $\delta\rho_0$.

A. Adare et al. (PHENIX Collab.)
Phys. Rev. C 77, 064907

- i.e least squares fitting with:

$$\chi^2 = \left(\sum_{j=1}^M \left(\sum_{i=1}^{n_j} \frac{(d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj} - th_{ij})^2}{\tilde{\sigma}_{ij}^2} + \epsilon_{bj}^2 + \epsilon_{cj}^2 \right) + \left(\frac{d_{\sigma_{tot}} - th_{\sigma_{tot}}}{\delta\sigma_{tot}} \right)^2 + \left(\frac{d_{\rho_0} - th_{\rho_0}}{\delta\rho_0} \right)^2 \right)$$

$$\tilde{\sigma}_{ij}^2 = \tilde{\sigma}_{aij} \left(\frac{d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj}}{d_{ij}} \right)$$

$$\tilde{\sigma}_{kij} = \sqrt{\sigma_{kij}^2 + (d'_{ij}\delta_k t_{ij})^2}, \quad k \in \{a, b\}$$

- minimization with **CERN Root MINUIT**, parameter error estimation by **MINOS**.

Fit method

The PHENIX method is validated by evaluating the χ^2 from a full covariance matrix fit of the $\sqrt{s} = 13$ TeV TOTEM differential cross-section data using the Lévy expansion method from [T. Csörgő, R. Pasechnik, & A. Ster, Eur. Phys. J. C 79, 62 \(2019\)](#).

- the t -independent σ_c normalization uncertainties $\rightarrow \epsilon_c$ parameters;
- the measured total cross-section $d_{\sigma_{tot}}$ and ratio d_{ρ_0} and their total uncertainties $\delta\sigma_{tot}$ and $\delta\rho_0$.

[A. Adare et al. \(PHENIX Collab.\)
Phys. Rev. C 77, 064907](#)

- i.e least squares fitting with:

$$\chi^2 = \left(\sum_{j=1}^M \left(\sum_{i=1}^{n_j} \frac{(d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj} - th_{ij})^2}{\tilde{\sigma}_{ij}^2} \right) + \epsilon_{bj}^2 + \epsilon_{cj}^2 \right) + \left(\frac{d_{\sigma_{tot}} - th_{\sigma_{tot}}}{\delta\sigma_{tot}} \right)^2 + \left(\frac{d_{\rho_0} - th_{\rho_0}}{\delta\rho_0} \right)^2$$

$$\tilde{\sigma}_{ij}^2 = \tilde{\sigma}_{aij} \left(\frac{d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj}}{d_{ij}} \right)$$

$$\tilde{\sigma}_{kij} = \sqrt{\sigma_{kij}^2 + (d'_{ij}\delta_k t_{ij})^2}, \quad k \in \{a, b\}$$

- minimization with **CERN Root MINUIT**, parameter error estimation by **MINOS**.

Fit method

The PHENIX method is validated by evaluating the χ^2 from a full covariance matrix fit of the $\sqrt{s} = 13$ TeV TOTEM differential cross-section data using the Lévy expansion method from [T. Csörgő, R. Pasechnik, & A. Ster, Eur. Phys. J. C 79, 62 \(2019\)](#).

The PHENIX method and the fit with the full covariance matrix result in the same minimum within one standard deviation of the fit parameters.

[A. Adare et al. \(PHENIX Collab.\)
Phys. Rev. C 77, 064907](#)

- i.e least squares fitting with:

$$\chi^2 = \left(\sum_{j=1}^M \left(\sum_{i=1}^{n_j} \frac{(d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj} - th_{ij})^2}{\tilde{\sigma}_{ij}^2} \right) + \epsilon_{bj}^2 + \epsilon_{cj}^2 \right) + \left(\frac{d_{\sigma_{tot}} - th_{\sigma_{tot}}}{\delta\sigma_{tot}} \right)^2 + \left(\frac{d_{\rho_0} - th_{\rho_0}}{\delta\rho_0} \right)^2$$

$$\tilde{\sigma}_{ij}^2 = \tilde{\sigma}_{aij} \left(\frac{d_{ij} + \epsilon_{bj}\tilde{\sigma}_{bij} + \epsilon_{cj}d_{ij}\sigma_{cj}}{d_{ij}} \right)$$

$$\tilde{\sigma}_{kij} = \sqrt{\sigma_{kij}^2 + (d'_{ij}\delta_k t_{ij})^2}, \quad k \in \{a, b\}$$

- minimization with **CERN Root MINUIT**, parameter error estimation by **MINOS**.

Proportionality between $\rho_0(s)$ and $\alpha(s)$

$$t_{el}(s, b) = i \left(1 - e^{i \alpha \tilde{\sigma}_{in}(s, b)} \sqrt{1 - \tilde{\sigma}_{in}(s, b)} \right)$$

$$\alpha \tilde{\sigma}_{in} \ll 1$$


$$\text{Im } t_{el}(s, b) \simeq \lambda(s) \exp\left(-\frac{b^2}{2R^2(s)}\right)$$



$$\rho_0(s) = \alpha(s) \left(2 - \frac{3}{2} \lambda(s) + \frac{1}{3} \lambda^2(s) \right)$$

$$\lambda(s) = \text{Im } t_{el}(s, b = 0)$$

→ by rescaling one can get additional α parameter values at energies where ρ_0 is measured (and vice versa)



The dependence of ρ_0/α on $\lambda = \text{Im } t_{el}(s, b = 0)$ in the TeV energy range. The data points are generated numerically by using the trends of the ReBB model scale parameters and the experimentally measured ρ -parameter values.

Measurable quantities

- differential cross section:

$$\frac{d\sigma}{dt}(s, t) = \frac{1}{4\pi} |T(s, t)|^2$$

- total, elastic and inelastic cross sections:

$$\sigma_{tot}(s) = 2\text{Im}T(s, t = 0)$$

$$\sigma_{el}(s) = \int_{-\infty}^0 \frac{d\sigma(s, t)}{dt} dt$$

$$\sigma_{in}(s) = \sigma_{tot}(s) - \sigma_{el}(s)$$

- ratio ρ_0 :

$$\rho_0(s) = \lim_{t \rightarrow 0} \rho(s, t) \equiv \frac{\text{Re}T(s, t \rightarrow 0)}{\text{Im}T(s, t \rightarrow 0)}$$

- slope of $d\sigma/dt$:

$$B(s, t) = \frac{d}{dt} \left(\ln \frac{d\sigma}{dt}(s, t) \right)$$

$$B_0(s) = \lim_{t \rightarrow 0} B(s, t)$$