

Entanglement entropy and proton's structure



Krzysztof Kutak



Motivation

Bounds and properties of EE may provide some new insight on behavior of pdfs

Links to other areas (thermodynamics, gravity, quantum information, conformal field theory)

Interesting in context of parton saturation and thermalization problem of Quark Gluon Plasma

Various approaches to entropy in the low x limit: entropy of gluon density, thermodynamic entropy, momentum space entanglement, coordinate space entanglement, Wehrl entropy,...

Based on:

Based on

Eur.Phys.J.C 82 (2022) 2, 111

M. Hentschinski, K. Kutak

Eur.Phys.J.C 82 (2022) 12, 1147

M. Hentschinski, K.Kutak, R. Straka

PRL'23

H. Hentschinski, D. Kharzeev, K. Kutak, Z. Tu

2404.07657

P. Caputa, K. Kutak

Boltzman and von Neuman entropy formulas – reminder

The entropy S of macrostate is given by the log of number W of distinct microstates that compose it

$$S = - \sum_{i=1}^W p(i) \ln p(i) \quad \text{Gibbs entropy}$$

For uniform distribution $p(i) = \frac{1}{W}$ the entropy is maximal Boltzmann entropy
 $S = \ln W$

Since partons are introduced as the microscopic constituents that compose the macroscopic state of the proton, it is natural to evaluate the corresponding entropy or entropy corresponding to parton density.

K. Kutak '11, Peschanski'12
 A. Kovner, M. Lublinsky '15
 D. Kharzeev, E. Levin '17,...

But proton as a whole is a pure state and the von Neuman entropy is 0. Can one get any nontrivial result?

For pure state (one state) density matrix is: For mixed state i.e. classical statistical mixture

$$\rho = |\psi\rangle\langle\psi|$$

$$S_{VN} = -\text{Tr}[\rho \ln \rho] = -1 \ln 1 = 0$$

$$\rho = \sum p(i) |\psi_i\rangle\langle\psi_i|$$

$$S_{VN} \neq 0$$

Kharzeev, Levin '17

Entanglement entropy in DIS

The composite system is described by

$$|\Psi_{AB}\rangle \text{ in } A \cap B$$

entangled

if the product can not be expressed as separable product state

$$|\Psi_{AB}\rangle = \sum_{i,j} c_{ij} |\varphi_i^A\rangle \otimes |\varphi_j^B\rangle$$

Schmidt decomposition

$$|\Psi_{AB}\rangle = \sum \alpha_n |\Psi_n^A\rangle |\Psi_n^B\rangle$$

orthonormal states belonging to A


orthonormal states belonging to B

related to matrix C

separable

if the product can be expressed as separable product state

$$|\Psi_{AB}\rangle = |\varphi^A\rangle \otimes |\varphi^B\rangle$$



Kharzeev, Levin '17

Entanglement entropy in DIS

$$|\Psi_{AB}\rangle = \sum_n \alpha_n |\Psi_n^A\rangle |\Psi_n^B\rangle$$


$$\rho_{AB} = |\Psi_{AB}\rangle \langle \Psi_{AB}|$$

$$\rho_A = \text{tr}_B \rho_{AB} = \sum_n \alpha_n^2 |\Psi_n^A\rangle \langle \Psi_n^A|$$

$\alpha_n^2 \equiv p_n$ probability of state with n partons

$$S = - \sum_n p_n \ln p_n$$

entropy results from the entanglement between the regions A and B, and can thus be interpreted as the entanglement entropy. Entropy of region A is the same as entropy in region B.





The density matrix of the mixed state probed in region A


Kharzeev, Levin '17

Momentum space vs coordinate space


momentum space - Bjorken frame



position space - Mueller frame



gluon \sim color dipole



from A. Stasto
Acta Phys.Polon.
B35 (2004) 3069-3102

$$\mathcal{F}(x, k) = \mathcal{F} + K_{ms} \otimes \mathcal{F}(x, k) - \frac{1}{R^2} TPV \otimes \mathcal{F}(x, k)^2 \quad N(x, r, b) = N_0 + K_{ps} \otimes (N(x, r, b) - N(x, r, b)^2)$$

dipole unintegrated gluon density

related by Fourier transform

Evolved with BK dipole amplitude – expectation value of product of Wilson lines in fundamental representation

Partonic, dipole cascade

$$p_n = P_n$$

$$\frac{dP_n(Y)}{dY} = -\lambda n P_n(Y) + (n-1)\lambda P_{n-1}(Y)$$

set of partons is described by set of dipoles with fixed sizes, Y is rapidity and is related to energy

Mueller 95, Lublinsky, Levin '03

$$P_n(Y) = e^{-\lambda Y} (1 - e^{-\lambda Y})^{n-1}$$

depletion of the probability to find n dipoles due to the splitting into $(n+1)$ dipoles.

the growth due to the splitting of $(n-1)$ dipoles into n dipoles.

$$S = -\sum_n p_n \ln p_n$$

model of BFKL dipole cascade

$$S(Y) \approx \lambda Y \quad \text{where} \quad Y = \ln 1/x$$

$$\langle n \rangle = \sum_n n P_n(Y) = \left(\frac{1}{x} \right)^\lambda$$

BFKL intercept = $4 \ln 2 \bar{\alpha}_S$

Kharzeev, Levin '17

Assumption $\langle n \rangle \equiv xg(x)$

The approach can be generalized to 3+1 d and one can account for hard scale dependence.

Density matrix in 1+1 D Nowak, Liu, Zahed '22

EE in DLL Nowak, Liu, Zahed '23

$$S(x) = \ln(xg(x))$$

$$S(x, Q) = \ln(xg(x, Q))$$

Entropy formula - interpretation

$$P_n(Y) = e^{-\lambda Y} (1 - e^{-\lambda Y})^{n-1}$$

At low x partonic microstates have equal probabilities

In this equipartitioned state the entropy is maximal – the partonic state at small x is maximally entangled.

In terms of information theory as Shannon entropy:

- equipartitioning in the maximally entangled state means that all “signals” with different number of partons are equally likely
- it is impossible to predict how many partons will be detected in a given event.
- structure function at small x should become universal for all hadrons.

From strict bounds on entanglement entropy (from conformal field theory) one can obtain that at low x (in conformal regime) one has

$$xg(x) \leq \text{const } x^{-1/3}$$

Kharzeev, Levin '17

Furthermore entropy of the final state hadrons can not be smaller than entropy of partons.

Comments

CFT result for EE

central charge

$$S = \frac{c}{3} \ln \frac{L}{\epsilon}$$

UV cutoff

Relation to Kharzeev-Levin formula

$$L = (mx)^{-1}$$


Length of tube
probed in DIS

$$S = \ln \left(\frac{1}{x} \right)^{1/3}$$

$$S(x) = \ln(xg(x))$$

$$\epsilon \equiv 1/m$$

Proton's Compton
wave length



Entanglement entropy obtained from CFT calculations as well as from gravity using Ryu-Takayanagi formula

See also
Callan, Wilczek '94
Calabrese, Cardy '04

and lectures by
Headrick

Studied also in the
context of 2 D QCD

Liu, Nowak, Zahed, '22

Casini, Huerta, Hosco '05

Entanglement entropy – calculation and measurement

For DIS at high energies, this entanglement entropy can be calculated using


$$S(x, Q^2) = \ln \left\langle n \left(\ln \frac{1}{x}, Q \right) \right\rangle$$

$$S_{hadron} = \sum P(N) \ln P(N)$$


N number of measured hadrons

The charged particle multiplicity distribution measured in either the current fragmentation region or the target fragmentation region.

Fraction of events with charged hadron



Data and EE



HERA pdf used

$$S(x, Q) = \ln(xg(x, Q))$$

Also attempt by Kharzeev and Levin
to use quarks instead of gluons
[Phys. Rev. D 104, 031503 \(2021\)](#)

H1

[Eur.Phys.J.C 81 \(2021\) 3, 212](#)

See also [Z. Tu, D. Kharzeev, T. Ulrich '20](#)
for calculations of EE in p-p.


Extension of KL entropy formula

Hentschinski, Kutak '21

$$\left\langle n \left(\ln \frac{1}{x}, Q \right) \right\rangle = xg(x, Q) + x\Sigma(x, Q)$$

To get the entropy of system of partons one needs to account for both quarks and gluons. One can view this as a higher order correction to KL formula. Furthermore it is impossible to isolate quarks from gluons therefore the complete entropy formula should receive contributions from quarks and gluons

Gluon and quark distribution



In the linear regime obeys BFKL equation. In our calculations we use NLO BFKL with kinematical improvements and running coupling. The gluon density has been fitted to F_2 data (exact kinematics was used)

Hentschinski, Sabio-Vera, Salas.
 Phys.Rev.D 87 (2013) 7, 076005
 Phys.Rev.Lett. 110 (2013) 4, 041601

We calculate the sea quarks distribution using

$$x\Sigma(x, Q) = P_{qg}(Q, \mathbf{k}) \otimes \mathcal{F}(x, \mathbf{k}^2)$$

$$xg(x, Q) = \int_0^{Q^2} d\mathbf{k}^2 \mathcal{F}(x, \mathbf{k}^2)$$

Transverse momentum dependent splitting function
 Catani, Hautmann
 Nucl.Phys. B427 (1994) 475-524

Other methods for resummation:
 KMS (Kwiecinski, Martin, Stasto);
 CCSS (Colferai, Ciafaloni, Staśto, Salam)

Gluon distribution

NLO BFKL with collinear resummation

$$\mathcal{F}(x, \mathbf{k}^2, Q) = \frac{1}{\mathbf{k}^2} \int_{\frac{1}{2}-i\infty}^{\frac{1}{2}+i\infty} \frac{d\gamma}{2\pi i} \hat{g}\left(x, \frac{Q^2}{Q_0^2}, \gamma\right) \left(\frac{\mathbf{k}^2}{Q_0^2}\right)^\gamma$$


$$\hat{g}\left(x, \frac{Q^2}{Q_0^2}, \gamma\right) = \frac{\mathcal{C} \cdot \Gamma(\delta - \gamma)}{\pi \Gamma(\delta)} \cdot \left(\frac{1}{x}\right)^{\chi(\gamma, Q, Q)} \left\{ 1 + \frac{\bar{\alpha}_s^2 \beta_0 \chi_0(\gamma)}{8N_c} \log\left(\frac{1}{x}\right) \left[-\psi(\delta - \gamma) + \log \frac{Q^2}{Q_0^2} - \partial_\gamma \right] \right\}$$

the low x growth





Hentschinski, Sabio-Vera, Salas.
 Phys.Rev.D 87 (2013) 7, 076005
 Phys.Rev.Lett. 110 (2013) 4, 041601

Proton structure function from HSS fit

F_2 data description







Results



Hint that the general idea works. Gluon dominates over quarks.
 One has to also take into account that only charged hadrons were measured.





Results





Hint that the general idea works. Gluon dominates over quarks.
 One has to also take into account that only charged hadrons were measured i.e 2/3 of partons contribute

Large scales - description

Martin Hentschinski, KK, Robert Straka '23





Small scales - prediction



See also [Hagivara, Hatta, Xiao '18](#)

The generalized KL model is used and entropy saturates in this approach
and [Nowak, Liu, Zahed '22](#)

Integrated gluon and entropy



$$\lim_{Q^2 \gg Q_s^2} S(x, Q^2) = \ln(S_{\perp} Q_s^2(x)) + \ln \frac{N_c}{8\alpha_s \pi^2} = \lambda \ln \frac{1}{x} + \text{const}$$

$$\lim_{Q^2 \ll Q_s^2} S(x, Q^2) = \ln \left(\frac{S_{\perp} Q^4}{Q_s^2(x)} \right) + \ln \frac{N_c}{16\alpha_s \pi^2}$$


Photon can not resolve proton anymore therefore the EE vanishes.

But it might be that the formalism breaks down for low scales.

There might be another source of entropy that keep the total entropy not vanishing → **generalized second law Bekenstein**

EE in Diffractive Deep Inelastic Scattering

H. Hentschinski, D. Kharzeev, K. Kutak, Z. Tu '23



$x_{\mathbb{P}}$ proton's momentum fraction carried by the Pomeron

β denotes the Pomeron's momentum fraction carried by the quark interacting with the virtual photon

$$x = \beta \cdot x_{\mathbb{P}} \quad \text{Bjorken } x$$

$$y_0 \simeq \ln 1/x_{\mathbb{P}} \quad \text{size of rapidity gap}$$



$$Y = \ln 1/x$$

$$y_X = Y - y_0 \simeq \ln 1/\beta$$

Analogous evolution equation as for non-diffractive case but Initial conditions are different and there is delay because of rapidity gap.


Munier, Mueller Phys. Rev. D 98, 034021 (2018)

Diffraction vs. nondiffraction



H1 detector

EE in DDIS



Asymptotic or maximal entangled region. All configurations have the same probability

High probability of configurations with few partons

H1 data region

$$\left\langle \frac{dn(\beta)}{d\beta} \right\rangle_{\text{charged}} \simeq \frac{2}{3} \left\langle \frac{dn(\beta)}{d\beta} \right\rangle$$


$$p_n^D(y_X) = \frac{1}{C} e^{-\Delta y_X} \left(1 - \frac{1}{C} e^{-\Delta y_X} \right)^{n-1}$$

$$p_n \equiv 1/Z$$

$$S(Z) = - \sum_n p_n \ln p_n = (1 - Z) \ln \frac{Z-1}{Z} + \ln Z$$

$$S_{\text{asym.}}(Z) = \ln Z + 1 + \mathcal{O}(1/Z)$$

EE in DDIS



momentum fraction carried by the
quark interacting with the virtual photon

Complexity and gluon density

P. Caputa, K. Kutak, 2404.07657

“The complexity of the task is defined as the minimum number of gates used to construct the circuit that accomplishes it” L. Susskind

The key point is to expand the state or the operator in the minimal basis that supports its unitary evolution

State complexity

$$|\Psi(t)\rangle = e^{-iHt} |\Psi_0\rangle = \sum_{n=0} \psi_n(t) |K_n\rangle$$

$$H |K_n\rangle = a_n |K_n\rangle + b_n |K_{n-1}\rangle + b_{n+1} |K_{n+1}\rangle$$

$$\mathcal{C}_K(t) = \langle n \rangle = \sum_n n p_n(t)$$

$$p_n(t) \equiv |\psi_n(t)|^2$$

$$|\psi(t)\rangle = e^{-i\alpha(L_1+L_{-1})t} |0\rangle \otimes |0\rangle$$

$$S_K(t) = - \sum_n p_n(t) \log p_n(t)$$

After transforming to rapidity variable one gets

boost operator

$$\partial_Y p_n(Y) = \alpha n p_{n-1}(Y) - \alpha(n+1) p_n(Y)$$


We can write the relation between the K-complexity and the gluon patron distribution

$$\mathcal{C}_K = xg(x)$$


QI measures and dipole equations

P. Caputa, K. Kutak, 2404.07657

quantum measures



quantum measures



$$\partial_Y p_n(Y) = -\lambda n p_n(Y) + \lambda(n-1)p_{n-1}(Y)$$

$$\begin{aligned} \partial_Y p_n(Y) = & -\lambda n p_n(Y) + \lambda(n-1)p_{n-1}(Y) \\ & + \beta n(n+1)p_{n+1}(Y) - \beta n(n-1)p_n(Y) \end{aligned}$$


Hagiwara, Hatta, Xiao, Yuan
Phys.Rev.D 97 (2018) 9, 094029

Conclusions and outlook

- We show evidences for the proposal for low x maximal entanglement entropy of proton constituents in .
- It can be systematically improved (quark contributions, NLO BFKL, rc BK) and can describe successfully H1 data.
- We obtain saturation of entropy at small resolution scales.
- We demonstrate that the proposal works for DDIS and that it can be used to study onset of maximal entanglement
- Saturation might be manifestation of provide mechanism for vanishing of EE at low resolution scale
- The thermodynamic based approach agrees with KL approach
- Formal study of dipole cascade model 2404.07657, P. Caputa, K. Kutak it follows that one can view gluon density as a measure of quantum complexity of proton

Backup

Unruh effect




Accelerated observer in its rest frame feels thermal radiation or Bose-Einstein distribution with temperature

$$T = \frac{|a|}{2\pi}$$

Colliding hadrons and Unruh effect

Stages of collision

Kharzeev, Tuchin '05



Hadron of mass m after interaction with external field transforms into hadrons of mass M

$$P(M \leftarrow m) = 2\pi |\mathcal{T}(M \leftarrow m)|^2 \rho(M),$$

probability for transition to final state

$$\int dM P(M \leftarrow m)$$

should be finite

density of states determined by typical momentum. Q_s emerges

$$|\mathcal{T}(M \leftarrow m)|^2 \sim \exp(-2\pi M/a)$$

$$T = \frac{Q_s(x)}{2\pi}$$

transition amplitude

Saturation and entropy

The relation $T = \frac{Q_s(x)}{2\pi}$ Can be understood in a generalized sense i.e. that **saturation** scale defines some **temperature**.

Equilibrium thermodynamics relations  Lower bound on produced entropy

It can be shown that the saturation line has an interpretation of a characteristics i.e. line along which the gluon density has a constant value.

Kutak 1103.3654.v1 and v2

$$dE = TdS$$

$$dM = TdS$$

$$\frac{dQ_s(x)}{Q_s(x)} = \frac{dS}{2\pi}$$


$$dE = dM$$

$$dM = dQ_s(x)$$

mass of system
of gluons

Gluon density and entropy

Kutak 1103.3654.v1 and v2



Many-body interactions



Medium generated mass of gluon.
Framework of Hard Thermal Loops.

Similarly in QED. Cut on photon's k_t
Is equivalent to introducing mass.


In presented approach **mass is not fixed it is x dependent**

$$\Delta S = \pi \lambda \Delta y$$

$$\Delta y = \ln(x_0/x)$$

$$Q_s^2 = Q_0^2 (x_0/x)^\lambda$$

Bining and KL formula



plot showing dependence of the result on the size of bins if binning is naive

Data binning takes place in rapidity

$$\bar{n}_g(\bar{x}) = \frac{1}{y_{\max} - y_{\min}} \int_{y_{\min}}^{y_{\max}} dy \frac{dn_g}{dy} = \frac{n_g(y_{\max}) - n_g(y_{\min})}{y_{\max} - y_{\min}}$$

$$y_{\max, \min} = \ln 1/x_{\min, \max}$$

for small bins

$$\bar{n}_g(x, Q^2) = \frac{dn_g}{d \ln(1/x)} = xg(x, Q^2)$$

$$\langle \bar{n}(x, Q^2) \rangle_{Q^2} = \frac{1}{Q_{\max}^2 - Q_{\min}^2} \int_{Q_{\min}^2}^{Q_{\max}^2} dQ^2 [xg(x, Q^2) + x\Sigma(x, Q^2)]$$

$$\langle S(x, Q^2) \rangle_{Q^2} = \ln \langle \bar{n}(x, Q^2) \rangle_{Q^2}$$

$$n_g(Q^2) = \int_0^1 dx g(x, Q^2)$$

Formal definition of number of gluons

$$n_g(\bar{x}) = \int_{x_{\min}}^{x_{\max}} dx g(x, Q^2)$$

$$\bar{x} \in [x_{\min}, x_{\max}]$$

$$\bar{x} = \frac{\int_{x_{\min}}^{x_{\max}} dx x g(x, Q^2)}{\int_{x_{\min}}^{x_{\max}} dx g(x, Q^2)}$$


average x

Gluon production and entropy – another assumptions

Kutak '11

Bialas; Janik; Fialkowski, Wit; Iancu, Blaizot, Peschanski,...

$\phi(x, k)$



Many-body interactions

Medium generated mass of gluon.
Framework of Hard Thermal Loops.

Similarly in QED. Cut on photon's kt
Is equivalent to introducing mass.

In presented approach mass is not fixed it is x dependent

$$M_G(x) = Q_s(x)$$

energy dependent
gluon's mass

$$M(x) = N_G(x) M_G(x)$$

mass of system
of gluons

$$N_G(x) \equiv \frac{dN}{dy} = \frac{1}{S_{\perp}} \frac{d\sigma}{dy}$$

number of gluons

$$dE = TdS$$

$$dM = TdS$$

$$d[N_G(x) M_G(x)] = \frac{Q_s(x)}{2\pi} dS$$

Entropy due to less
dense hadron

$$S = \frac{6C_F A_{\perp}}{\pi\alpha_s} Q_s^2(x) + S_0$$

$$S = 3\pi [N_G(x) + N_{G0}]$$