


www.cea.fr


Mapping PDAs and PDFs | Hervé MOUTARDE

 Sep. 14th, 2018

Covariant extension

Parameterizations

Definition
Polynomiality
Radon transform
Positivity
Inverse Radon
Examples

Computations

Design
Releases
Phenomenology

Fits

Status
Global CFF fit

Conclusion

Appendix

- 1 How can we parameterize GPDs?
- 2 How can we compute exclusive processes with increasing precision?
- 3 How can we fit GPDs from experimental data?

Parameterizing GPDs?

Covariant
extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status


Global CFF fit

Conclusion

Appendix

$$H_\pi^q(x, \xi, t) = \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \left\langle \pi, P + \frac{\Delta}{2} \right| \bar{q}\left(-\frac{z}{2}\right) \gamma^+ q\left(\frac{z}{2}\right) \left| \pi, P - \frac{\Delta}{2} \right\rangle_{z^+=0, z_\perp=0}$$

with $t = \Delta^2$ and $\xi = -\Delta^+/(2P^+)$.



References

- Müller *et al.*, Fortschr. Phys. **42**, 101 (1994)
Ji, Phys. Rev. Lett. **78**, 610 (1997)
Radyushkin, Phys. Lett. **B380**, 417 (1996)

■ PDF forward limit

$$H^q(x, 0, 0) = q(x)$$

Covariant
extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status


Global CFF fit

Conclusion

Appendix

$$H_\pi^q(x, \xi, t) = \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \left\langle \pi, P + \frac{\Delta}{2} \right| \bar{q}\left(-\frac{z}{2}\right) \gamma^+ q\left(\frac{z}{2}\right) \left| \pi, P - \frac{\Delta}{2} \right\rangle_{z^+=0, z_\perp=0}$$

with $t = \Delta^2$ and $\xi = -\Delta^+/(2P^+)$.



References

- Müller *et al.*, Fortschr. Phys. **42**, 101 (1994)
- Ji, Phys. Rev. Lett. **78**, 610 (1997)
- Radyushkin, Phys. Lett. **B380**, 417 (1996)

- PDF forward limit
- Form factor sum rule

$$\int_{-1}^{+1} dx H_\pi^q(x, \xi, t) = F_1^q(t)$$

Covariant
extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status


Global CFF fit

Conclusion

Appendix

$$H_\pi^q(x, \xi, t) = \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \left\langle \pi, P + \frac{\Delta}{2} \right| \bar{q}\left(-\frac{z}{2}\right) \gamma^+ q\left(\frac{z}{2}\right) \left| \pi, P - \frac{\Delta}{2} \right\rangle_{\substack{z^+ = 0 \\ z_\perp = 0}}$$

with $t = \Delta^2$ and $\xi = -\Delta^+/(2P^+)$.



References

- Müller *et al.*, Fortschr. Phys. **42**, 101 (1994)
- Ji, Phys. Rev. Lett. **78**, 610 (1997)
- Radyushkin, Phys. Lett. **B380**, 417 (1996)

- PDF forward limit
- Form factor sum rule
- H^q is an even function of ξ from time-reversal invariance.

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status


Global CFF fit

Conclusion

Appendix

$$H_\pi^q(x, \xi, t) = \frac{1}{2} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \left\langle \pi, P + \frac{\Delta}{2} \right| \bar{q}\left(-\frac{z}{2}\right) \gamma^+ q\left(\frac{z}{2}\right) \left| \pi, P - \frac{\Delta}{2} \right\rangle_{\substack{z^+ = 0 \\ z_\perp = 0}}$$

with $t = \Delta^2$ and $\xi = -\Delta^+/(2P^+)$.



References

- Müller *et al.*, Fortschr. Phys. **42**, 101 (1994)
- Ji, Phys. Rev. Lett. **78**, 610 (1997)
- Radyushkin, Phys. Lett. **B380**, 417 (1996)

- PDF forward limit
- Form factor sum rule
- H^q is an even function of ξ from time-reversal invariance.
- H^q is real from hermiticity and time-reversal invariance.

Covariant
extension

■ Polynomiality

$$\int_{-1}^{+1} dx x^n H^q(x, \xi, t) = \text{polynomial in } \xi$$

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

Covariant
extension

■ Polynomiality

Parameterizations

Definition

- Polynomiality
- Radon transform
- Positivity
- Inverse Radon
- Examples

Computations

- Design
- Releases
- Phenomenology

Fits

- Status
- Global CFF fit

Conclusion

Appendix

Lorentz covariance

Covariant
extension

■ Polynomiality

Lorentz covariance

Parameterizations

Definition

Polynomiality
Radon transform
Positivity
Inverse Radon
Examples

Computations

Design
Releases
Phenomenology

Fits

Status
Global CFF fit

Conclusion

Appendix

$$H^q(x, \xi, t) \leq \sqrt{q\left(\frac{x+\xi}{1+\xi}\right) q\left(\frac{x-\xi}{1-\xi}\right)}$$

Covariant extension

■ Polynomiality

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

Lorentz covariance

■ Positivity

Positivity of Hilbert space norm

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

■ Polynomiality

Lorentz covariance

■ Positivity

Positivity of Hilbert space norm

■ H^q has support $x \in [-1, +1]$.

Covariant
extension

■ Polynomiality

Lorentz covariance

■ Positivity

Positivity of Hilbert space norm

- H^q has support $x \in [-1, +1]$.

Relativistic quantum mechanics

Parameterizations

Definition

Polynomiality
Radon transform
Positivity
Inverse Radon
Examples

Computations

Design
Releases
Phenomenology

Fits

Status
Global CFF fit

Conclusion

Appendix

Covariant
extension

■ Polynomiality

Lorentz covariance

■ Positivity

Positivity of Hilbert space norm

■ H^q has support $x \in [-1, +1]$.

Relativistic quantum mechanics

■ Soft pion theorem (pion target)

$$H^q(x, \xi = 1, t = 0) = \frac{1}{2} \phi_\pi^q \left(\frac{1+x}{2} \right)$$

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

Covariant
extension

Parameterizations

Definition

Polynomiality
Radon transform
Positivity
Inverse Radon
Examples

Computations

Design
Releases
Phenomenology

Fits

Status
Global CFF fit

Conclusion

Appendix

■ Polynomiality

Lorentz covariance

■ Positivity

Positivity of Hilbert space norm

■ H^q has support $x \in [-1, +1]$.

Relativistic quantum mechanics

■ Soft pion theorem (pion target)

Dynamical chiral symmetry breaking

Covariant
extension

Parameterizations

Definition

- Polynomiality
- Radon transform
- Positivity
- Inverse Radon
- Examples

Computations

- Design
- Releases
- Phenomenology

Fits

- Status
- Global CFF fit

Conclusion

Appendix

■ Polynomiality

Lorentz covariance

■ Positivity

Positivity of Hilbert space norm

■ H^q has support $x \in [-1, +1]$.

Relativistic quantum mechanics

■ Soft pion theorem (pion target)

Dynamical chiral symmetry breaking

How can we implement *a priori* these theoretical constraints?

- In the following, focus on **polynomiality** and **positivity**.
- Do not discuss the reduction to form factors or PDFs.

Covariant
extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

- Write **polynomiality condition**:

$$\int_{-1}^1 dx x^m H^q(x, \xi, t) = \sum_{\substack{i=0 \\ \text{even}}}^m (2\xi)^i C_{mi}^q(t) + (2\xi)^{m+1} C_{mm+1}^q(t).$$

- Assume the existence of $D^q(z, t)$ such that:

$$\int_{-1}^{+1} dz z^m D(z, t) = C_{mm+1}^q(t).$$

- $H^q(x, \xi, t) - D(x/\xi, t)$ satisfies polynomiality at order m :

$$\int_{-1}^1 dx x^m (H^q(x, \xi, t) - D(x/\xi, t)) = \sum_{\substack{i=0 \\ \text{even}}}^m (2\xi)^i C_{mi}^q(t).$$

- **Ludwig-Helgason** condition: there exists F_D such that:

$$H(x, \xi, t) = D(x/\xi, t) + \int_{\Omega_{DD}} d\beta d\alpha F_D(\beta, \alpha, t) \delta(x - \beta - \alpha \xi).$$

Covariant
extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

- Most general representation of GPD:

$$H^q(x, \xi, t) = \int_{\Omega_{DD}} d\beta d\alpha \delta(x - \beta - \alpha\xi) (F^q(\beta, \alpha, t) + \xi G^q(\beta, \alpha, t))$$

- Support property: $x \in [-1, +1]$.
- Discrete symmetries: F^q is α -even and G^q is α -odd.
- **Gauge**: any representation (F^q, G^q) can be recast in one representation with a single DD f^q :

$$H^q(x, \xi, t) = x \int_{\Omega_{DD}} d\beta d\alpha f_{BMKS}^q(\beta, \alpha, t) \delta(x - \beta - \alpha\xi)$$

Belitsky *et al.*, Phys. Rev. **D64**, 116002 (2001)

$$H^q(x, \xi, t) = (1 - x) \int_{\Omega_{DD}} d\beta d\alpha f_P^q(\beta, \alpha, t) \delta(x - \beta - \alpha\xi)$$

Pobylitsa, Phys. Rev. **D67**, 034009 (2003)

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

For $s > 0$ and $\phi \in [0, 2\pi]$:


$$\mathcal{R}f(s, \phi) = \int_{-\infty}^{+\infty} d\beta d\alpha f(\beta, \alpha) \delta(s - \beta \cos \phi - \alpha \sin \phi)$$

and:

$$\mathcal{R}f(-s, \phi) = \mathcal{R}f(s, \phi \pm \pi)$$

Relation to GPDs:

$$x = \frac{s}{\cos \phi} \text{ and } \xi = \tan \phi$$



Relation between GPD and DD in Belitsky *et al.* gauge

$$\frac{\sqrt{1 + \xi^2}}{x} H(x, \xi) = \mathcal{R}f_{\text{BMKS}}(s, \phi),$$

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

For $s > 0$ and $\phi \in [0, 2\pi]$:


$$\mathcal{R}f(s, \phi) = \int_{-\infty}^{+\infty} d\beta d\alpha f(\beta, \alpha) \delta(s - \beta \cos \phi - \alpha \sin \phi)$$

and:

$$\mathcal{R}f(-s, \phi) = \mathcal{R}f(s, \phi \pm \pi)$$

Relation to GPDs:

$$x = \frac{s}{\cos \phi} \text{ and } \xi = \tan \phi$$



Relation between GPD and DD in Pobylitsa gauge

$$\frac{\sqrt{1 + \xi^2}}{1 - x} H(x, \xi) = \mathcal{R}f_P(s, \phi),$$

Covariant
extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

- The Mellin moments of a Radon transform are **homogeneous polynomials** in $\omega = (\sin \phi, \cos \phi)$.
- The converse is also true:

Theorem (Hertle, 1983)

Let $g(s, \omega)$ an even compactly-supported distribution. Then g is itself the Radon transform of a compactly-supported distribution if and only if the **Ludwig-Helgason consistency condition** hold:

- g is C^∞ in ω ,
- $\int ds s^m g(s, \omega)$ is a homogeneous polynomial of degree m for all integer $m \geq 0$.

- Double Distributions and the Radon transform are the **natural solution** of the polynomiality condition.

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit


Conclusion

Appendix

DGLAP and ERBL regions

$$(x, \xi) \in \text{DGLAP} \Leftrightarrow |s| \geq |\sin \phi| ,$$

$$(x, \xi) \in \text{ERBL} \Leftrightarrow |s| \leq |\sin \phi| .$$



Each point (β, α) with $\beta \neq 0$ contributes to **both** DGLAP and ERBL regions.

Covariant
extension

Parameterizations

- Definition
- Polynomiality
- Radon transform
- Positivity
- Inverse Radon
- Examples

Computations

- Design
- Releases
- Phenomenology

Fits

- Status
- Global CFF fit

Conclusion

Appendix

Theorem (simple case)

Let f be a compactly-supported summable function defined on \mathbb{R}^2 and $\mathcal{R}f$ its Radon transform.

Let $(s_0, \omega_0) \in \mathbb{R} \times S^1$ and U_0 an open neighborhood of ω_0 s.t.:

for all $s > s_0$ and $\omega \in U_0$ $\mathcal{R}f(s, \omega) = 0$.

Then $f(\mathbf{x}) = 0$ on the half-plane $\langle \mathbf{x} | \omega_0 \rangle > s_0$ of \mathbb{R}^2 .

Theorem (Boman and Todd Quinto, 1987)

Assume $(s_0, \omega_0) \in \mathbb{R} \times S^{n-1}$ and $f \in \mathcal{E}'(\mathbb{R})$. Let $\mu(\mathbf{x}, \omega)$ be a strictly positive real analytic function on $\mathbb{R}^n \times S^{n-1}$ that is even in ω . Let U_0 be an open neighborhood of ω_0 . Finally assume $R_\mu(s, \omega) = 0$ for $s > s_0$ and $\omega \in U_0$. Then $f = 0$ on the half space $\langle \mathbf{x} | \omega_0 \rangle > s_0$.

Covariant
extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

Theorem (simple case)

for all $s > s_0$ and $\omega \in U_0$ $\mathcal{R}f(s, \omega) = 0$.

Then $f(\mathbf{x}) = 0$ on the half-plane $\langle \mathbf{x} | \omega_0 \rangle > s_0$ of \mathbb{R}^2 .

Consider a GPD H being zero on the DGLAP region.

- Take $\xi_0 = \tan \phi_0 \in [0, 1]$, $x_0 \in]\xi_0, +\infty[$ and s_0 s.t.
 $x_0 \cos \phi_0 > s_0 > \sin \phi_0$.
- $\exists \epsilon > 0$ s.t. $s_0 > \sin \phi$ for $|\phi - \phi_0| < \epsilon$.
- Hyp: the underlying DD f has a zero Radon transform for all $\phi \in]\phi_0 - \epsilon, \phi_0 + \epsilon[$ and $s > s_0$ (DGLAP region).
- Then $f(\beta, \alpha) = 0$ for all (β, α) s.t.
 $\beta \cos \phi_0 + \alpha \sin \phi_0 = s > s_0$.
- At last select $s = x_0 \cos \phi_0$ to get $\beta + \alpha \xi_0 = x_0$.
- Cannot constrain the line $\beta = 0$. \square

Proof.

Covariant
extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

- Identify the matrix element defining a GPD as an **inner product** of two different states.
- Apply Cauchy-Schwartz inequality, and identify PDFs at specific kinematic points, e.g.:

$$|H^q(x, \xi, t)| \leq \sqrt{\frac{1}{1-\xi^2} q\left(\frac{x+\xi}{1+\xi}\right) q\left(\frac{x-\xi}{1-\xi}\right)}$$

- This procedure yields **infinitely many inequalities** stable under LO evolution.

Pobylitsa, Phys. Rev. D66, 094002 (2002)

- The **overlap representation** guarantees *a priori* the fulfillment of positivity constraints.

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

- Decompose an hadronic state $|H; P, \lambda\rangle$ in a Fock basis:

$$|H; P, \lambda\rangle = \sum_{N, \beta} \int [dx d\mathbf{k}_\perp]_N \psi_N^{(\beta, \lambda)}(x_1, \mathbf{k}_{\perp 1}, \dots, x_N, \mathbf{k}_{\perp N}) |\beta, k_1, \dots, k_N\rangle$$

- Derive an expression for the pion GPD in the DGLAP region $\xi \leq x \leq 1$:

$$H^q(x, \xi, t) \propto \sum_{\beta, j} \int [d\bar{x} d\bar{\mathbf{k}}_\perp]_N \delta_{j,q} \delta(x - \bar{x}_j) (\psi_N^{(\beta, \lambda)})^*(\bar{x}', \hat{\mathbf{k}}'_\perp) \psi_N^{(\beta, \lambda)}(\tilde{x}, \tilde{\mathbf{k}}_\perp)$$

with $\tilde{x}, \tilde{\mathbf{k}}_\perp$ (resp. $\hat{x}', \hat{\mathbf{k}}'_\perp$) generically denoting incoming (resp. outgoing) parton kinematics.

Diehl *et al.*, Nucl. Phys. **B596**, 33 (2001)

- Similar expression in the ERBL region $-\xi \leq x \leq \xi$, but with overlap of N - and $(N+2)$ -body LFWFs.

Covariant extension

Parameterizations

Definition
Polynomiality
Radon transform
Positivity
Inverse Radon
Examples

Computations

Design
Releases
Phenomenology

Fits

Status
Global CFF fit

Conclusion

Appendix

For **any model of LFWF**, one has to address the following three questions:

- 1 Does the extension exist?
- 2 If it exists, is it unique?
- 3 How can we compute this extension?

Modeling strategy

- 1 Ensure positivity by modeling the DGLAP region as an overlap of LFWFs.
- 2 Ensure polynomiality by inverting the Radon transform to identify an underlying DD.

Chouika *et al.*, Eur. Phys. J. **C77**, 906 (2017)

Covariant
extension

Parameterizations

Definition
Polynomiality
Radon transform
Positivity
Inverse Radon

Examples

Computations

Design
Releases
Phenomenology


Fits

Status
Global CFF fit

Conclusion

Appendix

- Numerical evaluation ***almost unavoidable*** (polar vs cartesian coordinates).
- Ill-posedness by **lack of continuity**.
- The **unlimited** Radon inverse problem is **mildly** ill-posed while the **limited** one is **severely** ill-posed.
- Even if it existed, an analytic expression of the invert Radon transform would be of **limited practical use**.



Mezrag, PhD
dissertation
(2015)

Covariant
extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

Fully discrete case

Assume f piecewise-constant with values f_m for $1 \leq m \leq M$.

For a collection of lines $(L_n)_{1 \leq n \leq N}$ crossing Ω_{DD} , the Radon transform writes:

$$g_n = \mathcal{R}f = \int_{L_n} f = \sum_{m=1}^M f_m \times \text{Measure}(L_n \cap C_m) \quad \text{for } 1 \leq n \leq N$$

A discretized problem

Consider $N+1$ Hilbert spaces H, H_1, \dots, H_N , and a family of continuous surjective operators $R_n : H \rightarrow H_n$ for $1 \leq n \leq N$. Being given $g_1 \in H_1, \dots, g_n \in H_n$, we search f solving the following system of equations:

$$R_n f = g_n \quad \text{for } 1 \leq n \leq N$$

Covariant extension

Parameterizations

Definition
Polynomiality
Radon transform
Positivity
Inverse Radon
Examples

Computations

Design
Releases
Phenomenology

Fits

Status
Global CFF fit

Conclusion

Appendix

Kaczmarz algorithm

Denote P_n the orthogonal projection on the *affine* subspace $R_n f = g_n$. Starting from $f^0 \in H$, the sequence defined iteratively by:

$$f^{k+1} = P_N P_{N-1} \dots P_1 f^k$$

converges to the solution of the system.

The convergence is exponential if the projections are randomly ordered.

Strohmer and Vershynin, Jour. Four. Analysis and Appl. 15,
437 (2009)

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology


Fits

Status

Global CFF fit

Conclusion

Appendix



Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

And if the input data are inconsistent?

- Instead of solving $g = \mathcal{R}f$, find f such that $\|g - \mathcal{R}f\|_2$ is **minimum**.
- The solution **always exists**.
- The input data are **inconsistent** if $\|g - \mathcal{R}f\|_2 > 0$.

Covariant extension

How can we get a DD from a GPD in the DGLAP region?

Parameterizations

- Definition
- Polynomiality
- Radon transform
- Positivity
- Inverse Radon**
- Examples

Computations



- Design
- Releases
- Phenomenology

Fits

- Status
- Global CFF fit

Conclusion

Appendix



Covariant
extension

How can we get a DD from a GPD in the DGLAP region?

- Restrict to quark GPDs ($\beta > 0$).

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology



Fits

Status

Global CFF fit

Conclusion

Appendix



Covariant
extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status



Global CFF fit

Conclusion

Appendix

How can we get a DD from a GPD in the DGLAP region?

- Restrict to quark GPDs ($\beta > 0$).
- Only ERBL region "sees" both $\beta > 0$ and $\beta < 0$.



Covariant
extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status



Global CFF fit

Conclusion

Appendix

How can we get a DD from a GPD in the DGLAP region?

- Restrict to quark GPDs ($\beta > 0$).
- Only ERBL region "sees" both $\beta > 0$ and $\beta < 0$.
- Use α -parity of the DD.



Covariant
extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits


Status

Global CFF fit

Conclusion

Appendix

Example of a P1 basis function



- Discretize the DD on a mesh with $n \simeq 800$ triangular cells.
- Compute the Radon transform of a P1 basis function.
- Sample $m \simeq 4n$ (x, ξ) -lines intersecting the DD support.
- Solve a linear system $AX = B$ with A a sparse $m \times n$ matrix.
- Adopt an iterative regularization method: LSMR.

Fong and Saunders, arXiv:1006.0758

Examples - benchmarks (1/4).

Algebraic Bethe-Salpeter model.

Covariant
extension

$$\Psi_{I=0}(x, \mathbf{k}_\perp) = 8\sqrt{15}\pi \frac{M^3}{(\mathbf{k}_\perp^2 + M^2)^2} (1-x)x,$$

Parameterizations

Definition

Polynomiality


Radon transform

Positivity


Inverse Radon

Examples

$H(x, \xi, t = 0)$



$H(x, \xi = 0.5, t)$



Examples - benchmarks (2/4).

Algebraic spectator model.

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion


Appendix

$$\varphi(x, \mathbf{k}_\perp) = \frac{gM^{2p}}{\sqrt{1-x}} x^{-p} \left(M^2 - \frac{\mathbf{k}_\perp^2 + m^2}{x} - \frac{\mathbf{k}_\perp^2 + \lambda^2}{1-x} \right)^{-p-1}$$

Hwang and Müller, Phys. Lett. **B660**, 350 (2008)


$H(x, \xi, t = 0)$

numerical result
analytical result



$H(x, \xi = 0.5, t)$

numerical result
analytical result



Chouika et al., Eur. Phys. J. **C77**, 906 (2017)

Covariant
extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status


Global CFF fit

Conclusion

Appendix

$$q\text{Regge}(x) = \frac{35}{32} \frac{(1-x)^3}{\sqrt{x}} .$$

$H(x, \xi, t = 0)$



Chouika *et al.*, Eur. Phys. J. **C77**, 906 (2017)

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Basis



8

Fig. 1.

End

Status

Appendix



Chouika *et al.*, Eur. Phys. J. **C77**, 906 (2017)

Computing exclusive processes with increasing precision?



PARtonic
Tomography
Of
Nucleon
Software

Covariant
extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

Experimental
data and
phenomenology

Full processes

Computation
of amplitudes

Small distance
contributions

First
principles and
fundamental
parameters

Large distance
contributions

Covariant extension

Parameterizations
Definition
Polynomiality
Radon transform
Positivity
Inverse Radon
Examples

Computations
Design
Releases
Phenomenology

Fits
Status
Global CFF fit

Conclusion
Appendix

Experimental data and phenomenology


Computation of amplitudes

First principles and fundamental parameters

Full processes

Small distance contributions

Large distance contributions



Computing chain design.

Differential studies: physical models and numerical methods.

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit


Conclusion

Appendix

Experimental data and phenomenology

Computation of amplitudes

First principles and fundamental parameters



Computing chain design.

Differential studies: physical models and numerical methods.

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion


Appendix

Experimental data and phenomenology

Need for modularity

Computation of amplitudes

First principles and fundamental parameters



- Many observables.
- Kinematic reach.

- Perturbative approximations.
- Physical models.
- Fits.
- Numerical methods.
- Accuracy and speed.

Computing chain design.

Differential studies: physical models and numerical methods.

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion


Appendix

Experimental data and phenomenology

Need for modularity

Computation of amplitudes

First principles and fundamental parameters



- Many observables.
- Kinematic reach.

- Perturbative approximations.
- Physical models.**
- Fits.
- Numerical methods.
- Accuracy and speed.

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion


Appendix

Experimental data and phenomenology

Need for modularity

Computation of amplitudes

First principles and fundamental parameters



- Many observables.
- Kinematic reach.

- Perturbative approximations.
- Physical models.
- Fits.
- Numerical methods.
- Accuracy and speed.

Computing chain design.

Differential studies: physical models and numerical methods.

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion


Appendix

Experimental data and phenomenology

Need for modularity

Computation of amplitudes

First principles and fundamental parameters



- Many observables.
- Kinematic reach.

- Perturbative approximations.
- Physical models.
- Fits.
- Numerical methods.
- Accuracy and speed.

Computing chain design.

Differential studies: physical models and numerical methods.

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion


Appendix

Experimental data and phenomenology

Need for modularity

Computation of amplitudes

First principles and fundamental parameters



- Many observables.
- Kinematic reach.

- Perturbative approximations.
- Physical models.
- Fits.
- Numerical methods.
- Accuracy and speed.

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology


Fits

Status

Global CFF fit

Conclusion

Appendix



The screenshot shows the main page of the PARTONS website. The header includes the CEA Paris-Saclay logo, the title "PARTONS PARtonic Tomography Of Nucleon Software", and a search bar. The navigation menu has links for Main Page, Download, Tutorials +, Reference documentation +, and About +. The main content area features a large "PARTONS" logo, a section titled "What is PARTONS?", and a "Table of Contents" sidebar. The "What is PARTONS?" section describes the software as a C++ framework for GPDs and provides details about its structure and applications. The "Table of Contents" sidebar lists various sections including "What is PARTONS?", "Get PARTONS", "Configure PARTONS", "How to use PARTONS", "Publications and talks", "License", and "Contact and newsletter". Below the main content, there is a "Get PARTONS" section and a reference to a 2018 publication.

PARTONS

PARtonic Tomography Of Nucleon Software

Main Page

What is PARTONS?

PARTONS is a C++ software framework dedicated to the phenomenology of Generalized Parton Distributions (GPDs). GPDs provide a comprehensive description of the partonic structure of the nucleon and contain a wealth of new information. In particular, GPDs provide a description of the nucleon as an extended object, referred to as 3-dimensional nucleon tomography, and give an access to the orbital angular momentum of quarks.

PARTONS provides a necessary bridge between models of GPDs and experimental data measured in various exclusive channels, like Deeply Virtual Compton Scattering (DVCS) and Hard Exclusive Meson Production (HEMP). The experimental programme devoted to study GPDs has been carrying out by several experiments, like HERMES at DESY (closed), COMPASS at CERN, Hall-A and CLAS at JLab. GPD subject will be also a key component of the physics case for the expected Electron Ion Collider (EIC).

PARTONS is useful to theorists to develop new models, phenomenologists to interpret existing measurements and to experimentalists to design new experiments.

Get PARTONS

Berthou *et al.*, Eur. Phys. J. C78, 478 (2018)

Covariant extension

Parameterizations

- Definition
- Polynomiality
- Radon transform
- Positivity
- Inverse Radon
- Examples

Computations


- Design
- Releases**
- Phenomenology

Fits

- Status
- Global CFF fit

Conclusion

Appendix



partons > core > Details

core

Filter by name... Last created

Project	Description	Rating	Actions
partons	PARTONS project	★ 2	...
elementary-utils	Utility softwares (logger, parser, threads, string and file manipulation)	★ 0	...
numa	NumA++: numerical analysis C++ routines	★ 0	...
partons-example	Running version of PARTONS with examples (C++ code and XML computing scenarios)	★ 0	...

Prev 1 Next

Berthou *et al.*, Eur. Phys. J. C78, 478 (2018)

Covariant extension

Parameterizations

- Definition
- Polynomiality
- Radon transform
- Positivity
- Inverse Radon
- Examples

Computations

- Design

Releases


- Phenomenology

Fits

- Status
- Global CFF fit

Conclusion

Appendix



The screenshot shows the PARTONS website (partons.cea.fr) with a dark blue header bar containing navigation links: Main Page, Download, Tutorials +, Reference documentation +, and About +. Below the header is a search bar. The main content area features several sections: Contact (with email partons@cea.fr), Technical support (with email partons@cea.fr), Newsletter (instructions for subscribing via email to sympa@saxifrage.saclay.cea.fr), Development team (mentioning the GitLab page), and a footer note about joining the development team. The website is styled with a clean, modern look, using white text on a dark background for the header and light gray text on white for the main content.

Generated on Wed Mar 21 2018 16:23:16 for PARTONS 1.0 by Doxygen 1.8.14

Berthou *et al.*, Eur. Phys. J. C78, 478 (2018)

Covariant extension

Parameterizations

Definition
Polynomiality
Radon transform
Positivity
Inverse Radon
Examples

Computations

Design
Releases



Phenomenology

Fits

Status
Global CFF fit

Conclusion

Appendix



- Only LO phenomenology achievable without extension to ERBL region.
- Computation of **various DVCS observables** in the valence region under **different pQCD assumptions** with PARTONS.

Chouika, PhD thesis (2018)

Fitting experimental data?

Covariant extension

Parameterizations

- Definition
- Polynomiality
- Radon transform
- Positivity
- Inverse Radon
- Examples

Computations

- Design
- Releases
- Phenomenology

Fits

- Status
- Global CFF fit

Conclusion

Appendix

- GPD fits **only in the small x_B region** with a **flexible** parameterization (kinematic simplifications).
- Global fits of CFFs in the sea and valence regions.
- Some GPD models with non-flexible parameterizations adjusted to experimental DVCS or DVMP data.

Kumerički *et al.*, Eur. Phys. J. **A52**, 157 (2016)

The situation can be improved!

- GPD parameterizations satisfying *a priori* all theoretical constraints on GPDs.
- Computing framework to go beyond leading order and leading twist analysis.

Selected DVCS measurements.

All existing sets except $d^4\sigma_{UU}^-$ from Hall A (2015-17) and HERA.

	No.	Collab.	Year	Ref.	Observable	Kinematic dependence	No. of points used / all
Covariant extension	1	HERMES	2001	[13]	A_{LU}^+	ϕ	10 / 10
	2		2006	[114]	$A_C^{\cos i\phi}$	$i = 1$	4 / 4
	3		2008	[115]	$A_C^{\cos i\phi}$	$i = 0, 1$	x_{Bj} 18 / 24
Parameterizations					$A_{UT,DVCS}^{\sin(\phi-\phi_S)\cos i\phi}$	$i = 0$	
Definition					$A_{UT,I}^{\sin(\phi-\phi_S)\cos i\phi}$	$i = 0, 1$	
Polynomiality					$A_{UT,I}^{\cos(\phi-\phi_S)\sin i\phi}$	$i = 1$	
Radon transform	4		2009	[116]	$A_{LU,I}^{\sin i\phi}$	$i = 1, 2$	x_{Bj} 35 / 42
Positivity					$A_{LU,DVCS}^{\sin i\phi}$	$i = 1$	
Inverse Radon	5		2010	[117]	$A_{UL}^{+, \sin i\phi}$	$i = 0, 1, 2, 3$	
Examples					$A_{UL}^{+, \cos i\phi}$	$i = 0, 1, 2$	
Computations	6		2011	[118]	$A_{LT,DVCS}^{\cos(\phi-\phi_S)\cos i\phi}$	$i = 0, 1$	x_{Bj} 24 / 32
Design					$A_{LT,DVCS}^{\sin(\phi-\phi_S)\sin i\phi}$	$i = 1$	
Releases					$A_{LT,I}^{\cos(\phi-\phi_S)\cos i\phi}$	$i = 0, 1, 2$	
Phenomenology	7		2012	[119]	$A_{LT,I}^{\sin(\phi-\phi_S)\sin i\phi}$	$i = 1, 2$	
Fits					$A_{LU,I}^{\sin i\phi}$	$i = 1, 2$	x_{Bj} 35 / 42
Status	8	CLAS	2001	[14]	$A_{LU}^{-, \sin i\phi}$	$i = 1, 2$	— 0 / 2
Global CFF fit	9		2006	[120]	$A_{UL}^{-, \sin i\phi}$	$i = 1, 2$	— 2 / 2
Conclusion	10		2008	[121]	A_{LU}^-	ϕ	283 / 737
Appendix	11		2009	[122]	A_{LU}^-	ϕ	22 / 33
	12		2015	[123]	$A_{LU}^-, A_{UL}^-, A_{LL}^-$	ϕ	311 / 497
	13		2015	[124]	$d^4\sigma_{UU}^-$	ϕ	1333 / 1933
	14	Hall A	2015	[112]	$\Delta d^4\sigma_{LU}^-$	ϕ	228 / 228
	15		2017	[113]	$\Delta d^4\sigma_{LU}^-$	ϕ	276 / 358
	16	COMPASS	2018	[55]	b	—	1 / 1
					SUM:	2600 / 3970	

Moutarde et al., arXiv:1807.07620

Covariant extension

Parameterizations

- Definition
- Polynomiality
- Radon transform
- Positivity
- Inverse Radon
- Examples

Computations





- Design
- Releases
- Phenomenology

Fits

- Status
- Global CFF fit**

Conclusion

Appendix



Conclusion

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

- We can now build generic GPD models satisfying *a priori* **all theoretical constraints**.
- We now have tools to **systematically relate** these models to **experimental data**. Open source release under GPLv3.0. of the PARTONS framework.
- We now have an **operating fitting engine** for global CFF fits.

New studies become possible!

- Global GPD fits.
- Energy-momentum structure of hadrons.
- Quantitative impact of nonperturbative QCD ingredients on 3D hadron structure studies.
- ???

Appendix

Covariant extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

Proof (elements)

Let $\mathbf{x}_0 = (\mathbf{b}, \mathbf{a}) \in \mathbb{R}^2$, $s \in \mathbb{R}$ and $\delta > 0$ such that

$$\langle \mathbf{x}_0 | \omega_0 \rangle = \mathbf{b} \cos \phi_0 + \mathbf{a} \sin \phi_0 = s > s_0 + \delta.$$

Denote \mathcal{B} a ball containing \mathbf{x}_0 and the support of f , which is bounded by assumption.

We will show that $f = 0$ in a neighborhood of \mathbf{x}_0 in \mathcal{B} .

Covariant extension

Parameterizations

- Definition
- Polynomiality
- Radon transform
- Positivity
- Inverse Radon
- Examples

Computations

- Design
- Releases
- Phenomenology

Fits

- Status
- Global CFF fit


Conclusion

Appendix

Step 1

Identification of a neighborhood T of \mathbb{N}_0 s.t.:

$$\forall s > s_0 + \delta, \quad \forall \omega \in T, \quad \int_{\mathbb{R}^2} d\mathbb{N} \delta(s - \langle \mathbb{N} | \omega \rangle) f(\mathbb{N}) = 0.$$



Covariant extension

Parameterizations

- Definition
- Polynomiality
- Radon transform
- Positivity
- Inverse Radon
- Examples

Computations

- Design
- Releases
- Phenomenology

Fits

- Status
- Global CFF fit


Conclusion

Appendix

Step 2

Prove by induction on the multi index $\mathbf{m} = (m, n)$ that, for all nonnegative integers m, n and $\omega \in T$:

$$\int_{[s_0 + \delta, +\infty[} ds \int_{\mathcal{B}} d\mathbb{N} \delta(s - \langle \mathbb{N} | \omega \rangle) \mathbb{N}^{\mathbf{m}} \langle \mathbb{N} | \omega \rangle^k f(\mathbb{N}) = 0 .$$



**Covariant
extension****Parameterizations**

- Definition
- Polynomiality
- Radon transform
- Positivity
- Inverse Radon
- Examples

Computations


- Design
- Releases
- Phenomenology

Fits

- Status
- Global CFF fit

Conclusion**Appendix****Step 2 (cont')**

Induction step: infinitesimal change of the first cartesian coordinate of ω . G-type contributions go to 0.



Covariant
extension

Parameterizations

Definition

Polynomiality

Radon transform

Positivity

Inverse Radon

Examples

Computations

Design

Releases

Phenomenology

Fits

Status

Global CFF fit

Conclusion

Appendix

Step 3

Apply previous result with $k = 0$ and $\omega = \omega_0$, let δ goes to 0:

$$\text{for all } n, m \geq 0 \quad \int_{H_{\omega_0}} d\beta d\alpha \beta^m \alpha^n f(\beta, \alpha) = 0 .$$

Conclude by injectivity of the Fourier transform from $L^1(\mathbb{R}^2)$ into the set of continuous functions on \mathbb{R}^2 .

[◀ Back to uniqueness statement.](#)

Commissariat à l'énergie atomique et aux énergies alternatives
Centre de Saclay | 91191 Gif-sur-Yvette Cedex
T. +33(0)1 69 08 73 88 | F. +33(0)1 69 08 75 84

DRF
Irfu
DPIN

Établissement public à caractère industriel et commercial | R.C.S. Paris B 775 685 019