


# Neutrino Transport in Holography



Utrecht  
University

Edwan PREAU



26/03/25

**Collaborators:** Elias KIRITSIS (APC), Francesco NITTI (APC) and Matti JÄRVINEN (APCTP)


[arXiv:2306.00192](https://arxiv.org/abs/2306.00192), [arXiv:2409.04630](https://arxiv.org/abs/2409.04630)

# Outline

- 1) Motivation
- 2) Introduction : Formalism for neutrino transport
- 3) Holographic Set-up
- 4) Holographic calculation of the chiral current correlators
- 5) *Towards isospin asymmetry*
- 6) Summary

# Motivation


- **Neutrino ( $\nu$ )** radiation is the main mechanism for **Neutron Star (NS) cooling**
- Requires the knowledge of  $\nu$  interaction with **dense QCD matter** in the core
- **Simulations** need an **input** from particle physics :  $\mathbf{j} \ \& \ \alpha \leftrightarrow \langle J_{L/R} J_{L/R} \rangle^R$



# Motivation

- Computing  $\langle J_{L/R} J_{L/R} \rangle^R$  in the **dense strongly-coupled** QCD matter is a difficult problem
- We consider the **holographic** approach

**Problem :** compute  $\langle J_{L/R} J_{L/R} \rangle^R$  in **holographic QCD** at finite  $T$  and  $n_B$   
→ This work : **simplest** toy model (quark matter in  $\mathcal{N} = 4$  SYM)



# Formalism for neutrino transport

**Exercise :** compute the **(exact) propagator**  $G_\nu(\vec{x}_1, t_1; \vec{x}_2, t_2)$  of  $\nu$ 's in a **dense QCD medium**

Quasi-particle approximation :

$G_\nu$  is described by the  **$\nu$  distribution function**  $f_\nu(\vec{x}, t; k_\nu)$

The transport of neutrinos is described by the **Boltzmann equation** obeyed by  $f_\nu$

$$(k_\nu \cdot \partial) f_\nu \equiv \underbrace{j(E_\nu)}_{\text{Emissivity}} (1 - f_\nu) - \underbrace{\alpha(E_\nu)}_{\text{Absorptivity}} f_\nu \equiv j(E_\nu) - \underbrace{\kappa(E_\nu)}_{\text{Opacity}} f_\nu$$

Emissivity

Absorptivity


Opacity

$$\kappa = j + \alpha$$

# Schwinger-Dyson equation

The kinetic equation can be derived from the finite temperature **Schwinger-Dyson equation**

The self-energy  $\Sigma$  is expanded at order  $\mathcal{O}(G_F^2)$  in the weak interaction




It is fully **non-perturbative** in the **strong** interaction

# Schwinger-Dyson equation

The kinetic equation can be derived from the finite temperature **Schwinger-Dyson equation**

$$\nu + n \leftrightarrow e^- + p$$



Dirac equation

$$j(E_\nu) = G_F^2 \int \frac{d\vec{k}_e^3}{(2\pi)^3} \underbrace{(\text{kins})^{\lambda\sigma}}_{\vec{k}_e, \vec{k}_\nu} \times \underbrace{(\text{stats})}_{f_e, f_W} \times \text{Im}(i\langle J_\lambda^- J_\sigma^+ \rangle^R),$$

Dense QCD  
 $\sim \langle J_\lambda^L J_\sigma^L \rangle^R$

# The holographic set-up




# The Holographic Set-up

**Simplest** bottom-up holographic toy model with **chiral currents**  $J_{L/R}^\mu$

$$\begin{array}{ccc}
 T_{\mu\nu} & \leftrightarrow & g_{MN} \\
 U(N_f)_L \times U(N_f)_R : \partial_\mu J_{L/R}^\mu = 0 & \leftrightarrow & U(N_f)_L \times U(N_f)_R : A_{L/R}^M
 \end{array}$$

$N_c \rightarrow \infty, \frac{N_f}{N_c} \text{ finite}$




$$S = M_{Pl}^3 N_c^2 \int dx^5 \sqrt{-g} \left( R + \frac{12}{\ell^2} - \frac{\kappa}{N_c} \text{Tr} \left\{ F_{MN}^{(L)} F_{(L)}^{MN} + F_{MN}^{(R)} F_{(R)}^{MN} \right\} \right),$$

# Background solution

We want to compute  $\langle J_\lambda^- J_\sigma^+ \rangle^R$  in an equilibrium state at **finite**  $(T, \mu_q)$  = dense strongly-coupled **quark matter**

→ Charged AdS **black hole**, with charge  $Q \propto \mu_q$




# Summary of parameters

Parameters of the <b>model</b>	$M_{Pl} \ell$	Fitted to lattice quark-gluon thermodynamics
	$\kappa$	
<b>Environmental</b> parameters	$\frac{\mu_q}{T}$	Varied
<b>Neutrino</b> properties	$\frac{E_\nu}{T}$	Varied

# Holographic calculation of the chiral current 2-point function

# Perturbations of AdS-RN



The boundary plasma has an **SO(3) rotational invariance**

$$\langle J_\lambda J_\sigma \rangle^R(\omega, \vec{k}) = P_{\lambda\sigma}^\perp(\omega, \vec{k}) i\Pi^\perp(\omega, \vec{k}) + P_{\lambda\sigma}^\parallel(\omega, \vec{k}) i\Pi^\parallel(\omega, \vec{k})$$

# Hydrodynamic approximation

The **long-range** behavior of a system **near equilibrium** is described by **hydrodynamics**

→ Equilibrium **correlators** follow a **universal** long-range structure :

- **Expansion** in  $(\omega/T, k/T)$ , with **transport coefficients**
- The **hydro modes** appear as **poles** at leading order

$$\langle J_\lambda J_\sigma \rangle^R(\omega, \vec{k}) = \underbrace{\sigma}_{\text{Conductivity}} \left( P_{\lambda\sigma}^\perp \omega + P_{\lambda\sigma}^\parallel \frac{\omega^2 - k^2}{\omega + iDk^2} \right) \left( 1 + \mathcal{O}\left(\frac{\omega}{T}, \frac{k^2}{T^2}\right) \right),$$

$\partial_t J^0 = D\Delta J^0$

# Hydrodynamic approximation at $\mu_q \gg T$

$$\langle J_\lambda J_\sigma \rangle^R(\omega, \vec{k}) = \sigma \left( P_{\lambda\sigma}^\perp \omega + P_{\lambda\sigma}^\parallel \frac{\omega^2 - k^2}{\omega + i\mathbf{D}k^2} \right) \left( 1 + \mathcal{O}\left(\frac{\omega}{T}, \frac{k^2}{T^2}\right) \right),$$

Hydro a priori **breaks down at  $\omega, k \gg T$**

**AdS-RN** : the LO **hydro** approximation remains valid as long as  **$\omega, k \ll \mu_q$**


→  **$\nu$  transport** in a NS:  **$E_\nu, \mu_e, \mu_\nu \ll \mu_q$**

[Davison & Parnachev '13]  
[Moitra, Sake & Trivedi '21]

At  **$\mu_q \gg T$** , we have  $\mu_e, \mu_\nu \simeq 0.7 \mu_q$

# Hydrodynamic approximation at $\mu_q \gg T$

- The scale where **hydrodynamics breaks down** is identified by analyzing the **poles** of the correlator  $\leftrightarrow$  **QNM's** of AdS-RN





[Edalati, Jottar & Leigh '10]



# Hydrodynamic approximation at $\mu_q \gg T$

- When **k is increased**, the poles **collide**, but a **diffusive pole** effectively remains
- For **hydrodynamics** to remain valid :  $Res(\omega_n) \rightarrow 0$  as  $T/\mu \rightarrow 0$


[Gursoy, Järvinen, Policastro & Zinnato '21]



From [Arean, Davison, Goutéraux & Suzuki '21]

# Numerical results


# Charged current correlators




$$\frac{\mu_q}{T} \simeq 65$$

# Opacities : comparison with hydro

$T = 10 \text{ MeV}$



$\nu$




$\bar{\nu}$

$$\kappa(E_\nu) = j(E_\nu) + \frac{1}{\lambda(E_\nu)}$$

# Comparison with weak coupling

[Iwamoto '82]




$$T = 10 \text{ MeV}, \quad n_B = 0.11 \text{ fm}^{-3}$$

Towards isospin asymmetry


# Phase diagram at finite $\mu_3$

- $\mu_3$  is introduced as a **source** for  $L_0^3 + R_0^3$
- **AdS RN** is still a solution of the Einstein-Yang-Mills equations (with  $Q_3 \propto \mu_3$ ), but other solutions exist : **p-wave superconductors (SC)**, with  **$U(1)_3$  spontaneously broken** [Gubser '08], [Gubser, Pufu, '08]



- Gubser and Pufu considered specific cases :
  - **General ansatz** but **probe** ( $\kappa \rightarrow 0$ )
  - **Back-reacted** but for a **specific ansatz** ( $\kappa$  finite)
- First step for our purpose: derive the **full 3-dimensional phase diagram** ( $\kappa, \mu_3/T, \mu_q/T$ )

# Phase diagram at finite $\mu_3$



Partial results at  $\mu_q = 0$  in  
[Ammon, Erdmenger, Grass, Kerner  
& O'Bannon '10]  
[Arias & Landea '13]



# General ansatz for SC solutions

- At  $T \neq 0$  and  $\mu_3 \neq 0$ , the theory has  $SO(3) \times U(1)_3$  symmetry ( $d = 3 + 1$ )


→ The **most general SC ansatz** for the gauge fields is

$$L = R = \frac{1}{2} \Phi(r) dt \mathbb{I}_2 + \frac{1}{2} \Phi_3(r) dt \sigma^3 + \frac{1}{2} A_z^1(r) dz \sigma^1 + \frac{1}{2} A_x^2(r) dx \sigma^2,$$

- We expect a **discrete set of solutions**, labeled by the **number of nodes**  $(n, m)$
- The **dominant solutions** should be the **nodeless** solutions  $(0,1)$  and  $(1,1)$ :
  - $(0,1) : A_x^2(r) = 0$ , **p-wave** [Gubser, Pufu, '08]
  - $(1,1) : A_z^1(r) = A_x^2(r) \neq 0$ , **(p+ip)-wave** [Gubser '08]
- Both **preserve**  $U(1) \subset SO(3) \times U(1)$ , so the **metric ansatz** can be chosen as

$$ds^2 = e^{2A(r)} (-f(r) dt^2 + f(r)^{-1} dr^2 + dx^2 + dy^2 + h(r) dz^2)$$

# The plane $\mu_q = 0$




## Solitonic solution

- Horizonless
- $AdS^5$  IR geometry

~ [Horowitz & Roberts, '09]


A plane at fixed  $\kappa < \kappa_c$  with  $\mu_q \neq 0$



**Extremal hairy BH**

- $AdS^2$  IR geometry
- $Q_q$  carried by the horizon
- $Q_3$  carried by the hair

# The full 3d phase diagram



# Summary and outlook

First step towards the description of holographic **neutrino transport** : toy model of **strongly-coupled quark matter**

- **Hydrodynamic** behavior
- **Opacity suppressed** compared with the weak coupling result
- More work is needed to **corroborate** these results

Several directions of improvement :


- Transport in an **isospin asymmetric** medium
- **Neutrino rates** from **neutral current** interactions
- **More realistic model** of holographic QCD

# Thank you !

arXiv:2306.00192 [astro-ph.HE]  
arXiv:2409.04630 [hep-th]

# Appendix

# Phases of QCD





## H. Details about the perturbations of AdS-RN

# Perturbations of AdS-RN

[Son & Starinets '02]

[Skenderis & van Rees '08]

$\langle J_\lambda J_\sigma \rangle^R$  is obtained by considering **perturbations** of the fields on top of **AdS-RN**

$$A_{L/R}^M \rightarrow \bar{A}_{L/R}^M + \delta A_{L/R}^M, \quad g_{MN} \rightarrow \bar{g}_{MN} + \delta g_{MN},$$

$$\forall \varphi, \delta \varphi = \int \frac{d^4 k}{(2\pi)^4} e^{ik \cdot x} C_k(z) \delta \varphi_0(k), \quad \text{At } z \sim z_H : C_k(z) \sim (z_H - z)^{-\frac{ik^0 z_H}{4}}$$

Infalling boundary condition

- Only  $\delta T_{MN} \propto \delta A_B$  **couple to  $\delta g$**
- The **charged current** gauge fields **decouple** from  $\delta g$


# Perturbations : Symmetries

The boundary plasma has an **SO(3) rotational invariance**

$$\langle J_\lambda J_\sigma \rangle^R(\omega, \vec{k}) = P^\perp(\omega, \vec{k})_{\lambda\sigma} i\Pi^\perp(\omega, \mathbf{k}) + P^\parallel(\omega, \vec{k})_{\lambda\sigma} i\Pi^\parallel(\omega, \mathbf{k})$$

For a given **mode**  $(\omega, \vec{k})$ , it reduces to an **SO(2) subgroup**

The perturbations are divided into **helicity sectors** that decouple



Helicity	Gauge field	Metric
$h = 0$	$\delta A_0, \delta A_3$	$\delta g_0^0, \delta g_0^3, \delta g_3^3, \delta g_1^1 + \delta g_2^2$
$h = 1$	$\delta A_{1,2}$	$\delta g_0^{1,2}, \delta g_3^{1,2}$
$h = 2$	—	$\delta g_2^1, \delta g_1^1 - \delta g_2^2$

# Sector decoupled from the metric

Consider  $\delta A_\mu$  that decouples from  $\delta g_{\mu\nu}$

The modes are organized in terms of the **gauge-invariants** under

$$U(1) : \delta A \rightarrow \delta A + d\delta\lambda$$

$h = 1$	$h = 0$
$\delta A_1, \delta A_2$	$E^\parallel \equiv \omega\delta A_3 + k\delta A_0$

The linearized **Maxwell equations** in each helicity sector can be written in terms of the gauge-invariants

The  $\Pi$ 's are extracted from the **solutions near the boundary** ( $z \rightarrow 0$ )

$$\Pi^\perp \propto -\frac{\ell}{z} \frac{\partial_z \delta A_1}{\delta A_1} \Big|_{z \rightarrow 0}, \quad \Pi^\parallel \propto -\frac{\ell}{z} \frac{\partial_z \delta E^\parallel}{\delta E^\parallel} \Big|_{z \rightarrow 0}.$$

# Sector coupled to the metric

$\delta T_{MN} \propto \delta A_B$  couples to  $\delta g_{\mu\nu}$

Again, organize the modes in terms of the **gauge-invariants** under :

○  $U(1) : \delta X \rightarrow \delta X + d\delta\lambda$

○ **Diffeomorphisms** :

$$\delta X_M \rightarrow \delta X_M + \delta\xi^N \partial_N \bar{X}_M + \bar{X}_N \partial_M \delta\xi^N$$

$$\delta g_{MN} \rightarrow \delta g_{MN} + \nabla_M \delta\xi_N + \nabla_N \delta\xi_M$$

$h = 1$	$h = 0$
$\delta X_{1,2}$	$\delta S_1 \equiv \omega \delta X_3 + k \delta X_0 + a(z) \mu k (\delta g_1^1 + \delta g_2^2)$
$\delta Y^{1,2} \equiv k \delta g_0^{1,2} + \omega \delta g_3^{1,2}$	$\delta S_2 \equiv 2\omega k \delta g_0^3 + \omega^2 \delta g_z^z - f(z) k^2 \delta g_0^0 + b(z, \omega/k) k^2 (\delta g_1^1 + \delta g_2^2)$

# Sector coupled to the metric

The linearized **Einstein-Maxwell equations** in each helicity sector can be written in terms of the **gauge-invariants** :

- $\mathbf{h} = \mathbf{1}$  : 2 coupled 2<sup>nd</sup> order ODE's for  $\delta X_{1,2}$  and  $\delta Y^{1,2}$
- $\mathbf{h} = \mathbf{0}$  : 2 coupled 2<sup>nd</sup> order ODE's for  $\delta S_1$  and  $\delta S_2$

The  $\Pi$ 's are extracted from the **solutions near the boundary** ( $z \rightarrow 0$ )

$$\mathbf{h} = \mathbf{1} : \quad \delta X_1 = \delta \hat{X}_1 + z^2 \delta \Pi_{X_1} + \dots, \quad \delta \Pi_{X_1} \equiv \mathbf{\Pi}_{\mathbf{X}\mathbf{X}}^\perp \delta \hat{X}_1 + \Pi_{\mathbf{X}\mathbf{Y}}^\perp \delta \hat{Y}^1,$$

Compute **2 solutions** and invert the linear relation

$$\left( \mathbf{\Pi}_{\mathbf{X}\mathbf{X}}^\perp \quad \Pi_{\mathbf{X}\mathbf{Y}}^\perp \right) = \left( \delta \Pi_{X_1}^{(1)} \quad \delta \Pi_{X_1}^{(2)} \right) \begin{pmatrix} \delta \hat{X}_1^{(1)} & \delta \hat{X}_1^{(2)} \\ \delta \hat{Y}_{(1)}^1 & \delta \hat{Y}_{(2)}^1 \end{pmatrix}^{-1}$$