


Precision nucleon and nuclear structure from light (muonic) atoms



Randolf Pohl

Johannes Gutenberg
Universität Mainz

ECT* Trento
28.07.25




Hydrogen-like atoms



“There's a reason physicists are so successful with what they do,
and that is they study the hydrogen atom and the helium ion
and then they stop.”

- Richard Feynman

Hydrogen-like atoms



“There's a reason physicists are so successful with what they do,
and that is they study the hydrogen atom and the helium ion
and then they stop.”

- Richard Feynman

Welcome to ECT* Trento



ECT*
EUROPEAN CENTRE
FOR THEORETICAL STUDIES
IN NUCLEAR PHYSICS AND RELATED AREAS


New perspectives in the charge radii
determination for light nuclei

July 28, 2025 to August 1, 2025

ECT*

Europe/Rome timezone

Cool muonic stuff|



ECT*, 28.7.2025

Randolf Pohl, JGU Mainz




Organizers: Loredana **Gastaldo**
Franziska **Hagelstein**
Nancy **Paul**
Randolf **Pohl**

Main topics:

- High-resolution X-ray and laser spectroscopy
- Microcalorimeter X-ray detector technology
- Experimental progress: target preparation, background suppression, calibration
- Theory of muonic atoms
- Nuclear and nucleon structure effects
- Searches for physics beyond the Standard model (BSM)

Welcome to ECT* Trento

		Monday	Tuesday	Wednesday	Thursday	Friday
9:40 – 10:20	30 + 10	van Kolck	Ohayon	Bacca	Gasparian	Craig
10:20 – 11:00	30 + 10	Pohl	Peset	Drissi	Ouf	Potvliege
11:00 – 11:30	Coffee					
11:30 – 12:10	30 + 10	Gorshteyn	Harman	Pastore	Pitelis	Feng
12:10 – 12:50	30 + 10	Fleischmann	Nörtershäuser	Oreshkina	Lensky	Closing
13:00 – 15:00	Lunch					
15:00 – 15:40	30 + 10	Kreuzberger	Weber	Excursion	Ji	
15:40 – 16:20	30 + 10	Godhino	Rathi		Egert	
16:20 – 16:50		Coffee				
16:50 – 17:30	30 + 10	Baptista	Cocolios			
17:30 – 18:10	30 + 10	Heiße (Zoom)	Deseyn			
18:10 – 18:30	Discussions					






Stuff with lasers

The “Proton Radius Puzzle”

Measuring R_p using **electrons**: 0.88 fm ($\pm 0.7\%$)
using **muons**: 0.84 fm ($\pm 0.05\%$)




μd 2016: RP et al (CREMA Coll.) Science 353, 669 (2016)

μp 2013: A. Antognini, RP et al (CREMA Coll.) Science 339, 417 (2013)

The “Proton Radius Puzzle”

Measuring R_p using **electrons**: 0.84 fm
using **muons**: 0.84 fm



μd 2016: RP et al (CREMA Coll.) Science 353, 669 (2016)
 μp 2013: A. Antognini, RP et al (CREMA Coll.) Science 339,

ECT*, 28.7.2025

Randolf Pohl, J



The “Proton Radius Puzzle”



The “Proton Radius Puzzle”



Muonic Atoms

A **bare** nucleus, orbited by **one negative muon**.

Muonic Atoms

A **bare** nucleus, orbited by **one negative muon**.

Hydrogen theory!

What is a Muon?



Carl David Anderson

Nobel prize 1936
(for the Positron!)



Seth Neddermeyer

Muons are **heavy electrons**:
200x the mass of an electron.

Discovered in 1936 by Carl D. Anderson and
Seth Neddermeyer as **part of the cosmic
radiation**.

Muon have a **lifetime of 2 microseconds**.
(millionths of a second).

Isidor Isaac Rabi

What is




Carl David Anderson

Nobel prize 1936
(for the Positron!)



Seth Neddermeyer



What is a Muon?



Carl David Anderson

Nobel prize 1936
(for the Positron!)



Seth Neddermeyer

Muons are **heavy electrons**:
200x the mass of an electron.

Discovered in 1936 by Carl D. Anderson and Seth Neddermeyer as **part of the cosmic radiation**.

Muon have a **lifetime of 2**

Who ordered that?!




Isidor Isaac Rabi

Muonic atoms in a nutshell

Regular hydrogen:

Bohr radius $\sim 50'000$ x nuclear radius



Muonic hydrogen:



Muon **mass** = **200** * electron mass

Bohr **radius** = **1/200** of H

200³ = a **ten million times** more
sensitive to nuclear size & structure

=> Our (laser) spectroscopy at **10⁻⁵** level
can compete with **10⁻¹²** from normal atoms


$$\Delta E_{\text{FNS}}(n, l) = \frac{2c^4}{3\hbar^2} \frac{(Z\alpha)^4}{n^3} m_r^3 r^2 \delta_{l0}$$




Vastly not to scale!!

Randolf Pohl, JGU Mainz


Lamb shift in Muonic Hydrogen




The accelerator at PSI




The muon beam line in $\pi E5$




Inside the laser hut




Inside the laser hut



Laser beam line




The hydrogen target




Time Spectra

13 hours of data




Time Spectra


13 hours of data




Time Spectra




Time Spectra




Resonance



Resonance vs Result



Resonance vs Result





The total predicted $2S_{1/2}^{F=1} - 2P_{3/2}^{F=2}$ energy difference, $\Delta\tilde{E}$, in muonic hydrogen is the sum of radiative, recoil, and proton structure contributions, and the fine and hyperfine splittings for our particular transition, and it is given^{8,11-15} by

$$\Delta\tilde{E} = 209.9779(49) - 5.2262 r_p^2 + 0.0347 r_p^3 \text{ meV} \quad (1)$$

where $r_p = \sqrt{\langle r_p^2 \rangle}$ is given in fm. A detailed derivation of equation (1) is given in Supplementary Information.

Measurement NOT useful without THEORY!

Resonance vs Result



$$\Delta\tilde{E} = 209.9779(49) - 5.2262 r_p^2 + 0.0347 r_p^3 \text{ meV} \quad (1)$$

Measurement NOT useful without THEORY!

Theory in muonic H

$$\Delta E_{\text{Lamb}} = 206.0336 (15) \text{ meV}_{\text{QED}} + 0.0332 (20) \text{ meV}_{\text{TPE}} - 5.2275 (10) \text{ meV/fm}^2 * R_p^2$$


Annals of Physics 331 (2013) 127–145



Contents lists available at SciVerse ScienceDirect

Annals of Physics

journal homepage: www.elsevier.com/locate/aop



Theory of the 2S–2P Lamb shift and 2S hyperfine splitting in muonic hydrogen



Aldo Antognini^{a,*}, Franz Kottmann^a, François Biraben^b, Paul Indelicato^b,
François Nez^b, Randolph Pohl^c

^a Institute for Particle Physics, ETH Zurich, 8093 Zurich, Switzerland

^b Laboratoire Kastler Brossel, École Normale Supérieure, CNRS and Université P. et M. Curie, 75252 Paris, CEDEX 05, France

^c Max-Planck-Institut für Quantenoptik, 85748 Garching, Germany

Our attempt to summarize all the original work by many theorists....


ECT*, 28.7.2025

Randolf Pohl, JGU Mainz


Theory in muonic H

$$\Delta E_{\text{Lamb}} = 206.0344 (3) \text{ meV}_{\text{QED}} + 0.0289 (25) \text{ meV}_{\text{TPE}} - 5.2259 (1) \text{ meV/fm}^2 * R_p^2$$


Uehling



Källen-Sabry




Muon SE+VP



and 20+ more....

Proton form factor



elastic and inelastic two-photon
exchange
(Friar moment and polarizability)

Theory I: “pure” QED

Table 1

All known radius-independent contributions to the Lamb shift in μp from different authors, and the one we selected. Values are in meV. The entry # in the first column refers to Table 1 in Ref. [13]. The “finite-size to relativistic recoil correction” (entry #18 in [13]), which depends on the proton structure, has been shifted to Table 2, together with the small terms #26 and #27, and the proton polarizability term #25. SE: self-energy, VP: vacuum polarization, LBL: light-by-light scattering, Rel: relativistic, NR: non-relativistic, RC: recoil correction.

#	Contribution	Pachucki [10,11]	Nature [13]	Borie-v6 [79]	Indelicato [80]	Our choice	Ref.
1	NR one-loop electron VP (eVP)	205.0074					
2	Rel. corr. (Breit–Pauli)	0.0169 ^a					
3	Rel. one-loop eVP		205.0282	205.0282	205.02821	205.02821	[80] Eq. (54)
19	Rel. RC to eVP, $\alpha(Z\alpha)^4$	(incl. in #2) ^b	−0.0041	−0.0041		−0.00208 ^c	[77,78]
4	Two-loop eVP (Källén–Sabry)	1.5079	1.5081	1.5081	1.50810	1.50810	[80] Eq. (57)
5	One-loop eVP in 2-Coulomb lines $\alpha^2(Z\alpha)^5$	0.1509	0.1509	0.1507	0.15102	0.15102	[80] Eq. (60)
7	eVP corr. to Källén–Sabry	0.0023	0.00223	0.00223	0.00215	0.00215	[80] Eq. (62), [87]
6	NR three-loop eVP	0.0053	0.00529	0.00529		0.00529	[87,88]
9	Wichmann–Kroll, “1:3” LBL		−0.00103	−0.00102	−0.00102	−0.00102	[80] Eq. (64), [89]
10	Virtual Delbrück, “2:2” LBL		0.00135	0.00115		0.00115	[74,89]
New	“3:1” LBL			−0.00102		−0.00102	[89]
20	μSE and μVP	−0.6677	−0.66770	−0.66788	−0.66761	−0.66761	[80] Eqs. (72) + (76)
11	Muon SE corr. to eVP $\alpha^2(Z\alpha)^4$	−0.005(1)	−0.00500	−0.004924 ^d		−0.00254	[85] Eq. (29a) ^e
12	eVP loop in self-energy $\alpha^2(Z\alpha)^4$	−0.001	−0.00150				[74,90–92]
21	Higher order corr. to μSE and μVP		−0.00169	−0.00171 ^g		−0.00171	[86] Eq. (177)
13	Mixed eVP + μVP		0.00007	0.00007		0.00007	[74]
New	eVP and μVP in two Coulomb lines				0.00005	0.00005	[80] Eq. (78)
14	Hadronic VP $\alpha(Z\alpha)^4 m_r$	0.0113(3)	0.01077(38)	0.011(1)		0.01121(44)	[93–95]
15	Hadronic VP $\alpha(Z\alpha)^5 m_r$		0.000047			0.000047	[94,95]
16	Rad corr. to hadronic VP		−0.000015			−0.000015	[94,95]
17	Recoil corr.	0.0575	0.05750	0.0575	0.05747	0.05747	[80] Eq. (88)
22	Rel. RC $(Z\alpha)^5$	−0.045	−0.04497	−0.04497	−0.04497	−0.04497	[80] Eq. (88), [74]
23	Rel. RC $(Z\alpha)^6$	0.0003	0.00030		0.0002475	0.0002475	[80] Eq. (86)+Tab.II
New	Rad. (only eVP) RC $\alpha(Z\alpha)^5$					0.000136	[85] Eq. (64a)
24	Rad. RC $\alpha(Z\alpha)^n$ (proton SE)	−0.0099	−0.00960	−0.0100		−0.01080(100)	[43] ^h [74]
	Sum	206.0312	206.02915	206.02862		206.03339(109)	

Antognini, RP et al., Ann. Phys. (N.Y.) 331, 127 (2013)

Theory

Comprehensive theory of the Lamb shift in light muonic atoms

K. Pachucki 

Faculty of Physics, University of Warsaw, Pasteura 5, 02-093 Warsaw, Poland

V. Lensky 

Institut für Kernphysik, Johannes Gutenberg-Universität Mainz, 55128 Mainz, Germany

F. Hagelstein 

*Institut für Kernphysik, Johannes Gutenberg-Universität Mainz, 55128 Mainz, Germany
and Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland*

S. S. Li Muli 

Institut für Kernphysik, Johannes Gutenberg-Universität Mainz, 55128 Mainz, Germany

S. Bacca 

*Institut für Kernphysik, Johannes Gutenberg-Universität Mainz, 55128 Mainz, Germany
and Helmholtz-Institut Mainz, Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany*

R. Pohl 

Institut für Physik, Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany



(published 24 January 2024)

A comprehensive theory of the Lamb shift in light muonic atoms such as μH , μD , $\mu^3\text{He}^+$, and $\mu^4\text{He}^+$ is presented, with all quantum electrodynamic corrections included at the precision level constrained by the uncertainty of nuclear structure effects. This analysis can be used in the global adjustment of fundamental constants and in the determination of nuclear charge radii. Further improvements in the understanding of electromagnetic interactions of light nuclei will allow for a promising test of fundamental interactions by comparison with “normal” atomic spectroscopy, in particular, with H-D and ^3He - ^4He isotope shifts.

DOI: [10.1103/RevModPhys.96.015001](https://doi.org/10.1103/RevModPhys.96.015001)

ECT*, 28.7.2025

Randolf Pohl, JGU Mainz

Pachucki, Lensky, Hagelstein, LiMuli,
Bacca, RP, RMP 96 (2024)

TABLE I. Contributions to the $2P_{1/2} - 2S_{1/2}$ energy difference E_L in meV, with the charge radii r_C given in fm. All corrections larger than 3% of the overall uncertainty are included. Theoretical predictions for E_L are $E_L(\text{theo}) = E_{\text{QED}} + Cr_C^2 + E_{\text{NS}}$. The last two rows show the values of r_C determined from a comparison of $E_L(\text{theo})$ to $E_L(\text{exp})$.

Section	Order	Correction	μH	μD	$\mu^3\text{He}^+$	$\mu^4\text{He}^+$
III.A	$\alpha(Z\alpha)^2$	eVP ⁽¹⁾	205.007 38	227.634 70	1641.886 2	1665.773 1
III.A	$\alpha^2(Z\alpha)^2$	eVP ⁽²⁾	1.658 85	1.838 04	13.084 3	13.276 9
III.A	$\alpha^3(Z\alpha)^2$	eVP ⁽³⁾	0.007 52	0.008 42(7)	0.073 0(30)	0.074 0(30)
III.B	$(Z, Z^2, Z^3)\alpha^5$	Light-by-light eVP	−0.000 89(2)	−0.000 96(2)	−0.013 4(6)	−0.013 6(6)
III.C	$(Z\alpha)^4$	Recoil	0.057 47	0.067 22	0.126 5	0.295 2
III.D	$\alpha(Z\alpha)^4$	Relativistic with eVP ⁽¹⁾	0.018 76	0.021 78	0.509 3	0.521 1
III.E	$\alpha^2(Z\alpha)^4$	Relativistic with eVP ⁽²⁾	0.000 17	0.000 20	0.005 6	0.005 7
III.F	$\alpha(Z\alpha)^4$	$\mu\text{SE}^{(1)} + \mu\text{VP}^{(1)}$, LO	−0.663 45	−0.769 43	−10.652 5	−10.926 0
III.G	$\alpha(Z\alpha)^5$	$\mu\text{SE}^{(1)} + \mu\text{VP}^{(1)}$, NLO	−0.004 43	−0.005 18	−0.174 9	−0.179 7
III.H	$\alpha^2(Z\alpha)^4$	$\mu\text{VP}^{(1)}$ with eVP ⁽¹⁾	0.000 13	0.000 15	0.003 8	0.003 9
III.I	$\alpha^2(Z\alpha)^4$	$\mu\text{SE}^{(1)}$ with eVP ⁽¹⁾	−0.002 54	−0.003 06	−0.062 7	−0.064 6
III.J	$(Z\alpha)^5$	Recoil	−0.044 97	−0.026 60	−0.558 1	−0.433 0
III.K	$\alpha(Z\alpha)^5$	Recoil with eVP ⁽¹⁾	0.000 14(14)	0.000 09(9)	0.004 9(49)	0.003 9(39)
III.L	$Z^2\alpha(Z\alpha)^4$	nSE ⁽¹⁾	−0.009 92	−0.003 10	−0.084 0	−0.050 5
III.M	$\alpha^2(Z\alpha)^4$	$\mu F_1^{(2)}, \mu F_2^{(2)}, \mu\text{VP}^{(2)}$	−0.001 58	−0.001 84	−0.031 1	−0.031 9
III.N	$(Z\alpha)^6$	Pure recoil	0.000 09	0.000 04	0.001 9	0.001 4
III.O	$\alpha(Z\alpha)^5$	Radiative recoil	0.000 22	0.000 13	0.002 9	0.002 3
III.P	$\alpha(Z\alpha)^4$	hVP	0.011 36(27)	0.013 28(32)	0.224 1(53)	0.230 3(54)
III.Q	$\alpha^2(Z\alpha)^4$	hVP with eVP ⁽¹⁾	0.000 09	0.000 10	0.002 6(1)	0.002 7(1)
IV.A	$(Z\alpha)^4$	r_C^2	−5.197 5 r_p^2	−6.073 2 r_d^2	−102.523 r_h^2	−105.322 r_a^2
IV.B	$\alpha(Z\alpha)^4$	eVP ⁽¹⁾ with r_C^2	−0.028 2 r_p^2	−0.034 0 r_d^2	−0.851 r_h^2	−0.878 r_a^2
IV.C	$\alpha^2(Z\alpha)^4$	eVP ⁽²⁾ with r_C^2	−0.000 2 r_p^2	−0.000 2 r_d^2	−0.009(1) r_h^2	−0.009(1) r_a^2
V.A	$(Z\alpha)^5$	TPE	0.029 2(25)	1.979(20)	16.38(31)	9.76(40)
V.B	$\alpha^2(Z\alpha)^4$	Coulomb distortion	0.0	−0.261	−1.010	−0.536
V.C	$(Z\alpha)^6$	3PE	−0.001 3(3)	0.002 2(9)	−0.214(214)	−0.165(165)
V.D	$\alpha(Z\alpha)^5$	eVP ⁽¹⁾ with TPE	0.000 6(1)	0.027 5(4)	0.266(24)	0.158(12)
V.E	$\alpha(Z\alpha)^5$	$\mu\text{SE}^{(1)} + \mu\text{VP}^{(1)}$ with TPE	0.000 4	0.002 6(3)	0.077(8)	0.059(6)
III	E_{QED}	Point nucleus	206.034 4(3)	228.774 0(3)	1644.348(8)	1668.491(7)
IV	Cr_C^2	Finite size	−5.225 9 r_p^2	−6.107 4 r_d^2	−103.383 r_h^2	−106.209 r_a^2
V	E_{NS}	Nuclear structure	0.028 9(25)	1.750 3(200)	15.499(378)	9.276(433)
	$E_L(\text{exp})$	Experiment ^a	202.370 6(23)	202.878 5(34)	1258.598(48)	1378.521(48)
	r_C	This review	0.840 60(39)	2.127 58(78)	1.970 07(94)	1.678 6(12)
	r_C	Previous work ^a	0.840 87(39)	2.125 62(78)	1.970 07(94)	1.678 24(83)

Theory

Pachucki, Lensky, Hagelstein, LiMuli, Bacca, RP, RMP 96 (2024)

TABLE I. Contributions to the $2P_{1/2} - 2S_{1/2}$ energy difference E_L in meV, with the charge radii r_C given in fm. All corrections larger than 3% of the overall uncertainty are included. Theoretical predictions for E_L are $E_L(\text{theo}) = E_{\text{QED}} + Cr_C^2 + E_{\text{NS}}$. The last two rows show the values of r_C determined from a comparison of $E_L(\text{theo})$ to $E_L(\text{exp})$.


Section	Order	Correction	μH	μD	$\mu^3\text{He}^+$	$\mu^4\text{He}^+$
III.A	$\alpha(Z\alpha)^2$	eVP ⁽¹⁾	205.007 38	227.634 70	1641.886 2	1665.773 1
III.A	$\alpha^2(Z\alpha)^2$	eVP ⁽²⁾	1.658 85	1.838 04	13.084 3	13.276 9
III.A	$\alpha^3(Z\alpha)^2$	eVP ⁽³⁾	0.007 52	0.008 42(7)	0.073 0(30)	0.074 0(30)
III.B	$(Z, Z^2, Z^3)\alpha^5$	Light-by-light eVP	−0.000 89(2)	−0.000 96(2)	−0.013 4(6)	−0.013 6(6)
III.C	$(Z\alpha)^4$	Recoil	0.057 47	0.067 22	0.126 5	0.295 2
III.D	$\alpha(Z\alpha)^4$	Relativistic with eVP ⁽¹⁾	0.018 76	0.021 78	0.509 3	0.521 1
III.E	$\alpha^2(Z\alpha)^4$	Relativistic with eVP ⁽²⁾	0.000 17	0.000 20	0.005 6	0.005 7
III.F	$\alpha(Z\alpha)^4$	$\mu\text{SE}^{(1)} + \mu\text{VP}^{(1)}$, LO	−0.663 45	−0.769 43	−10.652 5	−10.926 0
III.G	$\alpha(Z\alpha)^5$	$\mu\text{SE}^{(1)} + \mu\text{VP}^{(1)}$, NLO	−0.004 43	−0.005 18	−0.174 9	−0.179 7
III.H	$\alpha^2(Z\alpha)^4$	$\mu\text{VP}^{(1)}$ with eVP ⁽¹⁾	0.000 13	0.000 15	0.003 8	0.003 9
III.I	$\alpha^2(Z\alpha)^4$	$\mu\text{SE}^{(1)}$ with eVP ⁽¹⁾	−0.002 54	−0.003 06	−0.062 7	−0.064 6
III.J	$(Z\alpha)^5$	Recoil	−0.044 97	−0.026 60	−0.558 1	−0.433 0
III.K	$\alpha(Z\alpha)^5$	Recoil with eVP ⁽¹⁾	0.000 14(14)	0.000 09(9)	0.004 9(49)	0.003 9(39)
III.L	$Z^2\alpha(Z\alpha)^4$	nSE ⁽¹⁾	−0.009 92	−0.003 10	−0.084 0	−0.050 5
III.M	$\alpha^2(Z\alpha)^4$	$\mu F_1^{(2)}, \mu F_2^{(2)}, \mu\text{VP}^{(2)}$	−0.001 58	−0.001 84	−0.031 1	−0.031 9
III.N	$(Z\alpha)^6$	Pure recoil	0.000 09	0.000 04	0.001 9	0.001 4
III.O	$\alpha(Z\alpha)^5$	Radiative recoil	0.000 22	0.000 13	0.002 9	0.002 3
III.P	$\alpha(Z\alpha)^4$	hVP	0.011 36(27)	0.013 28(32)	0.224 1(53)	0.230 3(54)
III.Q	$\alpha^2(Z\alpha)^4$	hVP with eVP ⁽¹⁾	0.000 09	0.000 10	0.002 6(1)	0.002 7(1)
IV.A	$(Z\alpha)^4$	2	−5.107 5(2)	−6.073 2(2)	−102.522(2)	−105.222(2)
V.A	$(Z\alpha)^5$	TPE			0.029 2(25)	1.979(20)
V.B	$\alpha^2(Z\alpha)^4$	Coulomb distortion			0.0	−0.261
V.C	$(Z\alpha)^6$	3PE			−0.001 3(3)	0.002 2(9)
V.D	$\alpha(Z\alpha)^5$	eVP ⁽¹⁾ with TPE			0.000 6(1)	0.027 5(4)
V.E	$\alpha(Z\alpha)^5$	$\mu\text{SE}^{(1)} + \mu\text{VP}^{(1)}$ with TPE			0.000 4	0.002 6(3)
III	E_{QED}	Point nucleus	206.034 4(3)	228.774 0(3)	1644.348(8)	1668.491(7)
IV	Cr_C^2	Finite size	−5.225 9 r_p^2	−6.107 4 r_d^2	−103.383 r_h^2	−106.209 r_a^2
V	E_{NS}	Nuclear structure	0.028 9(25)	1.750 3(200)	15.499(378)	9.276(433)
	$E_L(\text{exp})$	Experiment ^a	202.370 6(23)	202.878 5(34)	1258.598(48)	1378.521(48)
	r_C	This review	0.840 60(39)	2.127 58(78)	1.970 07(94)	1.678 6(12)
	r_C	Previous work ^a	0.840 87(39)	2.125 62(78)	1.970 07(94)	1.678 24(83)

Theory

Pachucki, Lensky, Hagelstein, LiMuli, Bacca, RP, RMP 96 (2024)

The “Proton Radius Puzzle”

Measuring R_p using **electrons**: 0.88 fm ($\pm 0.7\%$)
using **muons**: 0.84 fm ($\pm 0.05\%$)



μd 2016: RP et al (CREMA Coll.) Science 353, 669 (2016)

μp 2013: A. Antognini, RP et al (CREMA Coll.) Science 339, 417 (2013)

ECT* Workshop 2012

“Fitting wars”

TPE and The Subtraction Term

Missing QED contributions?

Form factors vs. charge density


BSM Physics

Muon Scattering


muH-HFS



Hydrogen energy levels




Rp from H spectroscopy



Situation 2016


RP et al., Metrologia 54, L1 (2017)

2S-nP in Garching




- cryogenic H beam (6 K)
- optical 1S-2S excitation (2S, $F=0$)
- 2S-nP transition is 1-photon
→ retroreflector (“AFR”)
- split line to 10^{-4}
- 0.5 kHz vs. 9 kHz PRP

2S-nP in Garching




1st order Doppler cancellation (2S-4P)




2S-nP in Garching

Systematics (2S-6P)




Contribution	$\Delta\nu$ (kHz)	σ (kHz)
First-order Doppler shift	0.34	0.43
Extrapolation (statistical)	0.34	0.43
Simulation of atom speeds	—	0.01
Simulation corrections	1.05	0.17
Light force shift	1.15	0.17
Quantum interference shift	0.05	0.02
Second-order Doppler shift	−0.14	0.01
dc-Stark shift	0.05	0.07
BBR-induced shift	0.28	0.01
Zeeman shift	0.00	0.08
Pressure shift	0.00	0.02
Sampling bias	0.00	0.06
Signal background	0.00	0.03
Laser spectrum	0.00	0.07
Frequency standard	0.02	0.01
Subtotal (experiment-specific contributions)	1.74	0.48
Recoil shift	−1 176.03	0.00
HFS correction $\Delta\nu_{\text{HFS}}(\nu_{2\text{S-6P}})$	−132 985.25	0.01
Total (all contributions)	−134 159.54	0.48


The proton radius situation today





Not really “solved”

H (2S-6P)
preliminary













Muonic Deuterium



Pohl et al. (CREMA), Science (2016)

2.5 transitions in muonic D



$$\Delta E_{\text{LS}}^{\text{exp}} = 202.8785(31)_{\text{stat}}(14)_{\text{syst}} \text{ meV}$$

$$\Delta E_{\text{HFS}}^{\text{exp}} = 6.2747(70)_{\text{stat}}(20)_{\text{syst}} \text{ meV}$$

Pohl et al. (CREMA Coll.), Science 353, 669 (2016)

Theory in muonic D

$$\Delta E_{\text{Lamb}}^{\mu\text{D}} = 228.7854 (13) \text{ meV}_{\text{QED}} + 1.7150 (230) \text{ meV}_{\text{TPE}} - 6.1103 (3) \text{ meV/fm}^2 * R_d^2$$



Nuclear structure contributions to the Lamb shift in muonic deuterium.

Item	Contribution	Pachucki [55] AV18		Friar [60] ZRA		Hernandez <i>et al.</i> [58] AV18 N ³ LO [†]		Pach.& Wienczek [65] AV18		Carlson <i>et al.</i> [64] data	Our choice value source	
	Source	1		2		3 4		5		6		
p1	Dipole	1.910	$\delta_0 E$	1.925	Leading C1	1.907	1.926	$\delta_{D1}^{(0)}$	1.910	$\delta_0 E$	1.9165	± 0.0095 3-5
p2	Rel. corr. to p1, longitudinal part	-0.035	$\delta_R E$	-0.037	Subleading C1	-0.029	-0.030	$\delta_L^{(0)}$	-0.026	$\delta_R E$		
p3	Rel. corr. to p1, transverse part					0.012	0.013	$\delta_T^{(0)}$				
p4	Rel. corr. to p1, higher-order								0.004	$\delta_{HO} E$		
sum	Total rel. corr., p2+p3+p4	-0.035		-0.037		-0.017	-0.017		-0.022		-0.0195	± 0.0025 3-5
p5	Coulomb distortion, leading	-0.255	$\delta_{C1} E$						-0.255	$\delta_{C1} E$		
p6	Coul. distortion, next order	-0.006	$\delta_{C2} E$						-0.006	$\delta_{C2} E$		
sum	Total Coulomb distortion, p5+p6	-0.261				-0.262	-0.264	$\delta_C^{(0)}$	-0.261		-0.2625	± 0.0015 3-5
p7	El. monopole excitation	-0.045	$\delta_{Q0} E$	-0.042	C0	-0.042	-0.041	$\delta_{R2}^{(2)}$	-0.042	$\delta_{Q0} E$		
p8	El. dipole excitation	0.151	$\delta_{Q1} E$	0.137	Retarded C1	0.139	0.140	$\delta_{D1}^{(2)}$	0.139	$\delta_{Q1} E$		
p9	El. quadrupole excitation	-0.066	$\delta_{Q2} E$	-0.061	C2	-0.061	-0.061	$\delta_Q^{(2)}$	-0.061	$\delta_{Q2} E$		
sum	Tot. nuclear excitation, p7+p8+p9	0.040		0.034	C0 + ret-C1 + C2	0.036	0.038		0.036		0.0360	± 0.0020 2-5
p10	Magnetic	-0.008 $\diamond^a$	$\delta_M E$	-0.011	M1	-0.008	-0.007	$\delta_M^{(0)}$	-0.008	$\delta_M E$	-0.0090	± 0.0020 2-5
SUM.1	Total nuclear (corrected)	1.646		1.648 b		1.656	1.676		1.655		1.6615	± 0.0103
p11	Finite nucleus size			0.021	Retarded C1 f.s.	0.020 $\diamond^c$	0.021 $\diamond^c$	$\delta_{NS}^{(2)}$	0.020	$\delta_{FS} E$		
p12	n p charge correlation			-0.023	pn correl. f.s.	-0.017	-0.017	$\delta_{np}^{(1)}$	-0.018	$\delta_{FZE} E$		
sum	p11+p12			-0.002		0.003	0.004		0.002		0.0010	± 0.0030 2-5
p13	Proton elastic 3rd Zemach moment	} 0.043(3)	$\delta_P E$	0.030	$\langle r^3 \rangle_{(2)}^{pp}$	} 0.027(2)	δ_{pol}^N [64]	$\delta_{D1}^{(2)}$	} 0.043(3)	$\delta_P E$	} 0.028(2)	ΔE^{hadr}
p14	Proton inelastic polarizab.											
p15	Neutron inelastic polarizab.											
p16	Proton & neutron subtraction term											
sum	Nucleon TPE, p13+p14+p15+p16	0.043(3)		0.030		0.027(2)			0.059(9)		-0.0098	± 0.0098 Eq.(15) ^e
SUM.2	Total nucleon contrib.	0.043(3)		0.028		0.030(2)			0.061(9)		0.0471	± 0.0101 f
	Sum, published	1.680(16)		1.941(19)		1.690(20)			1.717(20)		2.011(740)	
	Sum, corrected			1.697(19) g		1.714(20) h			1.707(20) i		1.748(740) j	
											1.7096	± 0.0147

Krauth, RP et al., Ann. Phys. (N.Y.) 366, 168 (2016)

ECT*, 28.7.2025

Randolf Pohl, JGU Mainz

+ Pachucki et al., PRA 97, 062511 (2018)

+ Hernandez et al., PLB 778, 377 (2018)

Theory in muonic D

$$\Delta E_{\text{Lamb}}^{\mu\text{D}} = 228.7854 (13) \text{ meV}_{\text{QED}} + 1.7150 (230) \text{ meV}_{\text{TPE}} - 6.1103 (3) \text{ meV/fm}^2 * R_d^2$$

Nuclear structure contributions to the Lamb shift in muonic deuterium.

Item	Table 3 Nuclear polarization contributions to the 2S–2P Lamb shift ΔE [meV] in μD with different potentials.					
		Ref. [29]	AV18	N ³ LO-EM	N ³ LO-EGM	
p1						
p2						
p3						
p4	$\delta^{(0)}$					
sum	$\delta_{D1}^{(0)}$	−1.910	−1.907	−1.912	(−1.911, −1.926)	
p5	$\delta_L^{(0)}$	0.035	0.029	0.029	(0.029, 0.030)	
p6	$\delta_T^{(0)}$	−	−0.012	−0.012	−0.013	
sum	$\delta_C^{(0)}$	0.261	0.262	0.262	(0.262, 0.264)	
p7	$\delta_M^{(0)}$	0.016	0.008	0.007	0.007	
p8	$\delta_{Z3}^{(1)}$	−	0.357	0.359	(0.359, 0.363)	
p9	$\delta_{R2}^{(2)}$	0.045	0.042	0.041	0.041	
sum	$\delta_Q^{(2)}$	0.066	0.061	0.061	0.061	
p10	$\delta_{D1D3}^{(2)}$	−0.151	−0.139	−0.139	(−0.139, −0.140)	
SUM.1	$\delta_{NS}^{(1)}$	−	0.064	0.064	(0.064, 0.065)	
p11	$\delta_{np}^{(1)}$	−	0.017	0.017	0.017	
p12	$\delta_{NS}^{(2)}$	−	−0.015	−0.015	−0.015	
sum	δ_{pol}^A	−	−1.235	−1.237	(−1.236, −1.246)	
p13	δ_{Zem}^A	−	−0.421	−0.423	(−0.424, −0.428)	
p14	$\delta_{pol}^A + \delta_{Zem}^A$	−1.638	−1.656	−1.661	(−1.660, −1.674)	
p15						
p16						
sum						
SUM.2						

Hernandez <i>et al.</i> [58]	Pach.& Wienczek [65]	Carlson <i>et al.</i> [64]	Our choice	
18 N ³ LO [†]	AV18	data	value	source
4	5	6		
07 1.926 $\delta_{D1}^{(0)}$	1.910 $\delta_0 E$		1.9165 ± 0.0095	3-5
29 −0.030 $\delta_L^{(0)}$	−0.026 $\delta_R E$			
12 0.013 $\delta_T^{(0)}$	0.004 $\delta_{HO} E$			
17 −0.017	−0.022		−0.0195 ± 0.0025	3-5
	−0.255 $\delta_{C1} E$			
	−0.006 $\delta_{C2} E$			
32 −0.264 $\delta_C^{(0)}$	−0.261		−0.2625 ± 0.0015	3-5

O.J. Hernandez *et al.* / Physics Letters B 736 (2014) 344–349

nuclear structure theory with error bars!


0.027(2)	δ_{pol}^N [64]	$\left. \begin{array}{l} 0.043(3) \delta_P E \\ 0.016(8) \delta_N E \end{array} \right\}$	$\left. \begin{array}{l} 0.028(2) \Delta E^{\text{hadr}} \\ 0.0280 \pm 0.0020 \end{array} \right\}$	0.0289 ± 0.0015 Eq.(13) ^a
0.027(2)		0.059(9)		−0.0098 ± 0.0098 Eq.(15) ^c
0.030(2)		0.061(9)		0.0471 ± 0.0101 ^f
1.690(20)		1.717(20)	2.011(740)	
1.714(20) ^h		1.707(20) ⁱ	1.748(740) ^j	1.7096 ± 0.0147

Theory: Lamb shift in muonic D

$$\Delta E_{\text{Lamb}}^{\mu\text{D}} = 228.7740 (3) \text{ meV}_{\text{QED}} + 1.7503 (200) \text{ meV}_{\text{TPE}} - 6.1074 \text{ meV/fm}^2 * R_d^2$$

$$\Delta E_{\text{LS}}^{\text{exp}} = 202.8785(31)_{\text{stat}}(14)_{\text{syst}} \text{ meV}$$

Nuclear structure **two (and three!)-photon contributions** to the Lamb shift in muonic deuterium.



Pachucki, RP et al, arXiv 2212.13782

see also Krauth, RP et al. (2016) using calculations from
Pachucki (2011), Friar (2013), Carlson, Gorchtein, Vanderhaeghen
(2014), Hernandez et al. (2014), Pachucki + Wienczek (2015)



+ Pachucki et al., PRA 97, 062511 (2018): Sizeable three-photon !!

+ Hernandez et al., PLB 778, 377 (2018): χ EFT

+ Kalinowski (2019): eVP to nucl. struct.

+ Acharya et al., PRC 103, 024001 (2021)
 χ EFT + Dispersion relations

Muonic Deuterium



μD : 2.12758 (13)_{exp} (78)_{theo} fm

$\mu H + H/D(1S-2S)$: 2.12785 (17) fm

Theory in muonic D

$$\Delta E_{\text{Lamb}}^{\mu\text{D}} = 228.7740 (3) \text{ meV}_{\text{QED}} + 1.7503 (200) \text{ meV}_{\text{TPE}} - 6.1074 \text{ meV/fm}^2 * R_d^2$$

↓

$$\Delta E_{\text{TPE}} (\text{theo}) = 1.7503 \pm 0.0200 \text{ meV} \quad \text{Bacca group}$$

vs. $\pm 0.0034 \text{ meV}$ experimental uncertainty

(1) **charge radius**, using **calculated TPE**


$$r_d (\mu\text{D}) = 2.12758 (13)_{\text{exp}} (78)_{\text{theo}} \text{ fm}$$

(2) **polarizability**, using **charge radius from isotope shift**

$$\Delta E_{\text{TPE}} (\text{theo}) = 1.7503 (200) \text{ meV} \text{ vs.}$$


$$\Delta E_{\text{TPE}} (\text{exp}) = 1.7591 (59) \text{ meV} \quad \text{3x more accurate}$$

Muonic Deuterium




muonic

old electronic



$r_d^2 - r_p^2 = 3.82070(31) \text{ fm}^2$ H / D 1S-2S isotope shift
 $3.82028(232) \text{ fm}^2$ $\mu\text{H} / \mu\text{D}$ 2S-2P isotope shift (0.18 σ)

H/D isotope shift




electronic H/D (1S-2S): $r_d^2 - r_p^2 = 3.8207(3)_{\text{theo}} \text{ fm}^2$

muonic H/D (2S-2P): $r_d^2 - r_p^2 = 3.8200(7)_{\text{exp}}(30)_{\text{theo}} \text{ fm}^2$


→ Best bound on 5th force

K. Blaum, S. Sturm:
tomorrow

B. Ohayon (QUARTET)
μLi ... μNe
Wednesday



HFS in muonic D



PHYSICAL REVIEW A **98**, 062513 (2018)

Nuclear-structure corrections to the hyperfine splitting in muonic deuterium

Marcin Kalinowski* and Krzysztof Pachucki†

Faculty of Physics, University of Warsaw, Pasteura 5, 02-093 Warsaw, Poland

Vladimir A. Yerokhin






Center for Advanced Studies, Peter the Great St. Petersburg Polytechnic University, 195251 St. Petersburg, Russia




(Received 15 October 2018; revised manuscript received 7 November 2018; published 17 December 2018)

Nuclear structure corrections of orders $Z\alpha E_F$ and $(Z\alpha)^2 E_F$ are calculated for the hyperfine splitting of the muonic deuterium. The obtained results disagree with previous calculations and lead to a 5σ disagreement with the current experimental value of the $2S$ hyperfine splitting in muonic deuterium.

5 σ disagreement between theory and experiment !!!





Muonic Helium-4



Krauth et al. (CREMA), Nature (2021)

muonic ^4He ions



$2P_{3/2} : \pm 17 \text{ GHz}$

$2P_{1/2} : \pm 15 \text{ GHz}$

$$R(^4\text{He}) = 1.67854 (13)_{\text{exp}} (120)_{\text{theo}} \text{ fm}$$


$$(120)_{\text{theo}} = (112)_{2\text{PE}} (46)_{3\text{PE}}$$

2-photon exchange: Bacca group

3-photon exchange: our educated guess based on Pachucki et al.

Exp: Krauth, RP et al. (CREMA Coll.) Nature 589, 527 (2021) Theory updated in Pachucki, RP et al., arXiv 2212.13782

muonic ^4He ions



$$R(^4\text{He}) = 1.67854 (13)_{\text{exp}} (120)_{\text{theo}} \text{ fm}$$



$$(120)_{\text{theo}} = (112)_{2\text{PE}} (46)_{3\text{PE}}$$

2-photon exchange: Bacca group




3-photon exchange: our educated guess based on Pachucki et al.

Exp: Krauth, RP et al. (CREMA Coll.) Nature 589, 527 (2021) Theory updated in Pachucki, RP et al., arXiv 2212.13782

muonic ^4He ions




muonic ^4He ions





$2P_{1/2} : \pm 15 \text{ GHz}$

Muonic Helium-3



Schuhmann et al. (CREMA), Science (2025)
arXiv 2305.11679


muonic ^3He ions



exp: each line has ± 20 GHz(stat) ± 1 GHz (syst)

$$R(^3\text{He}) = 1.97007 (12)_{\text{exp}} (93)_{\text{theo}} \text{ fm}$$

$$\begin{aligned} \text{theo} = & \pm 0.00076 \text{ fm} & 2\text{PE} \\ & \pm 0.00052 \text{ fm} & 3\text{PE} \\ & \pm 0.00001 \text{ fm} & R^2 \text{ coeff.} \\ & \pm 0.00002 \text{ fm} & \text{QED} \end{aligned}$$




CREMA Coll., arXiv 2305.11679

ECT*, 28.7.2025

Randolf Pohl, JGU Mainz

Muonic Helium-3



e-scattering world average:

Sick 2014: PRC 90, 064002 (2014)

nuclear theory:


LENPIC: Maris et al., PRC 106, 064002 (2022)

Nevo Dinur et al., PRC 99, 034004 (2019)

Piarulli et al., PRC 87, 014006 (2013)

CREMA Coll., Science (2025)


Helium-3 – Helium-4 Isotope Shift



CREMA Coll., arXiv 2305.11679

Huang: PRA 101, 062507 (2020)
Rengelink: Nature Physics 14, 1132 (2018)
Zheng: PRL 119, 263002 (2017)
van Rooij: Science 333, 196 (2011)
Cancio Pastor: PRL 108, 143001 (2012)
Shiner: PRL 74, 3553 (1995)

Helium-3 – Helium-4 Isotope Shift




2nd order
hyperfine mixing

Qi et al, 2024
Pachucki et al. 2024

CREMA Coll., Science (2025)

Huang: PRA 101, 062507 (2020)
Rengelink: Nature Physics 14, 1132 (2018)
Zheng: PRL 119, 263002 (2017)
van Rooij: Science 333, 196 (2011)
Cancio Pastor: PRL 108, 143001 (2012)
Shiner: PRL 74, 3553 (1995)

Helium-3 – Helium-4 Isotope Shift




Huang: PRA 101, 062507 (2020)
 Rengelink: Nature Physics 14, 1132 (2018)
 Zheng: PRL 119, 263002 (2017)
 van Rooij: Science 333, 196 (2011)
 Cancio Pastor: PRL 108, 143001 (2012)
 Shiner: PRL 74, 3553 (1995)

Intermediate conclusions

Muonic atoms / ions provide:

- **~10x more accurate charge radii**, when combined with **calculated polarizability**

	${}^3\text{He}$ 1.9701* (10) 1.9730 (160)	${}^4\text{He}$ 1.6786 (12) 1.6810 (40)
${}^1\text{H}$ 0.8406 (4) 0.8751 (61)	${}^2\text{D}$ 2.1279 (2) 2.1413 (25)	${}^3\text{T}$ 1.7550 (860)




The New York Times

ECT*, 28.7.2025

Randolf Pohl, JGU Mainz

The future


5% ^3T




1S – 2S in (ordinary) atomic tritium




Tritium 1S-2S




**Triton-Radius EXperiment
Mainz**




Tritium 1S-2S




Triton-Radius EXperiment
inz





Tritium 1S-2S



Triton-Radius EXperiment
inz




Trapping and spectroscopy




- cryogenic H nozzle (4.2K)
- magnetic quadrupole guide
- Li MOT -> cold buffer gas
- magnetic trapping of H/D/T

Simulated trapping efficiency



Triton charge radius from Tritium 1S-2S




**Triton-Radius Experiment
Mainz**

	^3He 1.9679* (14) 1.9730 (160)	^4He 1.6782 (8) 1.6810 (40)
^1H 0.8409 (4) 0.8751 (61)	^2D 2.1279 (2) 2.1413 (25)	^3T 1.7xxx (200) 1.7550 (860)

**4x better radius
with 100 kHz measurement**

(vs. 0.01 kHz for H, D)


- Optogalvanic spectroscopy in a cell
- Syst. extrapolation w/ H,D
- Tritium confined.



staged approach

Hydrogen/Tritium Laser Spectroscopy in RF Discharge Cell

Spectroscopy Setup




optogalvanic detection → laser-induced impedance change of the plasma

- monitoring reflected (or forward) RF power via directional coupler (shown above)
- pickup-coil around plasma tube


- + avoid optical detection within the fluorescence background of the discharge glow
- + containment of radioactive tritium samples in a compact sealed glass cell
- large systematic effects expected due to electric fields and collision processes

Hydrogen/Tritium Laser Spectroscopy in RF Discharge Cell


Spectroscopy Setup



Resonator design inspired by [Tate, *Investigations of simple atomic systems by laser spectroscopy*, Phd thesis, Oxford (1987)]



▲ discharge tube (reduced power)






overview ▲



Hydrogen/Tritium Laser Spectroscopy in RF Discharge Cell

Balmer- β Transition / Overview


optogalvanic spectroscopy (OGS)




absorption spectroscopy



Towards trapping and spectroscopy




Triton-Radii EXperiment
Mainz




Towards H Beam Characterization

- Hydrogen beam dissociation
- Cryogenic nozzle design and atom beam shape
- Quadrupole cut-off velocity



Towards H Beam Characterization






Towards H Beam Characterization





Time-of-Flight velocity measurement
using Argon at Room Temperature


Atomic Lithium



2D MOT as cold Li beam source




2D MOT for cold Li (old)



2D-MOT design: Tiecke, Walraven et al, PRA 80, 013409 (2009)

Vacuum Setup / Beamline



Our 1st spectroscopy on cold Li





Fig.1: Fluorescence spectrum with a probe beam power μW measured over 1 s.

Next up: AFR


- Use of Active fiber-based Retroreflector:
→ Reduction of Doppler effect




[A. Beyer, "Active fiber-based retroreflector providing phase-retracing anti-parallel laser beams for precision spectroscopy", 2016]

[V. Wirthl, "Improved active fiber-based retroreflector with intensity stabilization and a polarization monitor for the near UV", 2022]


AFR: Actively stabilized Fiber-based Retro-Reflector



AFR: Actively stabilized Fiber-based Retro-Reflector



2D MOT + spectroscopy (AFR)



2D-MOT design: Tiecke, Walraven et al, PRA 80, 013409 (2009)

Thanks a lot
for your attention



Thanks a lot
for your attention




Welcome to ECT* Trento


		Monday	Tuesday	Wednesday	Thursday	Friday
9:40 – 10:20	30 + 10	van Kolck	Ohayon	Bacca	Gasparian	Craig
10:20 – 11:00	30 + 10	Pohl	Peset	Drissi	Ouf	Potvliege
11:00 – 11:30	Coffee					
11:30 – 12:10	30 + 10	Gorshteyn	Harman	Pastore	Pitelis	Feng
12:10 – 12:50	30 + 10	Fleischmann	Nörtershäuser	Oreshkina	Lensky	Closing
13:00 – 15:00	Lunch					
15:00 – 15:40	30 + 10	Kreuzberger	Weber	Excursion	Ji	
15:40 – 16:20	30 + 10	Godhino	Rathi		Egert	
16:20 – 16:50		Coffee				
16:50 – 17:30	30 + 10	Baptista	Cocolios			
17:30 – 18:10	30 + 10	Heiße (Zoom)	Deseyn			
18:10 – 18:30	Discussions					

Hyperfine structure in muonic H


CREMA-3 / HyperMu at PSI
(R16.02)



The sky in hydrogen



Hyperfine structure in H / μp



The **21 cm line** in hydrogen (1S hyperfine splitting) has been **measured** to **12 digits** (0.001 Hz) in **1971**:

$$\nu_{\text{exp}} = 1\,420\,405.751\,766\,7 \pm 0.000\,001 \text{ kHz}$$

Essen et al., Nature 229, 110 (1971)

QED test is limited to **6 digits** (800 Hz) because of **proton structure** effects:

$$\nu_{\text{theo}} = 1\,420\,403.1 \pm 0.6_{\text{proton size}} \pm 0.4_{\text{polarizability}} \text{ kHz}$$

Eides et al., Springer Tracts 222, 217 (2007)

Proton Zemach radius

HFS depends on “Zemach” radius:

$$\Delta E = -2(Z\alpha)m\langle r \rangle_{(2)}E_F$$


$$\langle r \rangle_{(2)} = \int d^3r d^3r' \rho_E(\mathbf{r}) \rho_M(\mathbf{r}') |\mathbf{r} - \mathbf{r}'|$$

Zemach, Phys. Rev. 104, 1771 (1956)

$$\Delta E = \frac{8(Z\alpha)m}{\pi n^3} E_F \int_0^\infty \frac{dk}{k^2} \left[\frac{G_E(-k^2) G_M(-k^2)}{1 + \kappa} \right]$$

Form factors and momentum space

From charge to magnetic properties




2S-2P = Lamb shift

is sensitive to CHARGE radius

1S-HFS = Hyperfine splitting


is sensitive to ZEMACH radius

Proton Zemach radius from μp



μp 2013: Antognini et al. (CREMA Coll.), Science 339, 417 (2013)

Proton Zemach radius from μp




PSI Exp. R-16-02: Antognini, RP et al. (CREMA-3 / HyperMu)

see e.g. Schmidt, RP et al., J. Phys. Conf. Ser 1138, 012010 (2018); arXiv 1808.07240

also: FAMU @ RIKEN/RAL, and a Collaboration at J-PARC

Backup


Electron scattering



Vanderhaeghen, Walcher: 1008.4225

Mainz MAMI data 2010

Correlation between R_∞ and R_p / R_d



$$E(n, \ell, j)/h = -\frac{cR_\infty}{n^2} \frac{m_{\text{red}}}{m_e} + \frac{E_{NS}}{n^3} \delta_{\ell 0} + \Delta(n, \ell, j). \quad (7)$$

$$\nu(1S - 2S) \approx \frac{3}{4} R_\infty - \frac{7}{8} E_{NS}$$

$$10^{-15} = 10 \text{ Hz} \quad 10^{-12} = 20 \text{ kHz}$$

The source of the 98.91% correlation of R_∞ and R_p

[Pohl et al., Metrologia 54, L1 (2017)]

Precision Physics of Simple Atoms and Constraints on a Light Boson with Ultraweak Coupling

S. G. Karshenboim*

D. I. Mendelev Institute for Metrology, St. Petersburg, 190005, Russia

and Max-Planck-Institut für Quantenoptik, Garching, 85748, Germany

(Received 12 April 2010; published 4 June 2010)





FIG. 1. Constraints on a long-range spin-independent interaction from hydrogen spectroscopy and $g_e - 2$, including a con-

modified Coulomb potential:

$$-\frac{\alpha}{r} \rightarrow -\frac{\alpha_{\text{eff}}(r)}{r}$$

$$= \frac{\alpha + \alpha' \exp(-m_X r)}{r}$$

Hydrogen 2S-6P: which contributions are being tested



Hydrogen $2S_{1/2}-6P_{1/2}$ (Hz)	
Dirac (with $m_e \rightarrow m_{\text{red}}$)	730 691 021 696 054
Rel. nuclear recoil	1 129 173
Radiative recoil	1540
1-loop QED	
self-energy	-1 071 679 859
vacuum-polarization	26 853 088
$\mu^+ \mu^-$ vacuum-pol.	634
hadronic vacuum-pol.	425
2-loop QED	-90 477
3-loop QED	-236
Finite nuclear size	
$\propto \alpha^4$	-138 394
$\propto \alpha^5$	5
$\propto \alpha^6$	-74
Nuclear polarizability	
$\propto \alpha^5$	8
$\propto \alpha^6$	-49
Nuclear self-energy	-584
Total	730 689 977 771 255
Theory uncertainty	199

Our 2S-6P meas. uncert.:

490 Hz

Start seeing muons and
hadrons in vacuum

Start seeing 3-loop
bound-state vacuum effects


Start seeing
Nuclear self-energy

CREMA: Laser spectroscopy of light muonic atoms




Radii of $Z=3$ 10

Xray spectroscopy of muonic atoms



X-ray spectroscopy





low-Z muonic atoms:
low X-ray energy → difficult!

muonic Li: 18 keV

$$E_n = -\frac{R_\infty}{n^2} + \frac{C_{FS}}{n^3} \langle r^2 \rangle \delta_{l0} + \Delta(n, l, j)$$

QUARTET Goals



10x improved nuclear charge radii → challenging nuclear few-body calculations

Current knowledge on radii of the lightest nuclei:



$Z=1,2$: muonic atom laser spectroscopy

$Z>3$: mostly e-scattering

$Z=6$: some muonic X-rays (crystal spectrometer)

$Z>8$: muonic X-rays (Ge detectors)

QUARTET Goals



QUARTET, muX, ReferenceRadii:

see Talk by Frederik Wauters

tomorrow

10x improved nuclear charge radii → challenging nuclear few-body calculations

Current knowledge on radii of the lightest nuclei:



$Z=1,2$: muonic atom laser spectroscopy

$Z>3$: mostly e-scattering

$Z=6$: some muonic X-rays (crystal spectrometer)

$Z>8$: muonic X-rays (Ge detectors)

Quantum Interference in 2S-nP





see
 Horbatsch, Hessels, PRA 82, 052519 (2010);
 PRA 84, 032508 ('11); PRA 86 040501 ('12)
 Sansonetti et al., PRL 107, 021001 (2011)
 Brown et al., PRA 87, 032504 (2013)

$$P(\omega) \propto \left| \frac{(\vec{d}_1 \vec{E}_0) \vec{d}_1}{\omega_1 - \omega_L + i\gamma_1/2} + \frac{(\vec{d}_2 \vec{E}_0) \vec{d}_2 e^{i\Delta\Phi}}{\omega_2 - \omega_L + i\gamma_2/2} \right|^2$$

= Lorentzian(1) + Lorentzian(2)
 + cross-term (QI)

Fitting this with 2 Lorentzians creates
line shifts

Studying Quantum Interference in 2S-4P




$$P(\omega) \propto \left| \frac{(\vec{d}_1 \vec{E}_0) \vec{d}_1}{\omega_1 - \omega_L + i\gamma_1/2} + \frac{(\vec{d}_2 \vec{E}_0) \vec{d}_2 e^{i\Delta\Phi}}{\omega_2 - \omega_L + i\gamma_2/2} \right|^2$$

= Lorentzian(1) + Lorentzian(2)

+ cross-term (QI)

Beyer, Maisenbacher, RP et al, Science 358, 79 (2017)

Studying Quantum Interference in 2S-4P



Beyer, Maisenbacher, RP et al, Science 358, 79 (2017)