Beta Decays at a loop level in SMEFT

Maria Dawid

Based on
M.D, V. Cirigliano, W. Dekens
ArXiv: (2402.06723)

Beta decay



The Cabibbo-Kobayashi-Maskawa matrix

$$V_{\text{CKM}} V^{\dagger}_{\text{CKM}} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} V^*_{ud} & V^*_{cd} & V^*_{td} \\ V^*_{us} & V^*_{cs} & V^*_{ts} \\ V^*_{ub} & V^*_{cb} & V^*_{tb} \end{pmatrix} = 1 \ ,$$



$$\mathscr{L}_{EW} \subset \frac{g}{\sqrt{2}} W^+_\mu \, \bar{d}_{i_L} \, V_{ij} \gamma^\mu u_{j_L}$$

Unitarity

$$\Delta_{CKM} \equiv |V_{ud}|^2 + |V_{us}|^2 - 1 \approx 0$$

Global fit

$$\Delta_{CKM} = -\,1.48(53) \times 10^{-3}$$

$$V_{ud} = 0.973670(25)$$

$$V_{us} = 0.22422(36)$$

Discrepancy with SM

- Underestimated SM uncertainties?
- BSM physics?

The CKM unitarity test



The Cabibbo-Kobayashi-Maskawa matrix



Scrutinizing CKM unitarity with a new measurement of the $K_{\mu 3}/K_{\mu 2}$ branching fraction
V. Cirigliano et al. (arXiv: 2208.11707)

Unitarity

$$\Delta_{CKM} \equiv |V_{ud}|^2 + |V_{us}|^2 - 1 \approx 0$$

Global fit

$$\Delta_{CKM} = -1.48(53) \times 10^{-3}$$

$$V_{ud} = 0.973670(25)$$

$$V_{us} = 0.22422(36)$$

Discrepancy with SM

- Underestimated SM uncertainties?
- BSM physics?

The CKM unitarity test



The Cabibbo-Kobayashi-Maskawa matrix

Probing physics beyond Standard Model

Work on three different fronts:

Enhancing the accuracy of experiments

Improving precision of SM calculations

Advancing theory for BSM physics

Standard Model Effective Field Theory

Unitarity

$$\Delta_{CKM} \equiv |V_{ud}|^2 + |V_{us}|^2 - 1 \approx 0$$

Global fit

$$\Delta_{CKM} = -1.48(53) \times 10^{-3}$$

$$V_{ud} = 0.973670(25)$$

$$V_{us} = 0.22422(36)$$

Discrepancy with SM

- Underestimated SM uncertainties?
- BSM physics?

The CKM unitarity test



The Cabibbo–Kobayashi–Maskawa matrix

Probing physics beyond Standard Model

Work on three different fronts:

Enhancing the accuracy of experiments

Improving precision of SM calculations

Advancing theory for BSM physics

Standard Model Effective Field Theory

One-loop corrections to two-body beta decay
with: M. d'Souza, V. Cirigliano, W. Dekens



Example

Tree-level three-body beta decay:
with: G. Chambers-Wall, E. Mereghetti, S. Pastore



Example

Effective Field Theories

To connect UV physics to beta decays, use EFT



Effective Field Theories

To connect UV physics to beta decays, use EFT



This talk

Standard Model EFT

The EFT constructed with Standard Model fields & symmetries

$$\mathscr{L}_{SMEFT} = \mathscr{L}_{SM} + \frac{1}{v} C^{(5)} Q^{(5)} + \frac{1}{v^2} \sum_k C_k^{(6)} Q_k^{(6)} + \mathcal{O}(\frac{1}{v^3}) + \dots$$

The Standard Model is a part of the leading order of SMEFT

With $C_k \sim v^2/\Lambda^2$

C - Wilson coefficients

Q - local interaction terms (operators)

Example of EFT: Fermi Theory



$$\frac{-g_{\mu\nu} + q_\mu q_\nu / M_W^2}{q^2 - M_W^2} \xrightarrow{q^2 << M_W^2} \frac{g_{\mu\nu}}{M_W^2}$$

Standard Model EFT

The EFT constructed with Standard Model fields & symmetries

$$\mathscr{L}_{SMEFT} = \mathscr{L}_{SM} + \frac{1}{v} C^{(5)} Q^{(5)} + \frac{1}{v^2} \sum_k C_k^{(6)} Q_k^{(6)} + \mathcal{O}(\frac{1}{v^3}) + \dots$$

6-dim operators

With $C_k \sim v^2/\Lambda^2$

No fermions







Two fermions







Four fermions



Standard Model EFT

The EFT constructed with Standard Model fields & symmetries

$$\mathscr{L}_{SMEFT} = \mathscr{L}_{SM} + \frac{1}{v} C^{(5)} Q^{(5)} + \frac{1}{v^2} \sum_k C_k^{(6)} Q_k^{(6)} + \mathcal{O}(\frac{1}{v^3}) + \dots$$

The Standard Model is a part of the leading order of SMEFT

With $C_k^i \sim v^{i-4}/\Lambda^{i-4}$

C - Wilson coefficients

Q - local interaction terms (operators)

Operators that contain particles directly involved in a process

SMEFT contributions to beta decay are well-known at a tree level

What about quantum fluctuations?

Goal of the work: identify leading contributions to beta decay at a one-loop level.

β-decay on loop level



β-decay on loop level



GeV scale effective Lagrangian



$$d \to u\, l^-\, \bar{\nu}_l$$

From muon decay

$$\epsilon_L = \epsilon_L^\beta - \epsilon_L^\mu$$

$\mathscr{L}_{LEFT} = -\frac{G_F^0 V_{ud}}{\sqrt{2}} [(1+\epsilon_L)\, \bar{e}\, \gamma_\mu (1-\gamma_5) \nu_l\, \bar{u}\gamma^\mu (1-\gamma_5) d$	left quarks
$+ \epsilon_R\, \bar{e}\gamma_\mu (1-\gamma_5) \nu_l\, \bar{u}\gamma^\mu (1+\gamma_5) d$	right quarks
$+ \epsilon_S\, \bar{e}\, (1-\gamma_5) \nu_l\, \bar{u} d$	scalar
$- \epsilon_P\, \bar{e} (1-\gamma_5) \nu_l\, \bar{u}\gamma_5 d$	pseudo-scalar
$+ \epsilon_T\, \bar{e}\sigma_{\mu\nu}\, (1-\gamma_5) \nu_l\, \bar{u}\sigma^{\mu\nu}\, (1-\gamma_5) d\,]$	tensor

ϵ_L^μ SMEFT correction to GF as extracted from muon decay

Anomalies in global SMEFT analyses. A case study of first-row CKM unitarity
V. Cirigliano, W. Dekens, J. de Vries, E. Mereghetti, T. Tong
(arXiv:2311.00021)

Global Fit

flavor-assumption-independent analysis

- Drell-Yan collider processes ('C')

- Low-energy charged current processes ('L')

  – Neutron and nuclear β decays
  – Kaon and pion decays ($\Gamma(K \to \mu\nu_\mu)$,

$$R_\pi = \Gamma(\pi \to e\nu_e)/\Gamma(\pi \to \mu\nu_\mu),$$
$$R_K = \Gamma(K \to e\nu_e)/\Gamma(K \to \mu\nu_\mu),$$
$$R_\mu = \Gamma(K \to \mu\nu_\mu)/\Gamma(\pi \to \mu\nu_\mu)\ )$$

- electroweak precision observables ('EW')

  – observables measured at the Z pole (decay widths, asymmetries, hadronic cross section obtained from $e + e^- \to Z \to \bar{q}q$, W mass)

→ put constraints on ϵ_i's $\qquad \sim v^2/\Lambda^2$

$$\epsilon_L^{(\mu)},\ \ \epsilon_L^{(c)},\ \ \epsilon_L^{(v)} \approx -\,8.3 \pm 2.5 \times 10^{-4}$$

$\to$ 20 TeV

Other fit:
A. Falkowski, M. González-Alonso, O. Naviliat-Cuncic
(arXiv:2010.1379))

Global Fit



- Current and projected 90% C.L. constraints on ϵ_S and ϵ_T defined at 2 GeV in the MS bar scheme
- Beta decay:
  - The current analysis includes all existing neutron and nuclear decay measurements,
  - Future projection assumes measurements of the various decay correlations with uncertainty of 0.1%
- LHC:
  - Obtained from $pp \rightarrow e + MET + X$
  - ATLAS results at $\sqrt{s} = 13$ TeV
  - The projected future LHC bounds are obtained by assuming that no events are observed at transverse mass greater than 3 TeV
- LHC cannot constrain $\epsilon_{L'}$ and ϵ_R

Isovector and flavor diagonal charges of the nucleon from 2+1+1 flavor QCD
Rajan Gupta et al. (arXiv:1812.03573)

β-decay on loop level



Beta Decay at the Tree Level



Four fermion operators

$$Q_{lq}^{(3)} = (\bar{l}_p \gamma_\mu \tau^I l_r)\,(\bar{q}_s \gamma^\mu \tau^I q_t),$$
$$Q_{ledq} = (\bar{l}_p e_r)\,(\bar{d}_s q_t),$$
$$Q_{lequ}^{(1)} = \left(\bar{l}_p^j e_r\right) \epsilon_{jk} (\bar{q}_s^k u_t),$$
$$Q_{lequ}^{(3)} = \left(\bar{l}_p^j \sigma_{\mu\nu} e_r\right) \epsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} u_t),$$



Vertex correction

$$Q_{Hud} = i\left(H^\dagger D_\mu H\right)\left(\bar{u}_p \gamma^\mu d_r\right),$$
$$Q_{Hl}^{(3)} = \left(H^\dagger i \overleftrightarrow{D}_\mu^I H\right)\left(\bar{l}_p\, \tau^I \gamma^\mu l_r\right),$$
$$Q_{Hq}^{(3)} = \left(H^\dagger i \overleftrightarrow{D}_\mu^I H\right)\left(\bar{q}_p\, \tau^I \gamma^\mu q_r\right),$$

p , r , s, t are flavor indices
here all 1

We work in the weak eigenstate basis

$$M_u = diag\,(m_u, m_c, m_t),\; M_d = diag\,(m_d, m_c, m_t)\; V^\dagger$$

$$l^i = \begin{pmatrix} \nu_L^i \\ e_L^i \end{pmatrix},\quad q^i = \begin{pmatrix} u_L^i \\ V\, d_L^i \end{pmatrix},\quad H = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix},$$

e, u, d – right-chiral fields,
$\tau - SU(2)$ generators,
D_μ - covariant derivative

Muon Decay

$\epsilon_L^{(\mu)}$ arises from the SMEFT correction to the Fermi constant extracted from muon decay and is given by

$$Q_{Hl}^{(3)} = \left(H^\dagger i \overleftrightarrow{D}_\mu^I H \right) (\bar{l}_p \, \tau^I \gamma^\mu l_r)$$

$$Q_{ll} = (\bar{l}_p \gamma_\mu l_r)(\bar{l}_s \gamma_\mu l_t)$$

flavour indices:
(1,2,2,1)



$$\epsilon_L^{(\mu)} = \left(C_{ll_{1221}} + C_{ll_{2112}} \right) - 2(C_{Hl_{11}}^{(3)} + C_{Hl_{22}}^{(3)}$$

$$l^i = \begin{pmatrix} \nu_L^i \\ e_L^i \end{pmatrix}, \quad q^i = \begin{pmatrix} u_L^i \\ V\, d_L^i \end{pmatrix}, \quad H = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix},$$

e, u, d – right-chiral fields,
$\tau - SU(2)$ generators,
D_μ - covariant derivative

Matching at the Electro-Weak Scale

$$[\mathcal{T}_{LEFT} = \mathcal{T}_{SMEFT}]_{\mu_W}$$

Vector interactions for left quarks:

- $\epsilon_L = \epsilon_L^{(v)} + \epsilon_L^{(c)} - \epsilon_L^{(\mu)}$
- $\epsilon_L^{(v)} = 2\, C_{Hl}^{(3)} + 2\frac{\left[V C_{Hq}^{(3)}\right]_{11}}{V_{ud}} + 4\frac{[V C_{Hl}^{(3)}]_{11}}{V_{ud}},$
- $\epsilon_L^{(c)} = -2\frac{\left[V C_{lq}^{(3)}\right]_{11}}{V_{ud}},$
- $\epsilon_L^{(\mu)} = \left(C_{ll_{1221}} + C_{ll_{2112}}\right) - 2(C_{Hl_{11}}^{(3)} + C_{Hl_{22}}^{(3)})$

$$\epsilon_R = -\frac{\left[V C_{Hud}^{(3)}\right]_{11}}{V_{ud}},$$

$$\epsilon_s - \epsilon_P = -2\frac{\left[V C_{ledq}^{(3)}\right]_{11}}{V_{ud}},$$

$$\epsilon_s + \epsilon_P = -2\frac{\left[V C_{lequ}^{(1)}\right]_{11}}{V_{ud}},$$

$$\epsilon_T = -\frac{C_{lequ}^{(3)}}{V_{ud}}\,.$$

$\epsilon_L^{(v)}$ - vertex correction
$\epsilon_L^{(c)}$ - contact correction
$\epsilon_L^{(\mu)}$ - correction to muon decay

$$\left[V C_{Hq}^{(3)}\right]_{11} = C_{Hq}^{(3)}{}_{11} V_{11} + C_{Hq}^{(3)}{}_{12} V_{21} + C_{Hq}^{(3)}{}_{13} V_{31}$$

Bounds from beta decay (CKM unitarity)

$$\Delta_{CKM} = 2\left|V_{ud}\right|^2 \epsilon_L^d + 2\left|V_{us}\right|^2 \epsilon_L^s$$
$$\epsilon_L^{(\mu)},\ \epsilon_L^{(c)},\ \epsilon_L^{(v)} \approx -8.3 \pm 2.5 \times 10^{-4},$$

β-decay on loop level



Mixing Procedure

- SMEFT operator



$Q^{(3)}_{Hq}$

Triplet

$$Q^{(3)}_{Hq} = (H^\dagger i\overleftrightarrow{D}^I_\mu H)(\bar{q}_p \tau^I \gamma^\mu q_r)$$

2 quarks and boson



$Q^{(3)}_{qq}{}_{1133} \rightarrow Q^{(3)}_{Hq}{}_{11}$

$$Q^{(3)}_{qq} = (\bar{q}_p \gamma_\mu \tau^I q_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$$

4 quarks

More Examples

- SMEFT operator



Contribution to $Q_{Hl}^{(3)}$ and $Q_{Hq}^{(3)}$
from $Q_{qq}^{(3)}$, $Q_{qq}^{(1)}$, $Q_{lq}^{(3)}$



Contribution to $Q_{lq}^{(3)}$
from $Q_{qq}^{(3)}$, $Q_{qq}^{(1)}$



Contribution to Q_{ll}
from Q_{ll}

Renormalization Group Equations

RGE

Tree level →
Loop level →

$$\begin{pmatrix} \dot{C}_\beta \\ \dot{C}_x \end{pmatrix} = \begin{pmatrix} \gamma_{\beta\beta} & \gamma_{\beta x} \\ \gamma_{x\beta} & \gamma_{xx} \end{pmatrix} \begin{pmatrix} C_\beta \\ C_x \end{pmatrix}$$

Initial conditions for $\Lambda = 5$ TeV

$$C_\beta(\Lambda) = 0$$

$$C_x(\Lambda) = 1$$

Mixing of operators

$$C_{\beta j}(\mu_W) = \sum_k U_{jk}(\mu_W, \Lambda) C_{xk}(\Lambda)$$

$$|\epsilon| = |\kappa_i \, C_i| \leq \bar{\epsilon}_L$$

mixing parameter from RGE's

bound from the experiment

Example of solutions of RGE's for $C_{Hq}^{(3)}$, $C_{Hl}^{(3)}$ and $C_{lq}^{(3)}$



RH, Scalar, Pseudoscalar, and Tensor Terms

This class of couplings are suppressed by y_b

Example:

$$\dot{C}_{\substack{Hud \\ 11}} \subset 4\left(C^{(1)}_{\substack{ud \\ 1331}} + \frac{4}{3} C^{(8)}_{\substack{ud \\ 1331}}\right)\left[\mathrm{Y}_u Y_d^{\dagger}\right]_{33} \xrightarrow{\text{Solving RGE}} \text{bounds of } \mathcal{O}(1)$$

Results for ϵ_i at $\mu_W = 246$ GeV in terms of C_β and C_x at $\Lambda = 5$ Tev

$$\epsilon_R = 0.950\, \boldsymbol{C_{Hud}} - 0.001\, C^{(1)}_{ud} - 0.0011\, C^{(8)}_{ud},$$
$$\epsilon_P = 0.603\, \boldsymbol{C^{(1)}_{lequ}} - 0.600\, \boldsymbol{C_{ledq}} + 0.143\, C^{(3)}_{lequ}$$
$$\epsilon_S = -0.603\, \boldsymbol{C^{(1)}_{lequ}} - 0.600 \boldsymbol{C_{ledq}} + 0.143\, C^{(3)}_{lequ}$$
$$\epsilon_T = 0.003\, C^{(1)}_{lequ} - 0465\, \boldsymbol{C^{(3)}_{lequ}}$$



Coefficienents in blue are matched to a given ϵ_i at a tree level

LH Currents

The largest contribution to ϵ_i at a loop level ($\kappa_{ij} > 10^{-2}$):

$$Q_{\beta_L}: \{Q_{ll}, Q_{lq}^{(3)}, Q_{Hl}^{(3)}, Q_{Hq}^{(3)}, Q_{qq}^{(1)}, Q_{qq}^{(3)}, Q_{H\Box}\}$$

$$Q_{qq}^{(1)} = \left(\bar{q}_p \gamma_\mu q_r\right)\left(\bar{q}_s \gamma^\mu q_t\right)$$
$$Q_{qq}^{(3)} = \left(\bar{q}_p \gamma_\mu \tau^I q_r\right)\left(\bar{q}_s \gamma^\mu \tau^I q_t\right)$$
$$Q_{H\Box} = \left(H^\dagger H\right)\Box\left(H^\dagger H\right)$$

Results for ϵ_L at $\mu_W = 246$ GeV in terms of C_β and C_x at $\Lambda = 5$ Tev

$$\begin{aligned}
\epsilon_L^{(d)} &= 0.946\, C^{(3)}_{\substack{Hq\\11}} - 1.052\, C^{(3)}_{\substack{lq\\1111}} + 1.012\, C_{\substack{ll\\1221}} - 0.965\, C^{(3)}_{\substack{Hl\\22}} \\
&+ 0.198\, C^{(3)}_{\substack{qq\\1133}} - 0.026\, C^{(3)}_{\substack{qq\\1331}} + 0.026\, C^{(1)}_{\substack{qq\\1331}} - 0.0471\, C_{\substack{ll\\1122}} - 0.100\, C^{(3)}_{\substack{lq\\2233}} ,
\end{aligned}$$

Beta decay

$$\begin{aligned}
\epsilon_L^{(s)} &= \frac{0.942}{\lambda} C^{(3)}_{\substack{Hq\\12}} - \frac{1.052}{\lambda} C^{(3)}_{\substack{lq\\1112}} + 1.010\, C_{\substack{ll\\1221}} - 0.960\, C^{(3)}_{\substack{Hl\\22}} \\
&+ \frac{0.198}{\lambda} C^{(3)}_{\substack{qq\\1233}} - \frac{0.028}{\lambda} C^{(3)}_{\substack{qq\\1332}} + \frac{0.026}{\lambda} C^{(1)}_{\substack{qq\\1332}} - 0.099\, C^{(3)}_{\substack{lq\\2233}} - 0.047\, C_{\substack{ll\\1122}} \\
&+ 0.0144\,(C^{(3)}_{\substack{Hq\\11}} + C^{(3)}_{\substack{Hq\\22}} + C^{(3)}_{\substack{Hq\\33}}) - 0.016\,(C^{(3)}_{\substack{lq\\1111}} + C^{(3)}_{\substack{lq\\1122}} + C^{(3)}_{\substack{lq\\1133}}) .
\end{aligned}$$

Kaon decay

Results

With $C_k \sim v^2/\Lambda^2$

$C_x(\Lambda = 5\,\text{TeV})$	Constraint from β decays	Strongest constraints from other processes	Process
$C^{(3)}_{qq\ 1133}$	-0.004 ± 0.0013	$-0.0073 \pm 0.006^{(1)}$ $\pm 0.00002^{(2)}$	Top production $K \to \pi\nu\bar{\nu}$
$C^{(3)}_{lq\ 2233}$	0.008 ± 0.0025	$-0.00024 \pm 0.00021^{(2)}$	B decays (R_K)
$C_{ll\ 1122}$	0.018 ± 0.005	-0.0018 ± 0.0029	Muon pair production
$C^{(1)}_{qq\ 1331}$	-0.031 ± 0.009	$-0.035 \pm 0.027^{(1)}$ $\pm 0.0009^{(2)}$	Top production Δm_K
$C^{(3)}_{qq\ 1331}$	0.03 ± 0.009	$-0.042 \pm 0.024^{(1)}$ $\pm 0.00004^{(2)}$	Top production $K \to \pi\nu\bar{\nu}$

(1) indicates that the fit was performed assuming $U(3)_l \times U(3)_e \times U(3)_d \times U(2)_u \times U(2)_q$ symmetry

(2) indicates that the constraint would disappear if one used the weak basis in which the down-type quark mass matrix M_d is diagonal.

M.D, V. Cirigliano, W. Dekens
ArXiv: (2402.06723)

Example of tree level process, $t\bar{t}$ production at the LHC



Result: Stronger constraints on BSM operators from loop-level processes than those achievable in other channels at tree level.

Results vs LHC

Transverse mass distributions for events satisfying all selection criteria in the electron channels.



Scattering amplitude $\mathcal{T} = \mathcal{T}_{SM} + \mathcal{T}_{SMEFT}$

Cross-section $\sigma = \left|\mathcal{T}_{SM}\right|^2 + 2Re\left[\mathcal{T}_{SM} \times \mathcal{T}_{SMEFT}\right] + \left|\mathcal{T}_{SMEFT}\right|^2$

Results vs LHC

Transverse mass distributions for events satisfying all selection criteria in the electron channels.



Search for a new heavy gauge boson resonance decaying into a lepton and missing transverse momentum in 36 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS experiment

Results

M.D, V. Cirigliano, W. Dekens
ArXiv: (2402.06723)

$$C \sim \frac{v^2}{\Lambda^2}$$



$\Lambda \sim 8$ TeV

Result: Beta decay at loop level can probe higher energies (that correspond to masses of BSM particles) than other processes, e.g., from LHC

β-decay on loop level



Beyond the SM: a model above the high-energy scale

Assumption of BSM above the Λ scale

1. $\mathscr{L}_{BSM}$ is invariant under linearly-realized $SU(3)_C \times SU(2)_L \times U(1)_Y$.
2. $\mathscr{L}_{BSM}$ containts only particles of spin ≤ 1.
3. Adding one new particle at a time.

Q_x operator	Heavy field						
$Q_{qq}^{(3)}$ Irrep	ω_1 $(3,1)_{-\frac{1}{3}}$	ζ $(3,3)_{-\frac{1}{3}}$	Ω_1 $(6,1)_{\frac{1}{3}}$	Υ^* $(6,3)_{\frac{1}{3}}$	$\mathcal{W}$ $(1,1)_0$	$\mathcal{G}$, $(8,1)_0$	$\mathcal{H}^*$ $(8,3)_0$
$Q_{qq}^{(1)}$ Irrep	ω_1 $(3,1)_{-\frac{1}{3}}$	ζ $(3,3)_{-\frac{1}{3}}$	Ω_1 $(6,1)_{\frac{1}{3}}$	Υ^* $(6,3)_{\frac{1}{3}}$	$\mathcal{B}$ $(1,3)_0$	$\mathcal{G}$, $(8,1)_0$	$\mathcal{H}^*$ $(8,3)_0$
$Q_{lq}^{(3)}$ Irrep	ω_1 $(3,1)_{-\frac{1}{3}}$	ζ $(3,3)_{-\frac{1}{3}}$	$\mathcal{W}$ $(1,3)_0$	$\mathcal{U}_2$ $(3,1)_0$	$\mathcal{X}$ $(3,3)_{\frac{2}{3}}$		
Q_{ll} Irrep	$\mathcal{S}_1$ $(1,1)_1$	Ξ $(1,3)_1$	β $(1,1)_0$	$\mathcal{W}$ $(1,3)_0$			

J. De Blas, J.C. Criado at al. (1711.10391)

Summary

- The CKM unitarity test is important for constraining ϵ_L and ϵ_R
- Beta decay probes BSM physics at a tree-level at ~ 20 TeV (needs experimental and theoretical scrutiny!)
- We put constraints on New Physics operators that include **top quarks** at ~ 8 TeV at a loop-level, which exceeds the LHC results!
- We connect these operators to the BSM model

FLAVOR-CHANGING NEUTRAL CURRENT OPERATORS

- At a loop level ($\Delta F = 2$):
  $\dot{C}_\beta = \gamma_{\beta x} C_x,$
  $\dot{C}_{FCNC} = \gamma_{FCNC}\, C_x$
- Rotation to the mass basis

The strongest constraints arise from the rare

$\Delta S = 1$ proces $K \to \pi\nu\bar{\nu}$ ⟵ $[\mathrm{L}^{\nu d}_{VLL}] < 0.97 \times 10^{-5}$ TeV

$$\upsilon^2[\mathrm{L}^{\nu d}_{VLL}] = \tilde{C}_{ll12} + \lambda\left(\tilde{C}_{ll11} - \tilde{C}_{ll22}\right) +$$
$$\tilde{C}_{llij} = \underset{llij}{C^{(1)}_{lq}} - \underset{llij}{C^{(3)}_{lq}} + \delta_{ll}\underset{ij}{C^{(1)}_{Hq}} + \delta_{ll}\underset{ij}{C^{(3)}_{Hq}}$$

⟹

$C^{(1)}_{lq\ 1133} < 2.0 \times 10^{-5}$, $C^{(1)}_{qq\ 1331} < 0.57 \times 10^{-3}$

$C^{(3)}_{lq\ 1133} < 2.0 \times 10^{-5}$, $C^{(3)}_{qq\ 1331} < 0.37 \times 10^{-4}$

With $C_k \sim v^2/\Lambda^2$