

# Heavy-quark transport coefficients Sensitivity to hadronization description

Hendrik van Hees

Goethe University Frankfurt

April 27, 2021



# Outline

- 1 Bulk-medium evolution
- 2 Heavy-quark interactions in the sQGP
- 3 Non-perturbative HQ interactions
  - Resonance model for HQ-q Scattering
  - T-matrix approach with lQCD potentials
- 4 Hadronization and HQs at FAIR


# Bulk-medium evolution (e.g., LHC)

- transport (UrQMD) + hydro (SHASTA) hybrid model
- nuclei: Woods-Saxon profile; binary interactions
- string excitation, fragmentation (PYTHIA)
- after  $t_{\text{start}} = 0.5 \text{ fm}/c$ : particle distributions, momenta, energy density from UrQMD → via Gaussian smearing to **initial conditions for hydro**
- during hydro evolution: **HG EoS or chiral DE-EoS**
- switch back to transport: **Cooper-Frye freezeout** at constant  $\tau$ -hypersurfaces for  $\epsilon \lesssim 5\epsilon_0 \simeq 730 \text{ MeV/fm}^3$

[H. Petersen, Phys.Rev. C **84** (2011) 034912; arXiv:1206.3371 [nucl-th]]


# Bulk-medium evolution

- Pb-Pb collisions  $\sqrt{s_{NN}} = 2.76$  TeV
- particle multiplicities at mid rapidity  $|\eta| < 0.5$




# Bulk-medium evolution

- Pb-Pb collisions  $\sqrt{s_{NN}} = 2.76$  TeV
- $p_T$  distribution of charged hadrons



# Bulk-medium evolution

- Pb-Pb collisions  $\sqrt{s_{NN}} = 2.76$  TeV
- $v_2$  of charged hadrons




# Bulk-medium evolution

- Pb-Pb collisions  $\sqrt{s_{NN}} = 2.76$  TeV
- $\epsilon_2$  and  $\epsilon_3$  distributions

[H. Petersen, G.-Y. Qin, S. A. Bass, B. Müller, Phys. Rev. C **82**, 041901 (2010); arXiv:1008.0625 [nucl-th]]

$$\epsilon_n = \frac{\sqrt{\langle r^n \cos(n\phi) \rangle^2 + \langle r^n \sin(n\phi) \rangle^2}}{\langle r^n \rangle}$$



# Relativistic Langevin process

- Langevin process: friction force + Gaussian random force
- in the (local) rest frame of the heat bath

$$d\vec{x} = \frac{\vec{p}}{E_p} dt,$$

$$d\vec{p} = -A \vec{p} dt + \sqrt{2dt} [\sqrt{B_0} P_{\perp} + \sqrt{B_1} P_{\parallel}] \vec{w}$$

- $\vec{w}$ : normal-distributed random variable
- $A$ : friction (drag) coefficient
- $B_{0,1}$ : diffusion coefficients
- Einstein dissipation-fluctuation relation  $B_1 = E_p T A$ .
- flow via Lorentz boosts between “heat-bath frame” and “lab frame”
- $A$  and  $B_0$  from microscopic models for  $qQ, gQ$  scattering
- medium: UrQMD → hydro → UrQMD vs. UrQMD/coarse-graining

[R. Rapp, HvH, R. C. Hwa and X. N. Wang (eds.), Quark-Gluon Plasma Vol. IV, World Scientific (2010), arXiv: 0903.1096 [hep-ph]; M. He, HvH, P. B. Gossiaux, R. J.

Fries, R. Rapp, Phys. Rev. E 88, 032138 (2013)]

# Free Lagrangian: Particle Content

- Chiral symmetry  $SU_V(2) \otimes SU_A(2)$  in light-quark sector of QCD

$$\mathcal{L}_D^{(0)} = \sum_{i=1}^2 [(\partial_\mu \Phi_i^\dagger)(\partial^\mu \Phi_i) - m_D^2 \Phi_i^\dagger \Phi_i] + \text{massive (pseudo-)vectors } D^*$$

- $\Phi_i$ : two doublets: **pseudo-scalar**  $\sim \begin{pmatrix} \bar{D}^0 \\ D^- \end{pmatrix}$  and **scalar**
- $\Phi_i^*$ : two doublets: **vector**  $\sim \begin{pmatrix} \bar{D}^{0*} \\ D^{-*} \end{pmatrix}$  and **pseudo-vector**

$$\mathcal{L}_{qc}^{(0)} = \bar{q} i \not{\partial} q + \bar{c} (i \not{\partial} - m_c) c$$

- $q$ : light-quark doublet  $\sim \begin{pmatrix} u \\ d \end{pmatrix}$
- $c$ : singlet

# Interactions


- Interactions determined by **chiral** symmetry
- For transversality of vector mesons:  
**heavy-quark effective theory vertices**

$$\begin{aligned}\mathcal{L}_{\text{int}} = & -G_S \left( \bar{q} \frac{1+\not{v}}{2} \Phi_1 c_v + \bar{q} \frac{1+\not{v}}{2} i \gamma^5 \Phi_2 c_v + h.c. \right) \\ & - G_V \left( \bar{q} \frac{1+\not{v}}{2} \gamma^\mu \Phi_{1\mu}^* c_v + \bar{q} \frac{1+\not{v}}{2} i \gamma^\mu \gamma^5 \Phi_{2\mu}^* c_v + h.c. \right)\end{aligned}$$


- $v$ : four velocity of heavy quark
- in **HQET**: spin symmetry  $\Rightarrow G_S = G_V$

# Non-perturbative interactions: Resonance Scattering

- General idea: Survival of  $D$ - and  $B$ -meson like **resonances** above  $T_c$
- model based on chiral symmetry (light quarks) HQ-effective theory
- elastic heavy-light-(anti-)quark scattering**



- $D$ - and  $B$ -meson like resonances in **sQGP**




- parameters

- $m_D = 2 \text{ GeV}, \Gamma_D = 0.4 \dots 0.75 \text{ GeV}$
- $m_B = 5 \text{ GeV}, \Gamma_B = 0.4 \dots 0.75 \text{ GeV}$

[HvH, R. Rapp, Phys. Rev. C 71, 034907 (2005); HvH, V. Greco, R. Rapp, Phys. Rev. C 73, 034913 (2006) ]



# Cross sections



- total pQCD and resonance cross sections: comparable in size
- BUT pQCD forward peaked  $\leftrightarrow$  resonance isotropic
- resonance scattering more effective for friction and diffusion

# Transport coefficients: pQCD vs. resonance scattering



- three-momentum dependence



- resonance contributions factor  $\sim 2 \dots 3$  higher than pQCD!


# Transport coefficients: pQCD vs. resonance scattering

- Temperature dependence



# T-matrix

- Brueckner many-body approach for elastic  $Qq, Q\bar{q}$  scattering




- $V$ : static  $q\bar{q}$  potential from lattice QCD ( $F$  and  $U$ )
- reduction scheme: 4D Bethe-Salpeter → 3D Lipmann-Schwinger
- $S$ - and  $P$  waves
- Relation to invariant **matrix elements**

$$\sum_q |\mathcal{M}(s)|^2 \propto \sum_q d_a \left( |\textcolor{red}{T}_{a,l=0}(s)|^2 + 3|\textcolor{red}{T}_{a,l=1}(s)|^2 \cos \theta_{\text{cm}} \right)$$

[HvH, M. Mannarelli, V. Greco, R. Rapp, Phys. Rev. Lett. **100**, 192301 (2008)]

# Static heavy-quark potentials from lattice QCD



- color-singlet free energy from lattice → internal energy


$$U_1(r, T) = F_1(r, T) - T \frac{\partial F_1(r, T)}{\partial T},$$

$$V_1(r, T) = U_1(r, T) - U_1(r \rightarrow \infty, T)$$

- Casimir scaling of Coulomb part for other color channels;  
confining part color blind [[F Riek, R. Rapp, Phys. Rev. C 82, 035201 \(2010\)](#)].


$$V_3 = \frac{1}{2} V_1, \quad V_6 = -\frac{1}{4} V_1, \quad V_8 = -\frac{1}{8} V_1$$

# T-matrix results



- resonance formation at lower temperatures  $T \simeq T_c$
- melting of resonances at higher  $T$
- model-independent assessment of elastic  $Qq, Q\bar{q}$  scattering!

# Transport coefficients



- $T$ -matrix resonance-scattering coefficients: **decrease** with  $T$
- from **non-pert.** interactions reach  $A_{\text{non-pert}} \simeq 1/(7 \text{ fm}/c) \simeq 4 A_{\text{pQCD}}$
- results for **free-energy potential**,  $F$  considerably smaller

# Hadronization

- Coalescence [Greco, Ko, Rapp, Phys. Lett. B 595, 202 (2004)]

$$P_{\text{coa}} = \exp \left\{ \left[ (\Delta p^0)^2 - \sum_{i=1}^3 (\Delta p^i)^2 - (\Delta_m)^2 \right] \sigma^2 \right\}, \quad \sigma^2 = \frac{8}{3} r_{\text{D}}^2$$

- Petersen fragmentation ( $z$ : momentum fraction c → D)

$$D(z) = \frac{H}{z[1 - (1/z) - \epsilon_p/(1-z)]^2}.$$

# Hadronic transport coefficients



- based on unitarized chiral HQ model

[Abreu et al, Ann. Phys. **326**, 2737 (2011); PRD **87**, 034019 (2013); Tolos, Torres-Rincon, PRD **88**, 074019 (2013)]

- coupled channel T-matrix approach with pseudo-Goldstone mesons ( $\pi, \eta, K$ ) and baryons ( $N, \bar{N}, \Delta, \bar{\Delta}$ )
- D-meson scattering cross sections used for HQ drag and diffusion coefficients



# Hadronic transport coefficients

- drag and diffusion coefficients for D from mesonic interactions





# Hadronic transport coefficients

- drag and diffusion coefficients for D from baryonic interactions





# Hadronic transport coefficients

- drag and diffusion coefficients for D at finite  $\mu_B$



# D-mesons at FAIR

- $E_{\text{kin}} = 25 \text{ AGeV}$
- bulk-medium evolution: UrQMD-hydro hybrid vs. UrQMD coarse graining
- QQP HQ transport coefficients from resonance model
- influence of hadronization description: sensitivity to
  - hadronization temperature
  - coalescence vs. fragmentation (dependence on  $r_D$ ,  $\epsilon_p$ )





- strong rise in  $R_{AA}$  due to energy constraint in pp ( $p_T \lesssim 2.5 \text{ GeV}$ )

[Inghrami, HvH, Endres, Torres-Rincon, Bleicher, Eur. Phys. J. 79:52 (2019)]

# D-mesons at FAIR

- $E_{\text{kin}} = 25 \text{ AGeV}$
- bulk-medium evolution: UrQMD-hydro hybrid vs. UrQMD coarse graining
- QQP HQ transport coefficients from resonance model
- influence of hadronization description: sensitivity to
  - hadronization temperature
  - coalescence vs. fragmentation (dependence on  $r_D$ ,  $\epsilon_p$ )





- strong rise in  $R_{AA}$  due to energy constraint in pp ( $p_T \lesssim 2.5 \text{ GeV}$ )

[Inghrami, HvH, Endres, Torres-Rincon, Bleicher, Eur. Phys. J. 79:52 (2019)]

# D-mesons at FAIR



- $E_{\text{kin}} = 25 \text{ AGeV}$
- bulk-medium evolution: UrQMD-hydro hybrid vs. UrQMD coarse graining
- QGP HQ transport coefficients from resonance model
- influence of hadronization description: sensitivity to
  - hadronization temperature
  - coalescence vs. fragmentation (dependence on  $r_D$ ,  $\epsilon_P$ )



- [Inghirami, HvH, Endres, Torres-Rincon, Bleicher, Eur. Phys. J. 79:52 (2019)]

# D-mesons at FAIR

- $E_{\text{kin}} = 25 \text{AGeV}$
- bulk-medium evolution: UrQMD-hydro hybrid vs. UrQMD coarse graining
- QQP HQ transport coefficients from resonance model
- influence of hadronization description: sensitivity to
  - hadronization temperature
  - coalescence vs. fragmentation (dependence on  $r_D, \epsilon_p$ )





- influence of longer hadronic phase negligible

[Inghirami, HvH, Endres, Torres-Rincon, Bleicher, Eur. Phys. J. 79:52 (2019)]

# D-mesons at FAIR

- $E_{\text{kin}} = 25 \text{AGeV}$
- bulk-medium evolution: UrQMD-hydro hybrid vs. UrQMD coarse graining
- QQP HQ transport coefficients from resonance model
- influence of hadronization description: sensitivity to
  - hadronization temperature
  - coalescence vs. fragmentation (dependence on  $r_D$ ,  $\epsilon_p$ )



- large influence on hadronization procedure

[Inghirami, HvH, Endres, Torres-Rincon, Bleicher, Eur. Phys. J. 79:52 (2019)]